

The tandem catalytic asymmetric allene diboration/imine allylation and the asymmetric transition-metal-catalyzed conjugate allylation of activated enones

Author: Joshua Daniel Sieber

Persistent link: <http://hdl.handle.net/2345/355>

This work is posted on [eScholarship@BC](#),
Boston College University Libraries.

Boston College Electronic Thesis or Dissertation, 2008

Copyright is held by the author, with all rights reserved, unless otherwise noted.

Boston College

The Graduate School of Arts and Sciences

Department of Chemistry

THE TANDEM CATALYTIC ASYMMETRIC ALLENE DIBORATION/IMINE
ALLYLATION AND THE ASYMMETRIC TRANSITION-METAL-CATALYZED
CONJUGATE ALLYLATION OF ACTIVATED ENONES

a dissertation

by

JOSHUA DANIEL SIEBER

Submitted in partial fulfillment of the requirements

for the degree of

Doctor of Philosophy

August 2008

ABSTRACT

JOSHUA D. SIEBER: The Tandem Catalytic Asymmetric Allene Diboration/Imine Allylation and the Asymmetric Transition-Metal-Catalyzed Conjugate Allylation of Activated Enones

(Under the direction of: Professor James P. Morken)

Described herein are methods for asymmetric allylation. Chapter 1 describes the scope of the Pd-catalyzed asymmetric diboration of prochiral allenes. The products of this process possess both a chiral allylboronate functional group and a vinylboronate moiety. The allylboronate functionality can subsequently be used for imine allylation, without isolation of the diboron intermediate, resulting in the formation of atypical allylation products through a tandem, one-pot sequence. Furthermore, enantioselection in the catalytic diboration and chirality transfer in the subsequent imine allylation are high; thus, non-racemic, protected homoallylic amines, and other derivatives, are produced in high enantiomeric excess. Chapter 2 describes the discovery and development of a transition-metal-catalyzed asymmetric conjugate allylation of allylboronate ester nucleophiles to activated enones. The scope, utility, and mechanistic aspects of this new reaction are discussed.

Dedicated to my loving family:

Richard J. Sieber
Mary E. Sieber
Tammy S. Sieber
Richard W. Sieber, D.P.M.
Rachel E. Sieber
Chumly M. M. Sieber

Also to:

Professor Stephen C. Hoops
for inspirational instruction in organic chemistry

ACKNOWLEDGMENTS

I am deeply grateful to all the excellent instructors I have been taught and influenced by during all of my educational training. As such, the work described in this dissertation represents a tiny portion of the “*fruits of their labor*.” I have been extremely blessed to have known them all.

In relation to this work, I thank my research advisor Prof. James Morken. It has been my pleasure to conduct research under an extremely intelligent and thoughtful chemist. In my experience, he has always been available to discuss problems and concerns related to research with an extremely positive attitude. As a result, I have many times come away from these interactions encouraged that the chemical problems I’ve been facing will be solved through new insights that Prof. Morken has given me. During my time in graduate school, Prof. Morken has taught me how to think about difficult problems as an independent scientist. I thank him for this. I am confident that much more insightful and elegant research will continue to flow from the Morken lab, and I am thankful for playing a small role in Prof. Morken’s independent career.

During my time in Prof. Morken’s lab, I have come into contact with many bright graduate students and post-doctoral fellows. All of these people have given me new theoretical and practical insights into the field of chemistry. I thank Prof. Jeremy Morgan who was a fourth year graduate student when I joined the lab. He showed me the ropes when I started and was always available for advice on how to conduct experiments. I also thank Dr. Nathan Fuller and Dr. Mike Evans (also fourth year graduate students at the time) for similar advice. Dr. Wesley Moran was a post-doc who worked in the same room as me for a while. He taught me many valuable lessons in chemistry and also made a fun environment to work in. I thank him for this. I also thank Dr. Sundaram Shanmugham, another post-doc, for many insightful discussions. Furthermore, I would like to thank Nick Pelz, Angela Woodward, and Heather Burks who all worked on the allene diboration project. I also thank Nick for providing an enjoyable working environment. He was a fun person to be around. In addition, I thank Heather for the good lunches we had at Sutton’s Drug Store. I also thank Rebecca Dueñes for her friendship and for helpful insights. In addition, I thank Laura Brozek and Laura Kliman for providing an enjoyable work environment while at Boston College. Finally, I thank the following Morken group members for their insights: Dr. Eric Ballard, Dr. Pavan Kumar, Dr. Stéphane Trudeau, Joe Nepveux, Diane Kalendra, Andy Ommen, Mike Tarselli, Li Yao, Ping Zhang, Hee Yeon Cho, Rob Ely, and Bob Kyne.

I thank the University of North Carolina at Chapel Hill for the opportunity to initiate my graduate studies there. I thank Prof. Maurice Brookhart for excellent instruction in physical organic chemistry and for supporting me by providing recommendation letters for post-doctoral applications. I also thank Prof. Jeffrey Johnson for excellent instruction in organic chemistry and for providing me with recommendation letters. In addition, I would like to thank Prof. Michel Gagné, Prof. Michael Crimmins, and Prof. Joseph DeSimone for their instruction. Finally, I thank Dr. Peter White for X-ray crystallographic analyses, and Dr. Shubin Liu for performing density functional theory (DFT) calculations pertinent to my research.

Further, I wish to thank the Boston College Chemistry Department for making the move from UNC to BC seamless. The research labs here are very helpful, and I thank all members of the department for making an environment that is conducive to learning. I thank my defense committee: Prof. Amir Hoveyda, Prof. T. Ross Kelly, and Prof. Kian Tan for their work in perfecting this dissertation. I also would like to thank Dr. John Boylan for always being very helpful in regard to the use of the NMR spectrometers here at BC. His help was crucial for the Bruker to Varian switch that I encountered upon moving to BC. His insights were extremely helpful and necessary for completing my research. I also thank Boston College for awarding me the LaMattina Graduate Fellowship in Organic Chemistry. In addition, I thank the Organic Division of the American Chemical Society for awarding me with a fellowship, and I thank Organic Syntheses, Inc. and Nelson J. Leonard for funding this award. I also thank Schering-Plough for awarding me the Schering-Plough Science and Innovation Award in 2008.

I wish also to thank many of the Professors who taught me at my undergraduate institution: The Pennsylvania State University. These people gave me a strong foundation in the physical sciences, and I am indebted to them for this. I thank Prof. Ayusman Sen for giving me the opportunity to conduct undergraduate research in his laboratory. The techniques which I learned in his group related to air and moisture sensitive chemistry have been extremely valuable in my graduate studies. I also thank Dr. Cecily Andes for instructing me while working in Prof. Sen's lab. Furthermore, I thank Prof. John Lowe, Prof. Kenneth Feldman, Prof. Kenneth Merz, Jr., Prof. Robert Minard, Prof. Christopher Falzone, Prof. William Hamilton, Prof. Fred Ridener, Prof. David Wells, Prof. Javier Gomez-Calderon, and Prof. Kwang Wang for excellent instruction. I also thank Prof. Stephen C. Hoops who introduced me to the field of organic chemistry. He is an amazing teacher, and I am indebted to him for the knowledge I obtained from his classes.

I thank my loving family for all their encouragement and support. My whole being has been shaped by them. I thank my grandparents Richard and Mary Sieber. They have been an excellent model for me to pattern my own life after. Words cannot express the impact they have had on shaping my life. I thank them for instilling in me a strong work ethic and for many enjoyable times. I have especially enjoyed the times of hunting, fishing, and working around the house. I also thank my mother, Tammy Sieber, for her love and support over the years. I thank my siblings Rich, Rachel, and Chumly for all the fun times. You all have made this work possible through your love and support; Thank You!

Finally, I thank the one, true, and living God who sent His son, Jesus Christ, to die on a cross and be raised from the dead to bear the sins of ALL mankind. He is alive today and through Him ALL men are saved. God's gift of salvation is for anyone who will receive it and believe in Him. I thank God for this wonderful gift and for all the blessings He has bestowed on me and my family. May the rest of my life be a testament to Him and bring glory to His name.

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	vii
LIST OF SCHEMES.....	viii
LIST OF FIGURES.....	xii
LIST OF ABBREVIATIONS.....	xiii
Chapter	
1. The Pd-Catalyzed Enantioselective Diboration of Allenes and its Application to the Allylation of Imines.....	1
I. Introduction.....	1
II. Background.....	3
A. Diastereoselective Imine Allylation using Substituted Allylmetals.....	3
B. Substrate Control for Diastereoselective Imine Allylation.....	7
C. Diastereoselective Imine Allylation Using Chiral Auxiliaries.....	11
D. Use of Chiral Allylmetals for Diastereoselective Imine Allylation.....	15
E. Catalytic Asymmetric Imine Allylation.....	20
F. Catalytic Diboration of Allenes.....	26
III. Development of the Pd-Catalyzed Enantioselective Diboration of Allenes and its Application to the Allylation of Imines.....	29
A. The Pd-Catalyzed Enantioselective Diboration of Allenes.....	29
B. Allylation of Imines with the Products of Allene Diboration.....	33

C. Scope of the Tandem Allene Diboration/Imine Allylation and Other Transformations.....	38
D. Stereochemical Analysis in the Imine Allylation.....	43
IV. Conclusion.....	46
V. Experimental Procedures.....	47
A. General.....	47
B. Experimental Procedures.....	49
2. Development of the Transition-Metal-Catalyzed Enantioselective Conjugate Allylation.....	101
I. Introduction.....	101
II. Background.....	103
A. Non-Enantioselective Conjugate Allylation Processes.....	103
B. Asymmetric Conjugate Allylation Processes.....	107
C. Conjugate Addition Reactions Catalyzed by Group 10 Metals.....	110
III. Discovery and Development of a Transition-Metal-Catalyzed Conjugate Allylation with Allylboronate Ester Nucleophiles.....	113
A. Reaction Discovery.....	113
B. Reaction Development.....	115
C. Scope and Utility of the Ni-Catalyzed Conjugate Allylation.....	117
D. Mechanistic Aspects of the Ni-Catalyzed Conjugate Allylation.....	121
IV. Development of a Transition-Metal-Catalyzed Asymmetric Conjugate Allylation with Allylboronate Ester Nucleophiles.....	133

A. Discovery of a Competent Chiral Catalyst.....	133
B. Scope and Utility in the Ni-Catalyzed Asymmetric Conjugate Allylation.....	137
C. Mechanistic Aspects of the Ni-Catalyzed Asymmetric Conjugate Allylation.....	143
D. Model for Chemoselectivity in the Conjugate Allylation.....	148
E. Extension to an Asymmetric Chemoselective Alkylidene Conjugate Allylation.....	152
F. Other Related Reactions.....	164
V. Conclusions.....	169
VI. Experimental Procedures.....	171
A. General.....	171
B. Experimental Procedures.....	174
1. Weinreb Amide Syntheses.....	174
2. Dialkylidene Ketone Syntheses.....	178
3. Chiral Ligand Syntheses.....	196
4. Allylation Procedures.....	237
5. Mechanistic Experiments.....	325
6. Synthetic Transformations.....	358

LIST OF TABLES

Table 1.1: Crotylation of Imines.....	4
Table 1.2: Diastereoselective Addition of Allylmetals to α -Chiral Imines.....	8
Table 1.3: Substrate Scope in the Asymmetric Pd-Catalyzed Allene Diboration.....	31
Table 1.4: Tandem Allene Diboration/Imine Allylation of Silylimines.....	39
Table 1.5: Tandem Diboration/Imine Allylation of <i>in situ</i> Prepared Imines.....	40
Table 2.1: Catalyst Optimization for the Conjugate Allylation of 2.62	116
Table 2.2: Synthesis of Unsymmetrical Dialkylidene Ketones.....	118
Table 2.3: Substrate Scope in the Ni-Catalyzed Conjugate Allylation.....	120
Table 2.4: Survey of Chiral Ligands in the Conjugate Allylation.....	134
Table 2.5: Optimization of the Activating Alkene in the DAK Substrate.....	137
Table 2.6: Synthesis of Unsymmetrical Dialkylidene Ketones.....	138
Table 2.7: Substrate Scope in the Asymmetric Conjugate Allylation.....	140
Table 2.8: Effect of Metal:Ligand Ratio in the Conjugate Allylation.....	147
Table 2.9: Correlation Between Ligand Basicity and Chemoselectivity.....	149
Table 2.10: Chiral Catalyst Survey with Pd ₂ (dba) ₃ as Precatalyst.....	153
Table 2.11: Chiral Catalyst Survey with 2.156 as Substrate.....	155
Table 2.12: Chiral Catalyst Survey with Newly Prepared Phosphoramidites.....	157
Table 2.13: Solvent Effects in the Asymmetric Conjugate Allylation of 2.156	158
Table 2.14: Attempts at Formation of All-Carbon Quaternary Stereocenters.....	160

LIST OF SCHEMES

Scheme 1.1: Stereoselective Addition of Allylic Metal Reagents to Imines.....	2
Scheme 1.2: A New Allylmetal Reagent for Imine Allylation.....	3
Scheme 1.3: Rational for Diastereoselectivity in the Crotylation of Imines.....	5
Scheme 1.4: Diastereoselective Crotylsilation.....	6
Scheme 1.5: Crotylboration of <i>in situ</i> Prepared Imines.....	7
Scheme 1.6: Allylmetal Additions to α -Heteroatom α -Chiral Imines.....	10
Scheme 1.7: 1,3-Asymmetric Induction in the Allylation of Imines.....	11
Scheme 1.8: α -Methylbenzylamine as a Chiral Auxiliary for Imine Allylation.....	12
Scheme 1.9: Aminoacid Based Auxiliaries for Imine Allylation.....	13
Scheme 1.10: Use of SAMP Hydrazones for Imine Allylation.....	14
Scheme 1.11: Allylation of Chiral Sulfinylketimines.....	15
Scheme 1.12: Chiral Allylic Silanes and Stannanes for Diastereoselective Imine Allylation.....	16
Scheme 1.13: Asymmetric Imine Allylation with Chiral Allylboron Reagents.....	17
Scheme 1.14: Asymmetric Allylation of Ketimines with Allylborane 1.75	18
Scheme 1.15: Use of Chiral Allylic Silanes for Asymmetric Imine Allylation and Crotylation.....	19
Scheme 1.16: Chiral Additives for Asymmetric Imine Allylation and Crotylation.....	20
Scheme 1.17: Enantioselective Pd-Catalyzed Allyl Additions to Imines.....	21
Scheme 1.18: Enantioselective Cu-Catalyzed Allyl Additions to Imines.....	23
Scheme 1.19: Substituted Chiral BINOL-Derived Catalysts for Imine Allylation.....	24

Scheme 1.20: Organocatalytic Asymmetric Addition of Allyl and Crotylboronates.....	26
Scheme 1.21: Pt-Catalyzed Diboration of Allenes.....	27
Scheme 1.22: Miyaura's Proposed Mechanism for the Pt-Catalyzed Diboration of Allenes.....	28
Scheme 1.23: Allylation of Benzaldehyde with a Product of Allene Diboration.....	28
Scheme 1.24: Enantiomeric Excess Determination of (<i>S</i>)- 1.122	32
Scheme 1.25: Hydrogenation of (<i>S</i>)- 1.132 with D ₂	33
Scheme 1.26: Diimide Reduction of (<i>S</i>)- 1.122	33
Scheme 1.27: Initial Attempts at Imine Allylation with 1.132	35
Scheme 1.28: Tandem Allene Diboration/Imine Allylation.....	36
Scheme 1.29: Allylation of <i>in situ</i> Prepared Imines with 1.132	38
Scheme 1.30: Access to Other Homoallylic Amides.....	42
Scheme 1.31: Isolation of the Allylation Intermediate.....	42
Scheme 2.1: General Metal-Catalyzed Conjugate Addition.....	101
Scheme 2.2: Reaction of Di(allyl)cuprate with Enones.....	103
Scheme 2.3: Conjugate Allylation with Allylsilanes.....	104
Scheme 2.4: Conjugate Allylation with Allyltrimethylstannane.....	104
Scheme 2.5: Effect of Cu-Salt Purity in Conjugate Allylation.....	106
Scheme 2.6: Use of Allylic Copper Reagents in Conjugate Allylation.....	107
Scheme 2.7: Asymmetric Sakurai Conjugate Allylation.....	108
Scheme 2.8: Asymmetric Conjugate Allylation with Allylic Stannanes and Allylmagnesium Chloride.....	109

Scheme 2.9: Ni-Catalyzed Conjugate Addition of AlMe ₃ to Enones.....	110
Scheme 2.10: Reaction of L ₂ Pd(enone) Complexes with Lewis Acids.....	111
Scheme 2.11: Mechanism of the Pd-Catalyzed Conjugate Addition of AlMe ₃	112
Scheme 2.12: Conjugate Allylation with Allene Diboration Products.....	114
Scheme 2.13: Pd-Catalyzed Conjugate Allylation of Dbal.....	115
Scheme 2.14: Transformations of the Benzyldiene Activating Group.....	121
Scheme 2.15: Proposed Catalytic Cycle for the Conjugate Allylation.....	122
Scheme 2.16: Trap of the Intermediate Boron Enolate.....	123
Scheme 2.17: Calculated Reductive Elimination Barriers in the Conjugate Allylation.	125
Scheme 2.18: Analysis of a Phosphino-Directing Dialkylidene Ketone.....	127
Scheme 2.19: Conjugate Allylation of a Phosphine Containing DAK.....	128
Scheme 2.20: Synthesis of (PPh ₃)Ni(2.96) Complex.....	129
Scheme 2.21: Deuterium Labeling Experiments.....	131
Scheme 2.22: Possible Explanation for Labeling Experiments.....	131
Scheme 2.23: Cleavage of the Alkylidene Activating Unit.....	141
Scheme 2.24: Synthesis of a Potent NHE Inhibitor.....	143
Scheme 2.25: Deuterium Labeling Experiments.....	148
Scheme 2.26: Model for Chemoselectivity in the Conjugate Allylation.....	151
Scheme 2.27: Reaction of 2.164 with Ni(cod) ₂ /PPh ₃	161
Scheme 2.28: Attempts at an Asymmetric Conjugate Allylation of 2.169	162
Scheme 2.29: Possible Mode of Asymmetric Induction in the Conjugate Allylation...	163
Scheme 2.30: Use of an α,β -Unsaturated α -Ketoesters.....	165

Scheme 2.31: Enantioselective Allylation of 2.173	166
Scheme 2.32: Catalytic Crotylation of 2.173	167
Scheme 2.33: Catalytic Addition of a Diboronate Esters to 2.173	168
Scheme 2.34: Catalytic Allylation of an $\alpha,\beta,\gamma,\delta$ -Unsaturated Enone.....	169

LIST OF FIGURES

Figure 1.1: Transition State Models for Allylation of α -Chiral Imines and Aldehydes....	9
Figure 1.2: Structure of (TADDOL)MONOPHOS.....	30
Figure 1.3: Crystal Structure of 1.145	43
Figure 1.4: Transition Structures for Allylation of (<i>E</i>)-Aldimines.....	44
Figure 1.5: Possible Transition Structures in the Imine Allylation.....	45
Figure 1.6: Crystal Structure of 1.161	89
Figure 1.7: Crystal Structure of 1.166	94
Figure 2.1: Electrophiles Tested in the Pd-Catalyzed Conjugate Allylation.....	115
Figure 2.2: Calculated Reductive Elimination Barriers for Bis(allyl)Pd-Complexes....	124
Figure 2.3: Calculated Transition Structure for the Reductive Elimination from 2.92	126
Figure 2.4: Proposed [(PCy ₃)Ni(dba)] ₂ Dimer.....	132
Figure 2.5: Chiral Ligands Used in the Chiral Catalyst Survey.....	135
Figure 2.6: Effect of Ligand Enantiomeric Purity on Reaction Selectivity.....	144
Figure 2.7: Effect of Ligand Enantiomeric Purity on Reaction Rate.....	145
Figure 2.8: Proposed Dimeric Complexes in the Asymmetric Conjugate Allylation....	146
Figure 2.9: New Ligands for Asymmetric Alkylidene Allylation.....	156

LIST OF ABBREVIATIONS

Å	angstrom
Ac ₂ O	acetic anhydride
AcOH	acetic acid
BBN	borabicyclo[3.3.1]nonane
BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthalene
BINOL	1,1'-binaphthalene-2,2'-diol
(Boc) ₂ O	di- <i>t</i> -butyl dicarbonate
B ₂ (pin) ₂	bis(pinacolato)diboron
brsm	based on recovered starting material
<i>n</i> -BuLi	<i>n</i> -butyllithium
<i>t</i> -BuLi	<i>t</i> -butyllithium
CDI	carbonyl diimidazole
cee	conservation of enantiomeric excess
cod	1,5-cyclooctadiene
COSY	¹ H- ¹ H correlation spectroscopy
DAK	dialkylidene ketone
dba	dibenzylideneacetone
<i>o</i> -DCB	1,2-dichlorobenzene
DCC	<i>N,N</i> -dicyclohexylcarbodiimide
DIBAL	diisobutylaluminum hydride
DMAP	4-(dimethylamino)pyridine

DMF	<i>N,N</i> -dimethylformamide
dppf	bis(diphenylphosphino)ferrocene
dr	diastereomeric ratio
EDCI	<i>N</i> -(3-dimethylaminopropyl)- <i>N</i> '-ethylcarbodiimide hydrochloride
ee	enantiomeric excess
eq	equation
equiv	equivalents
EtOAc	ethyl acetate
EtOH	ethanol
GLC	gas-liquid chromatography
HPLC	high performance liquid chromatography
HSQC	heteronuclear single-quantum correlation spectroscopy
LAH	lithium aluminum hydride
MeOH	methanol
MS	molecular sieves
n/a	not applicable
NBSH	<i>o</i> -nitrobenzenesulfonylhydrazine
n.d.	not determined
Ni(cod) ₂	bis(1,5-cyclooctadiene)nickel
NMR	nuclear magnetic resonance spectroscopy
nOe	nuclear Overhauser effect
NOESY	nuclear Overhauser effect spectroscopy

PCy ₃	tricyclohexylphosphine
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium
pin	pinacol
psig	pounds per square inch gauge
pyr	pyridine
RAMP	(<i>R</i>)-1-amino-2-(methoxymethyl)pyrrolidine
RBF	round-bottom flask
RCM	ring-closing metathesis
rt	room temperature
SAMP	(<i>S</i>)-1-amino-2-(methoxymethyl)pyrrolidine
SFC	supercritical fluid chromatography
TBAF	tetrabutylammonium fluoride
TBS	<i>t</i> -butyldimethylsilyl
TEA	triethylamine
THF	tetrahydrofuran
TLC	thin-layer chromatography
TsOH	<i>p</i> -toluenesulfonic acid monohydrate

Chapter 1


The Pd-Catalyzed Enantioselective Diboration of Allenes and its Application to the Allylation of Imines

I. Introduction

Optically active chiral amines are commonly found in natural products and commercial drugs. In addition, amine functional groups have found widespread use in chiral ligands for asymmetric transition-metal-catalyzed processes. Therefore, methods to produce chiral amines in enantiomerically enriched form are highly desirable. One of the most useful ways to create a stereogenic carbon atom with an attached nitrogen atom starting from simple prochiral starting materials is by the addition of an allylmetal reagent to an imine electrophile (Scheme 1.1).¹ This procedure affords chiral homoallylic amines, wherein facial selectivity can be dictated using substrate or reagent control. Furthermore, use of appropriately substituted allylmetal reagents simultaneously creates two new stereocenters. These methods represent powerful tools for the efficient construction of chiral amines.

¹ For recent reviews, see: (a) Ding, H.; Friestad, G. K. *Synthesis* **2005**, 2815. (b) Friestad, G. K.; Mathies, A. K. *Tetrahedron* **2007**, 63, 2541.

Scheme 1.1: Stereoselective Addition of Allylic Metal Reagents to Imines




With the goal of developing synthetically-useful catalytic bis(metalations) of unsaturated carbon-carbon bonds,² we envisioned that a bis(metalation) of the internal olefin of an allene would provide a practical allylmetal reagent for imine allylation (Scheme 1.2). For the overall process to be synthetically valuable, the catalyst for the bis(metalation) must afford the product in high enantioselectivity, and chirality transfer in the imine addition reaction must be high. If these requirements could be met, the imine allylation product resulting from this sequence would contain a vinylic carbon-metal bond, which may be converted into a wide variety of useful organic synthons with high enantiomeric purity. To accomplish these goals, we chose to study the addition of bis(boronate) esters to allenes. The high stability of organoboronate esters makes them attractive;^{2e} however, no catalytic asymmetric diboration of prochiral allenes was known prior to our work.³

² For reviews on metal catalyzed bis(metalations) of unsaturated carbon-carbon bonds, see: (a) Suginome, M.; Ito, Y. *Chem. Rev.* **2000**, *100*, 3221. (b) Marder, T. B.; Norman, N. C. *Topics in Catalysis* **1998**, *5*, 63. (c) Ishiyama, T.; Miyaura, N. *J. Organomet. Chem.* **2000**, *611*, 392. (d) Beletskaya, I. *Chem. Rev.* **2006**, *106*, 2320. (e) Burks, H. E.; Morken, J. P. *Chem. Commun.* **2007**, 4717.

³ (a) Pelz, N. F.; Woodward, A. R.; Burks, H. E.; Sieber, J. D.; Morken, J. P. *J. Am. Chem. Soc.* **2004**, *126*, 16328. (b) Burks, H. E.; Liu, S.; Morken, J. P. *J. Am. Chem. Soc.* **2007**, *129*, 8766.

Scheme 1.2: A New Allylmetal Reagent for Imine Allylation



II. Background

A. Diastereoselective Imine Allylation Using Substituted Allylmetals

The addition of crotylmetal reagents to imines was initially investigated by Yamamoto.^{4,5} A selection of these results is shown in Table 1.1, where it can be seen that selectivity in the crotylation is highly dependent on the choice of the metal used in the reaction. When Li, Mg, or Sn reagents are employed, the *syn* diastereomer is afforded in moderate diastereoselectivities. However, the use of crotylboron reagents (entry 4) gave exclusively the *anti* product. Interestingly, in the case of boron, the nature of the imine dictates which diastereomer is formed in the reaction. For both aryl and linear-aliphatic imines bearing small substituents on nitrogen ($R' = \text{linear and alkyl}$), the *syn*-product is favored in a 75:25 to 85:15 ratio. For linear-aliphatic imines where the substituent on nitrogen is branched ($R' = i\text{-Pr}$), complete *syn* selectivity results; however, the analogous aryl imines slightly favor the *anti*-allylation product (65:35 *anti:syn* when

⁴ Yamamoto, Y.; Komatsu, T.; Maruyama, K. *J. Am. Chem. Soc.* **1985**, *50*, 3115.

⁵ For a review of allylmetal addition reactions, see: Yamamoto, Y.; Asao, N. *Chem. Rev.* **1993**, *93*, 2207.

R = Ph). Furthermore, when branched aliphatic imines are used (R = *i*-Pr), moderate *anti* selectivity is again observed (~30:70).

Table 1.1: Crotylation of Imines


entry	R	R'	M	1.3:1.4 (<i>syn:anti</i>)	% yield
1	Ph	Ph	Li	60:40	50
2	Ph	Ph	MgCl	74:26	95
3	Ph	Ph	9-BBN	0:100	93
4	Ph	Ph	SnBu ₃ /BF ₃	75:25	80

Diastereoselectivity in these reaction can be explained by Zimmerman-Traxler⁶ chair-like transition states in the cases where M = Li, Mg, or B (Scheme 1.3). Chair-like transition structure **1.5** is preferred over the boat-like transition structure **1.6**.^{4,5} With allylboranes, the variation in selectivity with substrate structure is dependent more on the steric bulk of the group on nitrogen than on the size of the carbon substituent. That is to say, as the size of R' increases, a penalizing 1,3-diaxial interaction between this group and the bridgehead proton of the 9-BBN ring destabilizes **1.5** relative to **1.6**. This phenomenon is not observed with other metals because the weaker interaction between the lone pair on nitrogen and the metal alleviates the 1,3-interaction. In the case of

⁶ Zimmerman, H. E.; Traxler, M. D. *J. Am. Chem. Soc.* **1957**, 79, 1920.

crotylstannanes, these reagents react through open transition states and addition of a Lewis acid is required for reaction. Transition structure **1.7** is responsible for formation of the *syn* product in the crotylstannylation of imines.


Scheme 1.3: Rational for Diastereoselectivity in the Crotylation of Imines



Kobayashi⁷ has used crotylsilanes for diastereoselective crotylation of benzoylhydrazones (**1.9**, Scheme 1.4). Both aromatic and aliphatic hydrazones react under the optimized procedure with good diastereoselectivities. Importantly, the use of the *E*- or *Z*-crotylsilanes provides the *anti* or *syn* diastereomer of product, respectively. Thus, either diastereomer of product can be conveniently prepared by the correct choice of crotylsilane.

⁷ Kobayashi, S.; Hirabayashi, R. *J. Am. Chem. Soc.* **1999**, *121*, 6942.


Scheme 1.4: Diastereoselective Crotylsilation



Kobayashi has developed a novel allylation where the imine is formed *in situ* from an aldehyde and an ammonia source.⁸ Notably, this was done using allylboronate esters as the nucleophile in the crotylation reaction (Scheme 1.5). The use of crotylboronates (*E*)-**1.14** and (*Z*)-**1.14** affords amines **1.15** and **1.16**, respectively, in a stereospecific manner. In these cases, the ammonia derived imine (the condensation product between benzaldehyde and ammonia) is allylated in preference to benzaldehyde that may be present in solution due to the equilibrium between aldehyde and imine; thus, the homoallylic amine product is formed directly.

⁸ Sugiura, M.; Hirano, K.; Kobayashi, S. *J. Am. Chem. Soc.* **2004**, *126*, 7182.

Scheme 1.5: Crotylboration of *in situ* Prepared Imines



B. Substrate Control for Diastereoselective Imine Allylation

The study of diastereoselective additions of organometallic reagents to α -chiral aldehydes is arguably one of the earliest contributions to the chemical literature which led to the construction of useful models for the prediction of acyclic stereocontrol.⁹ Diastereocontrol in the addition of allylmetal reagents to α -chiral imines was first explored by Yamamoto.¹⁰ As depicted in Table 1.2, imine **1.17** is allylated with a variety of allylmetal reagents in good to excellent diastereoselectivity. The Cram-addition^{9a} product is the major diastereomer isolated in these reactions. Notably, the allylation of imines is generally more selective than the analogous allylation of α -chiral aldehydes.

⁹ (a) Cram, D. J.; Elhafez, F. A. A. *J. Am. Chem. Soc.* **1952**, *74*, 5828. (b) Karabatsos, G. J. *J. Am. Chem. Soc.* **1967**, *89*, 1367. (c) Chérest, M.; Felkin, H.; Prudent, N. *Tetrahedron Lett.* **1968**, *9*, 2199. (d) Anh, N. T.; Eisenstein, O. *Nouv. J. Chim.* **1977**, *1*, 61. (e) Paddon-Row, M. N.; Rondan, N. G.; Houk, K. N. *J. Am. Chem. Soc.* **1982**, *104*, 7162. (f) Houk, K. A.; Paddon-Row, M. N.; Rondan, N. G.; Wu, Y. -D.; Brown, F. K.; Spellmeyer, D. C.; Metz, J. T.; Li, Y.; Loncharich, R. J. *Science* **1986**, *231*, 1108. (g) Wong, S. S.; Paddon-Row, M. N. *J. Chem. Soc. Chem. Commun.* **1990**, 456. (h) Wong, S. S.; Paddon-Row, M. N. *J. Chem. Soc. Chem. Commun.* **1991**, 327. (i) Wu, Y. -D.; Tucker, J. D.; Houk, K. N. *J. Am. Chem. Soc.* **1991**, *113*, 5018. (j) Frenking, G.; Köhler, K. F.; Reetz, M. T. *Tetrahedron* **1991**, *47*, 8991. (k) Reetz, M. T. *Acc. Chem. Res.* **1993**, *26*, 462 and references therein.


¹⁰ (a) Yamamoto, Y.; Komatsu, T.; Maruyama, K. *J. Am. Chem. Soc.* **1984**, *106*, 5031. (b) Yamamoto, Y.; Nishii, S.; Maruyama, K.; Komatsu, T.; Ito, W. *J. Am. Chem. Soc.* **1986**, *108*, 7778.

This phenomenon results from the fact that the α -stereocenter of an α -chiral imine occupies the axial position in the cyclic chair-like transition state of the allylation (Figure 1.1, **1.21** and **1.22**); the chair preference arises because of the *E*-configuration of the imine. The allylmethyl reagent preferentially adds to the imine through **1.22** to avoid the penalizing steric interaction that is present between R_m and the coordination sphere of the allylmethyl in transition structure **1.21**. In comparison, since the α -stereocenter of an α -chiral aldehyde occupies the equatorial position in transition structure **1.23**, steric interactions between the α -chiral stereocenter of the aldehyde and the ligands of the allylmethyl are not present. Thus, lower diastereoselectivities are typically observed.⁵

Table 1.2: Diastereoselective Addition of Allylmethyls to α -Chiral Imines

entry	R'	M	1.19:1.20 (Cram:anti-Cram)
1	<i>n</i> -Pr	9-BBN	96:4
2	<i>i</i> -Pr	9-BBN	100:0
3	<i>n</i> -Pr	SnBu ₃ /TiCl ₄	93:7
4	<i>i</i> -Pr	SnBu ₃ /TiCl ₄	92:8
5	<i>n</i> -Pr	MgCl	84:16
6	<i>n</i> -Pr	MgBr	68:32
7	<i>i</i> -Pr	MgCl	70:30

Figure 1.1: Transition State Models for Allylation of α -Chiral Imines and Aldehydes



Chiral imines bearing α -heteroatoms undergo allylation under either Felkin^{9c} or chelation control, depending on the allylmethyl reagent used. For example, when imine **1.24** is subjected to allylmagnesium bromide, the major *syn* product results, as predicted by chelation control (Scheme 1.6, eq 1).¹¹ The use of allyl-9-BBN, however, results in formation of the *anti* diastereomer, as predicted by Felkin control (the benzyl ether being the large group). Furthermore, imines **1.27**¹² and **1.30**¹³ undergo Felkin-controlled allylation when allylic Grignard reagents are used (Scheme 1.6, eq 2 and 3). In addition, when hydrazones are employed as electrophiles, allylsilanes can be used as nucleophiles and excellent diastereoselectivity results (Scheme 1.6, eq 4 and 5).¹⁴


¹¹ Badorrey, R.; Cativiela, C.; Díaz-de-Villegas, M. D.; Díez, R.; Galvez, J. A. *Eur. J. Org. Chem.* **2002**, 3763.

¹² Cainelli, G.; Giacomini, D.; Galletti, P.; Quintavalla, A. *Eur. J. Org. Chem.* **2002**, 3153.

¹³ Reetz, M. T.; Jaeger, R.; Drewlies, R.; Hübel, M. *Angew. Chem. Int. Ed.* **1991**, 30, 103.

¹⁴ Kobayashi, S.; Hirabayashi, R. *J. Am. Chem. Soc.* **1999**, 121, 6942.

Scheme 1.6: Allylmetal Additions to α -Heteroatom α -Chiral Imines




Alkoxy substitution at the β -position has also been shown to control the stereochemistry of allylmetal additions.¹⁵ Allylation of chiral imine **1.39** with allylmagnesium chloride or allyltributylstannane gave the chelation control addition product (Scheme 1.7). When allylmagnesium chloride is used, the addition of CeCl₃ increases the diastereoselectivity from 67:33 to 89:11 *anti:syn*. Formation of the *anti*

¹⁵ Shimizu, M.; Morita, A.; Fujisawa, T. *Chem. Lett.* **1998**, 467.

product is believed to come from external addition of the allyl nucleophile to the chelated substrate. In the case of allyllithium, the *syn* diastereomer is favored and presumably does not arise from chelation control.

Scheme 1.7: 1,3-Asymmetric Induction in the Allylation of Imines




C. Diastereoselective Imine Allylation Using Chiral Auxiliaries

The use of chiral auxiliaries to control facial selectivity in the addition of allylmetals to imine electrophiles has been widely examined.^{1a} For an auxiliary to be useful, a convenient installation and cleavage protocol needs to be established. For these reasons, chiral auxiliaries attached to the nitrogen atom of the imine are particularly useful and are the topic of this section. Chiral amines, including naturally occurring amino-acids, are attractive auxiliaries because they are widely available, and condensation between the amine and an aldehyde provides a convenient installation method for the auxiliary.

Seminal work in this area was conducted by Yamamoto and coworkers,^{10b} who examined the allylation of imines prepared from the condensation of α -methylbenzylamine with aldehydes. When imine **1.42** is treated with allylmagnesium,

allylboron, or allylstannane nucleophiles, the Cram product **1.43** is formed in preference to anti-Cram product **1.44** (Scheme 1.8). The diastereoselectivity of the reaction can be explained through the cyclic transition structures **1.45** and **1.46**, where steric interactions between the stereocenter on the imine and the ligands are minimized by directing the hydrogen atom substituent toward the metal. The allyl group then adds to the more accessible face of the imine, away from the other substituents on the chiral center. In the case of allyltributylstannane, TiCl_4 is used as a Lewis acid. Good Cram selectivity in this case is rationalized by transmetalation of the allyl group from Sn to Ti, followed by allylation through **1.45** where $\text{M} = \text{Ti}$.


Scheme 1.8: α -Methylbenzylamine as a Chiral Auxiliary for Imine Allylation



Amino acids and vicinal aminoalcohols have been explored as chiral auxiliaries for imine allylation due to their wide availability. Methyl-(*S*)-valinate derived imines (Scheme 1.9, **1.47**) provide excellent diastereocontrol in the addition of a variety of allylmetal reagents including allylcopper, allylcerium, allylaluminum, allyllead,

allylbismuth, and allylmagnesium halides.¹⁶ Here, the nature of the R-group on the imine affects the diastereoselectivity of the reaction. If the R-group is aryl or a branched aliphatic group, the 1,3-*anti* product is obtained; however, if the R-group is simply linear, the 1,3-*syn* product is obtained in good diastereoselectivity. Furthermore, the reaction can be carried out under Barbier conditions using allylbromide, and comparable levels of diastereoselectivity are observed. (*R*)-Phenylglycinol has been used as an auxiliary, giving amination substrates (**1.49**, Scheme 1.9) upon condensation with aromatic aldehydes.¹⁷ When **1.49** is treated with an allylcerium reagent at low-temperature, the imine electrophile is produced *in situ*, furnishing the 1,3-*anti* homoallylic amine product with good diastereocontrol. Subsequent to the allylation, the auxiliary can be cleaved using Pb(OAc)₄; however, in some cases racemization occurred.

Scheme 1.9: Aminoacid Based Auxiliaries for Imine Allylation




¹⁶ Basile, T.; Bocoum, A.; Savoia, D.; Umani-Ronchi, A. *J. Org. Chem.* **1994**, 59, 7766.

¹⁷ Wu, M. -J.; Pridgen, L. M. *Synlett* **1994**, 636.

Ender's SAMP and RAMP hydrazones¹⁸ (**1.52**, Scheme 1.10) have been examined with allylcerium reagents as nucleophiles.¹⁹ Notably, good yields and excellent diastereoselectivities are obtained in the reaction. Furthermore, the N-N bond of the resultant hydrazine products can be readily cleaved, without loss of enantiomeric purity, using Li/NH₃.

Scheme 1.10: Use of SAMP Hydrazones for Imine Allylation



Chiral sulfinylimines developed by Hua²⁰ and Ellman²¹ undergo efficient allylation with high diastereocontrol when a variety of organometallic reagents are used (Scheme 1.11). Notably, ketimines also undergo allylation with allylmagnesium bromide in excellent diastereoselectivity. In addition, the auxiliary can be cleaved quite readily using strong acid. The *p*-toluenesulfinamide is cleaved using trifluoroacetic acid in MeOH, while the *t*-butylsulfinamide is cleaved to the primary amine hydrochloride salt using HCl in MeOH.


¹⁸ Review: Job, A.; Janeck, C. F.; Bettray, W.; Peters, R.; Enders, D. *Tetrahedron* **2002**, 58, 2253.

¹⁹ Enders, D.; Schankat, J.; Klatt, M. *Synlett* **1994**, 795.

²⁰ Hua, D. H.; Miao, S. W.; Chen, J. S.; Iguchi, S. *J. Org. Chem.* **1991**, 56, 4.

²¹ Cogan, D. A.; Liu, G.; Ellman, J. *Tetrahedron* **1999**, 55, 8883.

Scheme 1.11: Allylation of Chiral Sulfinylketimines




D. Use of Chiral Allylmetals for Diastereoselective Imine Allylation

The methods discussed above for stereocontrol in imine allylation reactions all use chirality on the imine electrophile to control the configuration of the forming C-N stereocenter. Introduction of chirality onto the allylmetal nucleophile is another important approach to control the facial selectivity in imine allylation reactions. Optically active allylic silanes²² and allylic stannanes²³ have been prepared by the Panek and Marshall groups, respectively (Scheme 1.12). In the case of allylic silane **1.61**, Lewis acid induced allylation provides aminoester **1.62** in good yield with high chirality transfer. An important feature is that the reactive acylimine electrophile intermediate can be produced *in situ* by condensation of an amine or carbamate with an aldehyde; the pre-formation of imine **1.60** is not required. Marshall's stannane (**1.64**) gives excellent chirality transfer in the Lewis acid-induced allylation of amina **1.63** by reaction through an acyliminium ion.

²² (a) Panek, J. S.; Jain, N. F. *J. Org. Chem.* **1994**, *59*, 2674. (b) Schaus, J. V.; Jain, N.; Panek, J. S. *Tetrahedron* **2000**, *56*, 10263.

²³ Marshall, J. A.; Gill, K.; Seletsky, B. M. *Angew. Chem. Int. Ed.* **2000**, *39*, 953.

Scheme 1.12: Chiral Allylic Silanes and Stannanes for Diastereoselective Imine



While the synthesis of optically active allylic silanes and stannanes, as in **1.61** and **1.64**, is somewhat difficult and requires multi-step synthesis, the use of chirality on the ligand scaffold of the allylmetal reagent is a convenient way to control facial selectivity in the allylation of imines. Seminal work in this area was performed by Itsuno.²⁴ The reaction of *N*-(trimethylsilyl)aldimine **1.66** with the Roush²⁵ chiral allylboronate (**1.70**) or Brown's²⁶ allylborane (**1.71**) affords the optically active imine allylation product, albeit in poor enantioselectivity (Scheme 1.13, eq 1).^{18a} However, application of the Itsuno allylboronate (**1.72**) gives good enantiocontrol in the allylation of **1.66**.^{24b} Subsequently, however, Brown and co-workers²⁷ showed that water is essential to the allylation of *N*-

²⁴ (a) Watanabe, K.; Ito, K.; Itsuno, S. *Tetrahedron Asymm.* **1995**, *6*, 1531. (b) Itsuno, S.; Watanabe, K.; Ito, K.; El-Shehawy, A. A.; Sarhan, A. A. *Angew. Chem. Int. Ed.* **1997**, *36*, 109.


²⁵ (a) Roush, W. R.; Walts, A. E.; Hoong, L. K. *J. Am. Chem. Soc.* **1985**, *107*, 8186. (b) Roush, W. R.; Palkowitz, A. D.; Palmer, M. A. *J. Org. Chem.* **1987**, *52*, 316. (c) Roush, W. R.; Hoong, L. K.; Palmer, M. A. J.; Straub, J. A.; Palkowitz, A. D. *J. Org. Chem.* **1990**, *55*, 4117.

²⁶ Brown, H. C.; Jadhav, P. K. *J. Am. Chem. Soc.* **1983**, *105*, 2092.

²⁷ (a) Chen, G.-M.; Ramachandran, P. V.; Brown, H. C. *Angew. Chem. Int. Ed.* **1999**, *38*, 825. (b) Ramachandran, P. V.; Burghardt, T. E. *Chem. Eur. J.* **2005**, *11*, 4387.


(trimethylsilyl)aldimines (Scheme 1.13, eq 2). The reaction between **1.66** and **1.71** does not occur until water is added. It was rationalized that the role of water was to cleave the weak N-Si bond in **1.66** forming benzaldimine, the reactive electrophile for the allylation reaction, *in situ*. Thus, under the conditions initially explored by Itsuno, the allylation reaction likely takes place during the aqueous workup, which results in poor enantiocontrol when using **1.70** or **1.71**. In support of this hypothesis, it was found that addition of water to **1.66** or **1.68** at -78 °C in the presence of **1.71** resulted in excellent enantiocontrol in the allylation of aryl substituted silylimines (Scheme 1.13, eq 2).

Scheme 1.13: Asymmetric Imine Allylation with Chiral Allylboron Reagents



Ketimines can also be allylated in good yield with good facial selectivity using Soderquist's allylboronate (**1.75**, Scheme 1.14).²⁸ Again, *N*-(trimethylsilyl)aldimines do not react with **1.75** without the addition of water; however, the allylation of *N*-(trimethylsilyl)ketimines (**1.73**) proceeds without the addition of water at -78 °C. Since a mixture of the ketimine along with its enamine tautomer **1.74** are treated with **1.75**, the reaction is argued to proceed via **1.76**, where enamine **1.74** allows access to the more reactive *Z* form of the ketimine. Notably, the homoallylic amine products are isolated in good yields, with moderate to excellent enantioselectivities, and the chiral auxiliary can be recovered.

Scheme 1.14: Asymmetric Allylation of Ketimines with Allylborane 1.75



Allylic silanes **1.79** and **1.82** have been applied to the asymmetric allylation and crotylation of hydrazones **1.78** and **1.81**; a reaction which employs strain-induced Lewis acidity (Scheme 1.15).²⁹ The chiral allylic silanes are readily prepared from the commercially available and relatively inexpensive aminoalcohol pseudoephedrine.³⁰


²⁸ Canales, E.; Hernandez, E.; Soderquist, J. A. *J. Am. Chem. Soc.* **2006**, 128, 8712.

²⁹ Berger, R.; Rabbat, P. M. A.; Leighton, J. L. *J. Am. Chem. Soc.* **2003**, 125, 9596.

³⁰ Kinniard, J. W. A.; Ng, P. Y.; Kubota, K.; Wang, X.; Leighton, J. L. *J. Am. Chem. Soc.* **2002**, 124, 7920.

Excellent yields and enantioselectivities are obtained in the allylation of these hydrazones. In addition, crotylation of benzoylhydrazone **1.81** occurs in good diastereoselectivity and enantioselectivity. Furthermore, use of the *Z* version of **1.82** gave the *anti* diastereomer of **1.83** in comparable levels of yield and selectivity. Notably, this methodology has also been applied to the asymmetric allylation of ketone derived benzoyl hydrazones in 46-92% yield and 86-97% ee.³¹

Scheme 1.15: Use of Chiral Allylic Silanes for Asymmetric Imine Allylation and Crotylation




In addition to the use of isolable chiral allyl and crotylmethyl reagents, the chiral allylmethyl can be generated *in situ* by the addition of a stoichiometric amount of a chiral modifier and an achiral allylmethyl. Kobayashi³² has disclosed the use of BINAP-derived phosphine oxides, in the presence of allyl and crotylsilanes, for the enantio- and diastereoselective allylation and crotylation of hydrazano-esters (**1.84**) with good yields

³¹ Berger, R.; Duff, K.; Leighton, J. L. *J. Am. Chem. Soc.* **2004**, *126*, 5686.

³² Ogawa, C.; Sugiura, M.; Kobayashi, S. *Angew. Chem. Int. Ed.* **2004**, *43*, 6491.

and selectivity's (Scheme 1.16). Significantly, the chiral additive can be recovered in near quantitative yield without loss of enantiomeric purity.

Scheme 1.16: Chiral Additives for Asymmetric Imine Allylation and Crotylation




E. Catalytic Asymmetric Imine Allylation^{1b}

All the methods described above require the use of stoichiometric amounts of a chiral group to affect stereoselective imine allylation. If the stereocenter used to control facial selectivity is not wanted in the final product, removal of this group is both time consuming and wasteful. Therefore the use of asymmetric catalysis to control the facial selectivity in the allylation reaction is the most efficient route to chiral amines via allylation. Recent advances in this area are discussed in this section.

Early reports of a catalytic enantioselective allylation of imines were disclosed by Yamamoto using chiral dimeric Pd-catalyst **1.92** and allyltributylstannane (**1.90**, Scheme 1.17, eq 1).³³ While enantioselectivity near 80% ee is obtained with aromatic imines, aliphatic or cinnamyl imines are less selective (40 and 61% ee, respectively). However, these results prompted further studies in this area. Subsequently, Yamamoto was able to apply this same Pd-catalyst to a catalytic asymmetric allylation of imines with tetra(allyl)silane (Scheme 1.17, eq 2).³⁴ Enantioselectivities and yields are typically improved under these conditions; however, aliphatic and cinnamyl imines are still problematic substrates (52% ee when R = Cy, 74% ee when R = cinnamyl).

Scheme 1.17: Enantioselective Pd-Catalyzed Allyl Additions to Imines



³³ Nakamura, H.; Nakamura, K.; Yamamoto, Y. *J. Am. Chem. Soc.* **1998**, *120*, 4242.


³⁴ Fernandes, R. A.; Yamamoto, Y. *J. Org. Chem.* **2004**, *69*, 735.

Copper complexes bearing chiral bis(phosphine) and diamine ligands have been developed as efficient catalysts for asymmetric imine allylation with allylsilane,³⁵ allylstannane,^{35b} and allylborane³⁶ reagents as nucleophiles. A (*R*)-tol-BINAP derived Cu-catalyst gives effective crotylstannylation of imine **1.94** in good diastereoselectivity and enantioselectivity (Scheme 1.18, eq 1).^{35b} Furthermore, various substituted allylsilanes, in the presence of the chiral Cu-catalyst derived from diamine ligand **1.99**, afford the allylation products in good enantioselectivities (Scheme 1.18, eq 2).^{35c} Recently, aromatic ketimines were shown to be allylated with the pinacol ester of allylboronic acid (**1.102**) in good enantioselectivities and yields (Scheme 1.18, eq 3).³⁶ When **1.101** is employed in the reaction with an aliphatic R-group (R = PhCH₂CH₂) only 23% ee is observed; however, the reaction is quite efficient, affording the allylation product in 98% yield.

³⁵ (a) Ferraris, D.; Dudding, T.; Yolung, B.; Drury, III, W. J.; Letka, T. *J. Org. Chem.* **1999**, *64*, 2168. (b) Fang, X.; Johannsen, M.; Yao, S.; Gathergood, N.; Hazelli, R. G.; Jorgenson, K. A. *J. Org. Chem.* **1999**, *64*, 4844. (c) Kiyohara, H.; Nakamura, Y.; Matsubara, R.; Kobayashi, S. *Angew. Chem. Int. Ed.* **2006**, *45*, 1615.

³⁶ Wada, R.; Shibuguchi, T.; Makino, S.; Oisaki, K.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2006**, *128*, 7687.

Scheme 1.18: Enantioselective Cu-Catalyzed Allyl Additions to Imines




The use of substituted, optically active BINOL derivatives has been exploited for catalytic asymmetric imine allylation. Under Barbier conditions using In-metal and allyl iodide, in the presence of 10% of modified BINOL **1.107**, acylhydrazones (**1.105**) are allylated in good yields and enantioselectivities (Scheme 1.19, eq 1).³⁷ While aromatic acylhydrazones (**1.105**, R = Ar) give the highest enantioselectivities in the reaction (87-99% ee), an aliphatic acylhydrazone (**1.105**, R = PhCH₂CH₂) gave moderate

³⁷ Kargbo, R.; Takahashi, Y.; Bhor, S.; Cook, G. R.; Lloyd-Jones, G. C.; Shepperson, I. R. *J. Am. Chem. Soc.* **2007**, 129, 3846.

enantioselectivity (70% ee). In addition, chiral BINOL-derived Zr-catalysts have been developed for the addition of allylstannanes to imines (Scheme 1.19, eq 2).³⁸ While good enantioselectivities can be obtained using this method, the allylstannane must bear an allylic alcohol moiety in order for high enantioselection to be observed. It should be noted that only aromatic imines were explored using these catalysts.

Scheme 1.19: Substituted Chiral BINOL-Derived Catalysts for Imine Allylation



Finally, chiral BINOL-derivatives have been used as organocatalysts for the enantioselective addition of allylboronate and crotylboronate esters to acylimines


³⁸ Gastner, T.; Ishitani, H.; Akiyama, R.; Kobayashi, S. *Angew. Chem. Int. Ed.* **2001**, *40*, 1896.

(Scheme 1.20).^{39,40} Notably, both aromatic and aliphatic imines are allylated in excellent enantioselectivities and yields. Importantly, the crotylation is not stereospecific. That is, when (*E*)-**1.118** is used in the crotylation, high *anti*-selectivity is observed (>99:1 dr), and the use of (*Z*)-**1.118** also preferentially forms the *anti*-diastereomer of product in high selectivity (98:2 dr). The authors rationalize this result by suggesting that the *Z*-configured imine, which is in equilibrium with the more stable *E*-configured imine, is more reactive. The reaction then proceeds through a chair-like transition state when (*E*)-**1.118** is employed, furnishing the *anti*-product. However, when (*Z*)-**1.118** is used as the crotylboronate reagent, the crotylation of the *Z*-configured imine must proceed through a boat-like transition state to afford the *anti*-product. The preference for a boat-like transition state for the crotylation with (*Z*)-**1.118** presumably results from a penalizing *pseudo*-1,3-diaxial interaction between the methyl group on the crotylboronate and the acyl group of the *Z*-configured imine that results from reaction through a chair-like transition state.

³⁹ Lou, S.; Moquist, P. N.; Schaus, S. E. *J. Am. Chem. Soc.* **2007**, *129*, 15398.

⁴⁰ For seminal work on the use of BINOL-derived allylboronates for use in asymmetric allylation of aldehydes, see: (a) Thormeier, S.; Carboni, B.; Kaufmann, D. E.; *J. Organomet. Chem.* **2002**, *657*, 136. (b) Wu, T. R.; Shen, L.; Chong, J. M. *Org. Lett.* **2004**, *6*, 2701.

Scheme 1.20: Organocatalytic Asymmetric Addition of Allyl and Crotylboronates




F. Catalytic Diboration of Allenes

The addition of bis(metal) reagents of type M-M or M-M' to unsaturated carbon-carbon bonds, using transition metal catalysis, is a highly useful process in organic synthesis.² Of the group 10 transition metals, Pt has found widespread use as a catalyst in this area.^{2d} The first example of an allene diboration was disclosed by Miyaura and co-workers in 1998 (Scheme 1.21).⁴¹ In general, good yields are obtained when commercially available diboron reagent **1.121** is used in the diboration reaction. Regiochemistry in the reaction depends on the substituent on the allene. For alkyl

⁴¹ Ishiyama, T.; Kitano, T.; Miyaura, N. *Tetrahedron Lett.* **1998**, 39, 2357.


substituted allenes, diboration occurs at the internal olefin of the allene; however, for aromatic or heteroatom substituted allenes, diboration occurs at the external olefin of the allene. $\text{Pt}(\text{PPh}_3)_4$ is also an effective catalyst for this transformation giving improved yields and regioselectivities in some cases.

Scheme 1.21: Pt-Catalyzed Diboration of Allenes




The proposed mechanism for this process is shown in Scheme 1.22. Oxidative addition of $\text{B}_2(\text{pin})_2$ to Pt gives diboronate Pt-complex **1.124**. Insertion of the allene then gives rise to complex **1.125** or **1.126**, which upon reductive elimination releases the Pt catalyst and affords product **1.122** or **1.123**.

Scheme 1.22: Miyaura's Proposed Mechanism for the Pt-Catalyzed Diboration of Allenes



The potential for the products of allene diboration to participate in a subsequent allylation reaction has been demonstrated (Scheme 1.23). When benzaldehyde is treated with diboronate ester **1.127** in dioxane, followed by Suzuki-Miyaura cross-coupling⁴² with iodobenzene, homoallylic alcohol **1.128** is isolated in good yield. Only diboronate **1.127** has been explored in this type of allylation reaction.

Scheme 1.23: Allylation of Benzaldehyde with a Product of Allene Diboration



⁴² Reviews: (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, 95, 2457. (b) Suzuki, A. *J. Organomet. Chem.* **1999**, 576, 147. (c) Kohta, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, 58, 9633.

III. Development of the Pd-Catalyzed Enantioselective Diboration of Allenes and its Application to the Allylation of Imines

A. The Pd-Catalyzed Enantioselective Diboration of Allenes³

As described in the introduction, the bis(metalation) of the internal olefin of an allene may provide a useful allylboronate for the allylation of imine electrophiles. Precedent from Miyaura's group verifies that these intermediates can be used in aldehyde allylation; however, to access synthetically useful chiral allylboronates, highly regioselective diboration of the internal olefin of the allene must occur. Furthermore, to arrive at optically active allylation products, the diboronate intermediate must be formed in enantiomerically enriched form. Thus, we began to study the allene diboration reaction using chiral phosphine ligands as a means to access chiral allene diboration products with high enantiomeric purity.


Before I joined this project, fellow graduate student Nicholas Pelz had determined that the use of electron-rich phosphines, such as PCy₃, HMPT, or P(OEt)₃, in the presence of Pd₂(dba)₃, furnished exclusive diboration to the internal olefin of decyl allene in only 1 h at room temperature. This was surprising since Pd-catalyzed diboration reactions are rare.^{2e,43} More importantly, after a short screen of chiral phosphine ligands, he and Angela Woodward discovered that the use of (TADDOL)MONOPHOS⁴⁴ (**1.129**, Figure 1.2) in the diboration of decyl allene led to product formation with complete regiocontrol

⁴³ Cui, Q.; Musaev, D.; Morokuma, K. *Organometallics* **1998**, 17, 742.

⁴⁴ (a) Keller, E.; Maurer, J.; Naasz, R.; Schader, T.; Meetsma, A.; Feringa, B. L. *Tetrahedron Asymm.* **1998**, 9, 2409. (b) Alexakis, A.; Vastra, J.; Burton, J.; Benhaim, C.; Manganey, P. *Tetrahedron Lett.* **1998**, 39, 7869.


in good yield and enantioselectivity. It was at this point that I joined this project and was involved in examining the substrate scope of the allene diboration reaction.

Figure 1.2: Structure of (TADDOL)MONOPHOS




The substrate scope in the asymmetric Pd-catalyzed allene diboration is shown in Table 1.3. Diboration of either aromatic or aliphatic allenes occurred at the internal olefin of the allene with complete regiocontrol. Furthermore, the products were purified using silica gel chromatography and isolated in good yield and enantiomeric purity. In general, diboration of aliphatic allenes occurred with higher enantioselectivity, but in lower yields relative to aromatic allenes. In addition, the sterically hindered substrate *t*-butyl allene participated in the reaction; however, longer reaction times were required to afford useful product yields (entry 6).

Table 1.3: Substrate Scope in the Asymmetric Pd-Catalyzed Allene Diboration



1.120

$+ \text{B}_2(\text{pin})_2 \xrightarrow[\text{PhMe, rt, 14 h}]{2.5\% \text{ Pd}_2(\text{dba})_3, 6\% \text{ 1.129}}$



1.129

entry	R	% yield ^a	% ee ^b
1	decyl	61	91
2	Cy	62	89
3	PhCH ₂ CH ₂	73	90
4	Me	68	92
5	Ph	75	87
6	<i>t</i> -Bu	42 (58) ^c	89 (88) ^c
7	BnOCH ₂ CH ₂	57	91
8	Bn	65	90


^aIsolated yield of diboronate ester product. Yield represents an average of 2 experiments in each case.

^bEnantiomeric excess was determined by chiral GLC or SFC analysis of the diol derived from hydrogenation followed by oxidation with H₂O₂. ^cValue in parentheses represents data for a 48 h reaction.

The procedure for determining the enantiomeric purity of the diboronate ester products (**1.122**) from the allene diboration deserves mentioning, as **1.122** could not be directly analyzed by typical methods. Initially, the olefin in **1.122** was hydrogenated using Pd/C and H₂ gas, followed by oxidation of the C-B bonds with alkaline H₂O₂ to afford a diol, as a mixture of diastereomers (Scheme 1.24). The diol products from this reduction/oxidation sequence could then be analyzed using chiral GLC or chiral SFC to determine their enantiomeric purities. Surprisingly, this method gave non-reproducible

results, and the *syn* and *anti* diastereomers of the resultant diols did not possess the same enantiomeric purity.


Scheme 1.24: Enantiomeric Excess Determination of (S)-1.122



Heterogeneous Pd-catalysts for the hydrogenation of olefins are known to isomerize alkenes during the hydrogenation process.⁴⁵ We reasoned that isomerization of (S)-1.122 to a mixture of internal alkene isomers, during the heterogeneous hydrogenation, was the reason for the differences in enantiomeric purity between the two diastereomeric diols formed after oxidation. Since this ultimately leads to deterioration in the enantiomeric purity of 1.130 during the hydrogenation of (S)-1.120, this method would not give accurate enantioselectivity data for the diboration reaction. To ascertain if β -hydride elimination (presumed mechanism for isomerization) was occurring in the hydrogenation of (S)-1.122 with Pd/C, diboronate ester 1.132 was subjected to the hydrogenation conditions using D₂ gas in place of H₂ (Scheme 1.25). Deuterium scrambling was observed in the resultant diols, indicating that β -hydride elimination was likely occurring during the hydrogenation.


⁴⁵ Siegel, S. Heterogeneous Catalytic Hydrogenation of C=C and C \equiv C. In *Comprehensive Organic Synthesis*; Fleming, I., Trost, B. M. Eds.; Pergamon Press: New York, 1991; Vol. 8, pp 417-442.

Scheme 1.25: Hydrogenation of (S)-1.132 with D₂



This problem was easily solved by conducting the hydrogenation with diimide (Scheme 1.26).⁴⁶ Diimide is known to reduce olefins through a concerted process via *syn* addition of H₂ with concomitant loss of N₂. NBSH⁴⁷ was used to produce diimide *in situ* for the reduction of (S)-1.122.⁴⁸ After oxidation with H₂O₂, the resultant diastereomeric diols had the same enantiomeric purity, and reproducible results could be obtained.

Scheme 1.26: Diimide Reduction of (S)-1.122



B. Allylation of Imines with the Products of Allene Diboration

With access to enantiomerically enriched diboronate esters from the diboration of allenes, the use of these reagents in allylation reactions with imines could now be explored. Initially, the reaction between diboron **1.132** and (*E*)-aldimines (**1.135**) was


⁴⁶ (a) Corey, E. J.; Mock, W. L.; Pasto, D. J. *Tetrahedron Lett.* **1961**, 347. (b) Corey, E. J.; Pasto, D. J.; Mock, W. L. *J. Am. Chem. Soc.* **1961**, 83, 2957. (c) Review: Pasto, D. J.; Taylor, R. T. *Org. React.* **1991**, 40, 91.

⁴⁷ (a) Dann, A. T.; Davies, W. *J. Chem. Soc.* **1929**, 1050. (b) Myers, A. G.; Zheng, B. *J. Am. Chem. Soc.* **1996**, 118, 4492.

⁴⁸ (a) Haukaas, M. H.; O'Doherty, G. H. *Org. Lett.* **2002**, 4, 1771. (b) Hunig, S.; Muller, H. R.; Their, W. *Angew. Chem. Int. Ed.* **1965**, 4, 271.


examined (Scheme 1.27, eq 1). No reaction was observed with benzyl, tosyl, or silyl-substituted imines, even after heating to 60 °C. Considering the chair-like transition structures **Chair-TS-1** and **Chair-TS-2** as possible reaction pathways for the desired allylation reaction, it was postulated that **1.135** was inert to allylation because of the high steric demands present in **Chair-TS-1** and **Chair-TS-2**. In an attempt to alleviate strain in the transition state, the ammonia-derived imine of benzaldehyde (**1.135**, R = H) was employed. This imine could be generated *in situ* from imine **1.66** by protodesilylation with water (Scheme 1.27, eq 2). ¹H NMR analysis of this reaction indicated that the starting material was completely consumed. Subsequently, the allylation product was acylated, and oxidized with buffered H₂O₂ to give β-amidoketone **1.136**. Notably, the reaction occurred with a high level of chirality transfer in good yield, with >99% ee material being obtained after one recrystallization.

Scheme 1.27: Initial Attempts at Imine Allylation with 1.132



To make this process more efficient, development of a single-flask tandem allene diboration/imine allylation was pursued to obviate the need for isolation of the diboronate ester intermediate. Gratifyingly, treatment of the unpurified allene diboration reaction mixture with silylimine **1.66** and 1.1 equiv of MeOH, followed by acetylation and buffered oxidation, led to formation of β -amidoketone **1.136** in excellent overall yield with complete chirality transfer (Scheme 1.28). These conditions proved to be general, and the substrate scope is described in Section C of this chapter (see page 39).

Scheme 1.28: Tandem Allene Diboration/Imine Allylation




In addition to protonolysis of silyl-derived imines, direct condensation of benzaldehyde with an ammonia source provides access to reactive N-H substituted aldimines. Use of this procedure for imine allylation with the products of allene diboration would obviate the need for synthesizing and isolating the imine electrophile. As noted previously, Kobayashi⁸ has developed this protocol for imine allylboration, where the imine is formed *in situ* by reaction of an ammonia source with an aldehyde. In his report, the ammonia source consists of either liquid NH_3 in EtOH or a 28-30 wt% ammonium hydroxide solution in EtOH. In our initial experiments, a solution of diboronate ester **1.132** was added to a stirred solution of benzaldehyde in an ethanolic solution of liquid NH_3 or 29 wt% NH_4OH according to Kobayashi's procedure. It appeared as if some imine allylation product was present in the 1H NMR spectrum of the unpurified reaction mixture when NH_4OH was used as the ammonia source. However, after acetylation and oxidation, only a 30% yield of the desired β -amidoketone product was isolated (Scheme 1.29, eq 1). Further analysis of the 1H NMR spectrum of the unpurified reaction mixture, before acetylation and oxidation, suggested that unreacted diboronate ester **1.132** was present. However, the chemical shifts for the benzylic proton in **1.132** and one of the olefinic protons were shifted slightly downfield. Furthermore,

when **1.132** was mixed with 29 wt% NH₄OH in EtOH overnight, these same resonances shifted downfield in the ¹H NMR spectrum. These observations may be explained by a change in the ligand environment around the B-atom in **1.132**, possibly by interaction of hydroxide or ammonia with the vacant orbital on boron. This interaction may decrease the reactivity of **1.132** by blocking the vacant orbital on boron that is needed for coordination to the imine electrophile in the allylation reaction. We postulated that if hydroxide was the culprit, then this problem could be avoided with the use of NH₄OAc, since the acetate ion is less basic and nucleophilic relative to hydroxide. This hypothesis was initially examined by performing the reaction as before, using an ethanolic solution of 29 wt% NH₄OH, but with an equimolar amount of AcOH relative to NH₄OH added to the mixture (Scheme 1.29, eq 2). While the yield did improve, chirality transfer was poor. However, when NH₄OAc was used directly (in MeOH with activated 4 Å molecular sieves), good yield and chirality transfer was obtained (Scheme 1.29, eq 3).⁴⁹ Notably, the unpurified allene diboration reaction mixture was employed, obviating the need to purify the diboronate ester nucleophile. The scope of this method is described in the next section (see page 40).

⁴⁹ The ligand used in the diboration reaction (**1.138**, Scheme 1.26, eq 3) was identified by Heather Burks in the group to give improved enantioselection in the allene diboration. See: Woodward, A. R.; Burks, H. E.; Chan, L. M.; Morken, J. P. *Org. Lett.* **2005**, 7, 5505 and reference 3b.

Scheme 1.29: Allylation of *in situ* Prepared Imines with 1.132



C. Scope of the Tandem Allene Diboration/Imine Allylation and Other Transformations

The scope of the tandem asymmetric allene diboration/imine allylation of silyl-derived imines is given in Table 1.4. It should be noted that while this method was being optimized, phosphoramidite ligand **1.138** was identified by Heather Burks in the group to give enhanced enantioselectivity in the diboration reaction, and therefore, this ligand was now used in the allene diboration.⁴⁹ Both aromatic and aliphatic allenes were effective in the tandem diboration/allylation, and good yields were obtained (~60-70%). The only exception was the reaction employing cyclohexyl allene and *N*-(trimethylsilyl)benzaldimine (entry 7). Considering that four reactions are carried out

consecutively in one flask (diboration, allylation, acetylation, and oxidation), the yield per step is >87%. Importantly, only imines devoid of α -protons could be used, since the synthesis of *N*-silyl imines having α -protons is an unsolved problem. However, an aliphatic, α,β -unsaturated imine could be prepared and allylated in good yield and selectivity (entries 3, 6, and 9). In general, the conservation of ee (% cee) was the lowest for this imine electrophile.⁵⁰

Table 1.4: Tandem Allene Diboration/Imine Allylation of Silylimines

entry	R	R'	% yield ^a	% ee ^{b,c}	% cee ^d
1	Ph	Ph	68	97	99
2	Ph	2-furyl	68	97	99
3	Ph	(<i>E</i>)-hexenyl	70	92	94
4	PhCH ₂ CH ₂	Ph	64	93	95
5	PhCH ₂ CH ₂	2-furyl	69	92	94
6	PhCH ₂ CH ₂	(<i>E</i>)-hexenyl	70	89	91
7	Cy	Ph	46	91	98
8	Cy	2-furyl	66	92	99
9	Cy	(<i>E</i>)-hexenyl	59	87	94

^aYield of β -amidoketone after silica gel chromatography; average of 2 experiments in each case.


^bEnantiomeric excess determined by chiral HPLC or SFC analysis of the β -amidoketone. ^cEnantiomeric purity for the diboron intermediate for R = Ph, PhCH₂CH₂, and Cy was 98, 98, and 93% ee, respectively.

^dDefined as (% ee **1.140** \div % ee **1.122**) \times 100.

⁵⁰ The term conservation of ee (cee) was introduced by Evans and co-workers. See: Evans, P. A.; Robinson, J. E.; Nelson, J. D. *J. Am. Chem. Soc.* **1999**, *121*, 6761.

Imines prepared *in situ*, from the condensation of an aldehyde with NH₄OAc in MeOH, were examined in the tandem diboration/allylation (Table 1.5). In general, comparable levels of chirality transfer were observed employing this method; however, yields were somewhat lower. In the case of cyclohexyl allene, poor yields were obtained, partly due to competitive allylation of the aldehyde present in the reaction pot. Aliphatic aldehydes did not give satisfactory yields (<8% with butyraldehyde) and were not examined further. Since the silyl-derived imines are moisture sensitive and somewhat difficult to prepare and isolate, this process offers a convenient route to access the same allylation products without the need of isolating these sensitive imines.

Table 1.5: Tandem Diboration/Imine Allylation of *in situ* Prepared Imines

					
entry	R	aldehyde	% yield ^a	% ee ^{b,c}	% cee ^d
1	Ph	benzaldehyde	59	96	98
2	Ph	2-furfural	69	96	98
3	PhCH ₂ CH ₂	benzaldehyde	59	92	94
4	PhCH ₂ CH ₂	2-furfural	56	91	93
5	Cy	benzaldehyde	30	91	98
6	Cy	2-furfural	39	93	100

^aYield of β-amidoketone after silica gel chromatography; average of 2 experiments in each case.


^bEnantiomeric excess determined by chiral HPLC or SFC analysis of the β-amidoketone. ^cEnantiomeric purity for the diboron intermediate for R = Ph, PhCH₂CH₂, and Cy was 98, 98, and 93% ee, respectively.

^dDefined as (% ee **1.140** ÷ % ee **1.122**) x 100.

A unique feature of the tandem allene diboration/imine allylation product is that it contains a vinylboronate functional group which may be further manipulated. Beside oxidation, as discussed above, this vinylboronate group participated in Suzuki-Miyaura cross-coupling⁴² (Scheme 1.30, eq 1); protection of the amine in this sequence with (Boc)₂O gave the Boc-derived vinylboronate ester **1.142** in good yield as one olefin isomer. Suzuki-Miyaura coupling with iodobenzene gave the coupling product **1.142** in good yield and excellent enantiomeric excess. When this coupling reaction was performed using the acetyl protected amine analogous to **1.141**, low yields were obtained, which was presumably due to deprotection of the amine by aqueous KOH during the cross-coupling. Use of the Boc protecting group solved this problem. Aside from oxidation and cross-coupling, the vinylboronate functionality was also protonated in refluxing propionic acid to afford **1.144** (Scheme 1.30, eq 2).⁵¹ This method allows access to *Z*-configured allylation products that can be difficult to prepare by other allylation methods.

⁵¹ (a) Brown, H. C.; Gupta, S. K. *J. Am. Chem. Soc.* **1972**, *94*, 4370. (b) Brown, H. C.; Zweifel, G. J. *J. Am. Chem. Soc.* **1961**, *83*, 3834. (c) Brown, H. C.; Hébert, N. C. *J. Organomet. Chem.* **1983**, *255*, 135.

Scheme 1.30: Access to Other Homoallylic Amides



The olefin geometry of the allylation products was determined by crystallographic analysis of the unprotected amine, and by analysis of the coupling constant of the olefinic protons in **1.144**. To obtain crystals suitable for analysis, the unpurified allene diboration reaction was treated with **1.66** and MeOH to afford allylation product **1.145** (Scheme 1.31). A white solid was obtained that could be purified by crystallization. The X-ray structure of **1.145** (Figure 1.3) verified the configuration of the alkene.

Scheme 1.31: Isolation of the Allylation Intermediate






Figure 1.3: Crystal Structure of 1.145



D. Stereochemical Analysis in the Imine Allylation

Considering that allylboron reagents add to aldehyde and imine electrophiles through cyclic transition states,⁵ a model for the observed chirality transfer in the imine allylation with allene diboration products could be developed. Upon analysis of the possible cyclic transition states for the reaction of (*E*)-aldimines with diboronate ester **1.122**, it was not surprising that silylaldimines were not reactive until the N-Si bond was cleaved with H₂O or MeOH (Figure 1.4). The possible cyclic transition structures for reaction of **1.122** with an (*E*)-aldimine are shown in Figure 1.4. All possible chair-like and boat-like transition structures have considerable 1,3-diaxial interactions or allylic strain.


Figure 1.4: Transition Structures for Allylation of (*E*)-Aldimines



Ammonia derived aldimines were required for successful reaction, which presumably is due to a relief in strain exhibited in the allylation transition state. Furthermore, it is likely that these N-H derived aldimines are formed as an *E/Z* mixture during the reaction. This then allows for eight possible transition structures for the allylation reaction (Figure 1.5). Transition structures **CTS-1**, **CTS-2**, **BTS-1**, and **BTS-2** represent reaction of the (*Z*)-aldimine, and transition structures **CTS-3**, **CTS-4**, **BTS-3**, and **BTS-4** represent reaction of the (*E*)-aldimine. The only product observed from the allylation reaction is of the *R*-configuration at the newly formed stereocenter and possesses a *Z*-configured alkene. According to the models in Figure 1.5, the allylation reaction product results from either chair-like transition state **CTS-1** or boat-like transition state **BTS-4**. Of the possible transition states for the (*Z*)-aldimine, reaction most likely occurs through **CTS-1**, where the imine substituent resides in an equatorial position, with the substituent on the diboron reagent axial to avoid a penalizing A(1,2) interaction present in **CTS-2**. The (*E*)-aldimine most likely reacts through **BTS-4**. All

other transition states have penalizing 1,3-diaxial or allylic strain. While **BTS-4** correctly predicts the outcome observed in the allylation reaction, the lowest energy pathway is likely reaction of the (*Z*)-aldimine through **CTS-1**. Isomerization between the *E*- and *Z*-aldimines under the reaction conditions may allow for complete product formation through **CTS-1**.

Figure 1.5: Possible Transition Structures in the Imine Allylation



IV. Conclusions

Examination of the substrate scope in the asymmetric Pd-catalyzed allene diboration has been examined. These products were used as allylmetal reagents for imine allylation reactions to furnish atypical allylation products without the need for isolation of the diboronate ester intermediates. The following insights have been gained from this study:

1. Aliphatic and aromatic allenes undergo allene diboration exclusively at the internal olefin of the allene in good yield and with high enantioselectivity.
2. The unpurified diboronate ester products obtained from the allene diboration allylated *N*-(trimethylsilyl)aldimines with excellent chirality transfer when water or MeOH was added to the reaction medium. Subsequent oxidation, protonation, or Suzuki-Miyaura cross-coupling could be carried out in acceptable yield without loss of enantiomeric purity.
3. The unpurified diboronate ester products obtained from the allene diboration allylated ammonia-derived aldimines prepared *in situ* from an aldehyde and NH₄OAc. The chirality transfer for this process was comparable to the analogous reaction employing *N*-(trimethylsilyl)aldimines.
4. A model to explain the stereochemical outcome in the imine allylation has been developed based on cyclic chair-like transition structures.

V. Experimental Procedures

A. General. Melting points were determined using a Mel-Temp II melting point apparatus and are uncorrected. ^1H NMR spectra were recorded on Bruker DRX 300 or 400 MHz spectrometers. Chemical shifts are reported in ppm from tetramethylsilane with the residual solvent resonance as the internal standard (CHCl_3 : 7.24 ppm). Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, h = hextet, br = broad, m = multiplet), coupling constants (Hz) and assignment. ^{13}C NMR spectra were recorded on a Bruker 400 spectrometer (100 MHz) with complete proton decoupling. Chemical shifts are reported in ppm from tetramethylsilane with the solvent as the internal standard (CDCl_3 : 77.0 ppm). High-resolution mass spectrometry was performed by the University of Minnesota Mass Spectrometry Service Laboratory. Low-resolution mass spectrometry was performed by the University of North Carolina, Department of Chemistry Mass Spectrometry Facility. Infrared (IR) spectra were obtained on a Nicolet 560 Magna-FTIR. X-ray crystallography was performed by Dr. Peter White of the University of North Carolina X-ray facility.

Liquid chromatography was performed using forced flow (flash chromatography) on silica gel (SiO_2 , 230 X 450 mesh) purchased from Sorbent Technologies. Thin layer chromatography was performed on 250 μm silica gel plates from EMD Chemicals Inc. Visualization was achieved using UV light, phosphomolybdic acid in ethanol, potassium permanganate in water, or cerium sulfate and ammonium molybdate in sulfuric acid, each followed by heating.

Analytical gas-liquid chromatography (GLC) was performed on a Hewlett-Packard 6890 Series chromatograph equipped with a CTC Analysis Combi Pal autosampler by Leap Technologies (Carrboro, NC), a split mode capillary injection system, a flame ionization detector, and a Supelco β -dex 120 column with helium as the carrier gas. Analytical high performance liquid chromatography (HPLC) was performed on a Hewlett-Packard Series 1100 liquid chromatograph equipped with a UV detector and a Daicel Chiralcel OD-H column. Analytical supercritical fluid chromatography (SFC) was performed on a Berger Instruments supercritical chromatograph equipped with an Alcott autosampler and a Knauer UV detector.

All reactions were conducted in oven- or flame-dried glassware under an inert atmosphere of nitrogen or argon. Toluene was distilled over CaH_2 and degassed by freeze-pump-thaw cycles prior to use. Anhydrous THF was freshly distilled from Na metal and benzophenone. Tris(dibenzylideneacetone)dipalladium $[\text{Pd}_2(\text{dba})_3]$ and tricyclohexylphosphine were purchased from Strem Chemical Company. (TADDOL)MONOPHOS (**1.129**) was generously donated by Dr. Andre de Vries and Dr. David Ager of DSM Chemical. $[(R,R)\text{-xylylTADDOL}]\text{PNMe}_2$ (**1.138**) was synthesized according to the literature.⁵² Bis(pinacolato)diboron $[\text{B}_2(\text{pin})_2]$ was purchased from BASF or Aldrich Chemical Companies and used directly. Allenes were synthesized according to the literature.⁵³ Aldehydes were purchased from Aldrich Chemical Company and distilled prior to use. *N*-(trimethylsilyl)aldimines were synthesized via the


⁵² Boele, M. D. K.; Kamer, P. C. J.; Lutz, M.; Spek, A. L.; de Vries, J. G.; van Leeuwen, P. W. M. N.; van Strijdonck, G. P. F. *Chem. Eur. J.* **2004**, *10*, 6232.

⁵³ Vermeer, P.; Meijer, J.; Brandsma, L. *Recl. Trav. Chim. Pays-Bas* **1975**, *94*, 112.

known literature procedure⁵⁴ and distilled prior to use. Acetic anhydride was purchased from Fisher Chemical and distilled under N₂. All other reagents were purchased from either Fisher or Aldrich Chemical Companies and used directly.

B. Experimental Procedures

Synthesis of Diboronate Ester **1.132**



An oven-dried 20 mL scintillation vial, equipped with a magnetic stir-bar, was charged with 9.9 mg (0.011 mmol) of tris(dibenzylideneacetone)dipalladium(0), 13.9 mg (0.0258 mmol) of **1.129**, and 2.7 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min, and then 131 mg (0.516 mmol) of B₂(pin)₂ was added, followed by 50.0 mg (0.430 mmol) of phenyl allene. The vial was capped, taped with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 14 h. Volatile material was removed under reduced pressure, and silica gel chromatography of the mixture afforded 127 mg (0.343 mmol, 80%) of **1.132** as a viscous oil.

⁵⁴ Colvin, E. W.; McGarry, D.; Nugent, M. J. *Tetrahedron* **1988**, *44*, 4157.


(S)-4,4,5,5-Tetramethyl-2-[1-phenyl-1-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-en-2-yl]-1,3,2-dioxaborolane (1.132). $R_f = 0.22$ (15:1 hexanes:EtOAc). IR (neat): 2970 (br, s), 2924 (br, m), 1604 (w), 1417 (br, m), 1313 (br, s), 1142 (br, s) cm^{-1} ; ^1H NMR: δ 7.10-7.26 (5H, m, aromatic), 5.81 (1H, br s, CBCHH), 5.27 (1H, br s, CBCHH), 3.36 (1H, s, ArCHB), 1.20-1.25 (24H, m, OCCH₃); ^{13}C NMR: δ 141.0, 130.4, 128.6, 128.4, 125.8, 83.93, 83.89, 25.38, 25.23, 24.99, 24.93. Carbons with directly attached boron atoms were not observed, most likely due to quadrupolar relaxation.⁵⁵ HRMS (ESI) Calcd for C₂₁H₃₂B₂O₄ (M + Na)⁺: 393.2379, Found (M + Na)⁺: 393.2393.


Proof of stereochemistry. The diboronate ester product was converted to a 1,2-diol using hydrogenation (diimide)/oxidation (Scheme 1.26) and is described on page 54. Analysis of the diols by chiral GLC was used to determine the enantioselectivity of the reaction. Stereochemical ratios were determined in comparison to authentic racemic materials prepared by the diboration of phenylallene using PCy₃ as the achiral ligand, followed by the above reduction-oxidation sequence. The absolute configuration was established in comparison to authentic (1*R*,2*R*)-1-phenylpropane-1,2-diol synthesized via Sharpless asymmetric dihydroxylation of β -methylstyrene.⁵⁶

⁵⁵ Wrackmeyer, B. *Prog. NMR Spectrosc.* **1979**, 12, 227.

⁵⁶ Norrby, P-O.; Becker, H.; Sharpless, K. B. *J. Am. Chem. Soc.* **1996**, 118, 35.

Chiral GLC (β -dex, Supelco, 140 °C) analysis of diol products resulting from the reduction-oxidation of **1.132**:






(*S*)-2-[4,4-Dimethyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pent-1-en-2-yl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.150).

The product was prepared according to the general allene diboration procedure used to prepare **1.132** on page 49. A white solid. $R_f = 0.21$ (15:1 hexanes:EtOAc). IR (KBr pellet): 2986 (br, s), 2943 (br, s), 2866 (br, m) 1926 (w), 1600 (w), 1344 (br, m) cm^{-1} ; ^1H NMR: δ 5.89 (1H, d, $J = 3.6$ Hz, CBCHH), 5.71 (1H, d, $J = 3.6$ Hz, CBCHH), 1.96 (1H, s, $(\text{CH}_3)_3\text{CBH}$), 1.17-1.22 (24H, m, OCCCH_3), 0.89 (9H, s, $(\text{CH}_3)_3\text{CB}$); ^{13}C NMR: δ 132.5, 83.7, 82.9, 33.4, 29.8, 25.3, 25.2, 25.1, 25.0. Carbons with directly attached boron atoms were not observed, most likely due to quadrupolar relaxation.⁵³ HRMS (ESI) Calcd for $\text{C}_{19}\text{H}_{36}\text{B}_2\text{O}_4$ ($\text{M} + \text{Na}$)⁺: 373.2692, Found ($\text{M} + \text{Na}$)⁺: 373.2701.

Proof of Stereochemistry. The diboronate ester product was converted to a 1,2-diol using hydrogenation (diimide)/oxidation (Scheme 1.26) and is described on page 54. Analysis of the diols by chiral GLC was used to determine the enantioselectivity of the reaction. Stereochemical ratios were determined in comparison to authentic racemic materials prepared by the diboration of *t*-butylallene using PCy_3 as the achiral ligand, followed by the above reduction-oxidation sequence. The absolute configuration was established in comparison to authentic (*2R,3R*)-4,4-dimethylpentane-2,3-diol synthesized via Sharpless asymmetric dihydroxylation of (*E*)-4,4-dimethyl-2-pentene.⁵⁷

⁵⁷ Kolb, H. C.; VanNieuwenhze, M. S.; Sharpless, K. B. *Chem. Rev.* **2004**, 94, 2483.


Chiral GLC (β -dex, Supelco, 80 - 140°C at 1°C/min) analysis of diol products resulting from the reduction-oxidation of **1.150**:



All other allene diboration products listed in Table 1.3 were characterized previously by other group members.^{3a}


Representative Procedure for Hydrogenation/Oxidation of the Allene Diboration

Products to Assay Enantiomeric Purity




To 40.0 mg (0.108 mmol) of **1.132** in a 20 mL scintillation vial with magnetic stir-bar was added 1.08 mL of anhydrous dioxane in a dry-box. NBSH⁴⁷ (235 mg, 1.08 mmol) was added and the vial capped with a septum and removed from the dry-box. Next, 0.23 mL (1.62 mmol) of TEA was added by syringe, and the septum was replaced with a polypropylene cap, sealed with electrical tape, and heated at 95 °C for 3 h. This mixture was allowed to reach room temperature, and 3 mL of 3 M NaOH was added, followed by slow addition of 1 mL of aqueous 30% H₂O₂. The vial was purged with N₂, loosely capped, and allowed to stir at ambient temperature for an additional 3 h. This mixture was subsequently cooled to 0 °C and quenched by slow addition of 6 mL of saturated aqueous Na₂S₂O₃. The resultant reaction mixture was extracted with EtOAc (3x, ~30 mL). The combined organics were washed with 10 mL of 1M NaOH followed by 15 mL of brine. Finally, drying with anhydrous Na₂SO₄ followed by removal of volatile material under reduced pressure afforded diol **1.151**. The diol was purified by silica-gel chromatography (hexanes/EtOAc), and its structure confirmed by comparison of the ¹H NMR spectrum with the literature.

Representative Procedure for Reduction-Oxidation Sequence with Catalytic Pd/C




A dry 2-dram vial with magnetic stir-bar was charged with 8.6 mg of 10% Pd/C, capped with a septum, and purged with N₂. A solution of 29.8 mg (0.0805 mmol) of **1.132** in 0.4 mL of EtOH was added by canula, and the flask rinsed over with another 0.1 mL of EtOH. The resultant reaction mixture was sparged with H₂, and put under 1 atm of H₂ using a balloon. After 24 h at ambient temperature, the mixture was filtered through a pad of celite and washed with EtOH. To the resulting solution was added 0.25 mL of 3M NaOH, followed by slow addition of 0.15 mL of aqueous 30% H₂O₂. The solution was then left to stir at ambient temperature, under N₂, for 3 h. Finally, the reaction was cooled to 0 °C and quenched by slow addition of 2 mL saturated aqueous Na₂S₂O₃. Next, 10 mL of 1 M NaOH was added, and this mixture extracted with EtOAc (2x, ~25 mL). The combined organics were washed with brine (1x, 10 mL) and dried with anhydrous Na₂SO₄. Removal of volatile material under reduced pressure afforded an oil, which was purified by silica gel chromatography (hexanes/EtOAc) to afford the diols as a mixture of diastereomers. The structure was confirmed by comparison of the isolated material's ¹H NMR data with that reported in the literature.

Procedure for Reduction-Oxidation Sequence with Catalytic Pd/C and D₂



To 42 mg (0.11 mmol) of *rac*-**1.132** in a dry 2-dram vial with magnetic stir-bar under N₂ was added 0.7 mL of EtOH. Next, 12.1 mg of 10% Pd/C was added, and the reaction mixture was sparged with D₂ and put under 1 atm of D₂ using a balloon. After 17 h at ambient temperature, the mixture was filtered through a pad of celite and washed with EtOH. To the resulting solution was added 0.35 mL of 3 M NaOH followed by slow addition of 0.21 mL of aqueous 30% H₂O₂. The solution was then left to stir at ambient temperature, under N₂, for 3 h. Finally, the reaction was cooled to 0 °C and quenched by slow addition of 2 mL saturated aqueous Na₂S₂O₃. Next, 10 mL of 1 M NaOH was added, and this mixture extracted with EtOAc (2x, ~25 mL). The combined organics were washed with brine (1x, 10 mL) and dried with anhydrous Na₂SO₄. Removal of volatile material under reduced pressure afforded an oil, which was purified by silica gel chromatography (hexanes/EtOAc) to afford the diols as a mixture of diastereomers.

**Representative Procedure for the Tandem Diboration/Imine Allylation with *N*-
(Trimethylsilyl)aldimines (Table 1.4)**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 9.9 mg (0.011 mmol) of tris(dibenzylideneacetone)dipalladium(0), 16.8 mg (0.0258 mmol) of **1.138**, and 0.86 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min; then, 131 mg (0.516 mmol) of B₂(pin)₂ was added followed by 50.0 mg (0.430 mmol) of phenyl allene. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 14 h. After this time period, the vial was taken back into the dry-box, and 114 mg (0.645 mmol) of *N*-(trimethylsilyl)benzaldimine (**1.66**) was weighed in. Next, 28.7 μ L (0.710 mmol) of anhydrous MeOH was added dropwise while stirring the reaction. The vial was capped, removed from the dry-box, and allowed to stir at ambient temperature for 1 h. Next, 0.12 mL (1.3 mmol) of acetic anhydride was added, followed by 0.5 mL of CHCl₃. The vial was purged with N₂, sealed with a cap, and left to stir for an additional 1 h. Volatile material was then removed under reduced pressure, and the resulting residue was diluted with 1.6 mL of THF and 1.3 mL of pH 7 buffer. To this solution was added 0.48 mL of aqueous 30% H₂O₂, and the mixture was allowed to stir for 13 h at ambient temperature under N₂. The final mixture was then transferred to a separatory funnel with 25 mL of CH₂Cl₂ and 10 mL of pH 7 buffer. 2 mL of saturated

aqueous Na₂S₂O₃ was added, and the organic layer was collected. The aqueous phase was extracted with CH₂Cl₂ (2 x 20 mL), and the combined organics were washed with 15 ml of brine, dried with anhydrous Na₂SO₄, and concentrated using reduced pressure. Silica gel chromatography (hexanes/EtOAc) of the mixture afforded 80.7 mg (0.287 mmol, 67 %) of (*R*)-*N*-(3-oxo-1,4-diphenylbutyl)acetamide (**1.136**) as a cream-colored solid.

(*R*)-*N*-(3-Oxo-1,4-diphenylbutyl)acetamide (1.136). A cream-colored solid. Mp 123-126 °C. *R*_f = 0.17 (1:2 hexanes:EtOAc); IR (KBr): 3312 (br, s), 3060 (br, m), 1953 (w), 1871 (w), 1716 (s), 1650 (s), 1534 (s), 1406 (s), 1367 (s) cm⁻¹; ¹H NMR: δ 7.10-7.40 (8H, m, ArH), 7.04 (2H, d, *J* = 6.4 Hz, ArH), 6.63 (1H, br d, *J* = 7.6 Hz, NH), 5.37 (1H, q, *J* = 7.6 Hz, ArCHN), 3.61 (1H, d, *J* = 16 Hz, ArCH_aH_bC(O)), 3.55 (1H, d, *J* = 16 Hz, ArCH_aH_bC(O)) 3.08 (1H, dd, *J* = 16 Hz, *J* = 6.0 Hz, C(O)CH_aH_bCN), 2.89 (1H, dd, *J* = 16 Hz, *J* = 5.4 Hz, C(O)CH_aH_bCN), 1.93 (3H, s, NC(O)CH₃); ¹³C NMR: δ 207.1, 169.3, 140.6, 133.2, 129.4, 128.7, 128.6, 127.4, 127.1, 126.2, 50.7, 49.5, 46.4, 23.3. LRMS (AP+) Calcd for C₁₈H₁₉NO₂ (M + Na)⁺: 304.1, Found (M + Na)⁺: 304.3.


Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/allylation/acetylation/oxidation procedure using tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was determined by X-ray analysis of (*S*)-2-methyl-*N*-((*R*)-3-oxo-1,4-diphenylbutyl)butanamide (**1.161**, see page 87), which was prepared by using (*S*)-(+)-2-methylbutyric anhydride (>99% ee) in place of Ac₂O in the protection step when using phenyl allene and *N*-(trimethylsilyl)benzaldimine (**1.66**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 5.0 % MeOH) analysis of β-amidoketone product 1.136:



Representative Procedure for the Tandem Diboration/Imine Allylation with *In Situ*

Generated Aldimines (Table 1.5)




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 9.9 mg (0.011 mmol) of tris(dibenzylideneacetone)dipalladium(0), 16.8 mg (0.0258 mmol) of **1.138**, and 0.86 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min; then, 131 mg (0.516 mmol) of bis(pinacolato)diboron was added, followed by 50.0 mg (0.430 mmol) of phenyl allene. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 14 h. After this time period, the mixture was taken up in a syringe and added to a solution of benzaldehyde and NH_4OAc in anhydrous MeOH with activated 4 Å molecular sieves that had been stirring under N_2 at room temperature for 2 h (see below for the preparation of this solution). The residue in the vial from the diboration was then rinsed into the aldehyde solution using 0.5 mL of MeOH. This mixture was then allowed to stir for 1 h. Next, 0.41 mL (4.3 mmol) of acetic anhydride was syringed in, and the reaction was allowed to stir for an additional 1 h. The reaction was then poured into 20 mL of Et_2O and filtered over celite until a homogeneous solution was obtained. Volatile material was removed under reduced pressure, and the resulting mixture was then diluted with 1.2 mL of THF and 1.1 mL of pH 7 buffer. To this


solution was added 0.37 mL of aqueous 30% H₂O₂, and it was then allowed to stir at ambient temperature, under N₂, for 13 h. Lastly, it was worked up as described for the *N*-(trimethylsilyl)aldimine procedure (page 56). Silica gel chromatography (hexanes/EtOAc) of the mixture afforded 74.5 mg (0.265 mmol, 62 %) of (*R*)-*N*-(3-oxo-1,4-diphenylbutyl)acetamide (**1.136**) as a cream-colored solid.

Preparation of the aldehyde solution: To a 5-mL RBF with a magnetic stir-bar and activated 4 Å molecular sieves, in a dry-box under an argon atmosphere, was added 249 mg (3.23 mmol) of NH₄OAc. Next, 0.81 mL of anhydrous MeOH was added, followed by 65.5 µL (0.645 mmol) of benzaldehyde. The flask was capped with a septum, removed from the dry-box, and allowed to stir, under N₂, for 2 h before addition of the unpurified diboration mixture.

Determination of Enantiomeric Purity with Ligand **1.138** in the Allene Diboration for Table 1.4 and 1.5



The allene diboration was carried out as described on page 49 using 50.0 mg of allene in 0.86 mL of toluene. Following the same workup procedure, the diboron product was converted to the 1,2-diol, as described on page 54, and analyzed by chiral GLC to determine the enantiomeric purity.





(R)-N-(1-(Furan-2-yl)-3-oxo-4-phenylbutyl)acetamide (1.152).

The product was prepared according to the general allene diboration/imine allylation procedures on pages 57 and 60. A cream-colored solid. Mp 125-127 °C. $R_f = 0.20$ (4:1 $\text{CH}_2\text{Cl}_2:\text{Et}_2\text{O}$); IR (KBr): 3293 (br, s), 3064 (br, m), 2901 (m), 2803 (w), 1953 (w), 1712 (s), 1631 (s), 1542 (s), 1363 (s), 1289 (s) cm^{-1} ; ^1H NMR: δ 7.20-7.45 (4H, m, ArH), 7.15 (2H, d, $J = 9.6$ Hz, ArH), 6.49 (1H, d, $J = 11$ Hz, NH), 6.28 (1H, m, ArH), 6.09 (1H, m, ArH), 5.49 (1H, m, ArCHN), 3.69 (2H, s, ArCH₂C(O)), 3.16 (1H, dd, $J = 22$ Hz, $J = 6.4$ Hz, C(O)CH_aH_bCN), 2.90 (1H, dd, $J = 22$ Hz, $J = 8.0$ Hz, C(O)CH_aH_bCN), 1.97 (3H, s, NC(O)CH₃); ^{13}C NMR: δ 206.7, 169.2, 153.1, 141.7, 133.3, 129.4, 128.8, 127.2, 110.4, 106.2, 50.49, 43.98, 43.83, 23.26. LRMS (AP+) Calcd for $\text{C}_{16}\text{H}_{17}\text{NO}_3$ ($\text{M} + \text{Na}$)⁺: 294.3, Found ($\text{M} + \text{Na}$)⁺: 294.3.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was assumed to be analogous to that obtained for (*R*)-*N*-(3-oxo-1,4-diphenylbutyl)acetamide (**1.136**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 5.0 % MeOH) analysis of β -amidoketone product 1.152:







(*R,E*)-*N*-(2-Oxo-1-phenylnon-5-en-4-yl)acetamide (1.153).

The product was prepared according to the general allene diboration/imine allylation procedure on page 57. An off-white solid. Mp 63-67 °C. R_f = 0.23 (4:1 CH_2Cl_2 : Et_2O); IR (KBr): 3293 (br, s), 2963 (br, m), 1712 (s), 1642 (s), 1541 (s), 1413 (m), 1366 (m) cm^{-1} ; ^1H NMR: δ 7.23-7.45 (3H, m, ArH), 7.18 (2H, d, J = 9.6 Hz, ArH), 6.19 (1H, d, J = 10 Hz, NH), 5.47 (1H, dt, J = 15 Hz, J = 6.4 Hz, vinyl), 5.35 (1H, dd, J = 15 Hz, J = 6.4 Hz, vinyl), 4.73 (1H, m, $\text{CH}=\text{CHCHN}$), 3.69 (2H, s, $\text{ArCH}_2\text{C}(\text{O})$), 2.77 (1H, dd, J = 17 Hz, J = 4.8 Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.69 (1H, dd, J = 17 Hz, J = 5.2 Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 1.75-2.15 (5H, m, allylic CH_2 + $\text{NC}(\text{O})\text{CH}_3$), 1.29 (2H, h, J = 7.4 Hz, CH_2CH_3), 0.81 (3H, t, J = 7.2 Hz, CH_2CH_3); ^{13}C NMR: δ 207.4, 169.1, 133.4, 132.3, 129.37, 128.7, 128.4, 127.1, 50.54, 47.43, 45.88, 34.08, 23.34, 22.03, 13.51. LRMS (AP+) Calcd for $\text{C}_{17}\text{H}_{23}\text{NO}_2$ ($\text{M} + \text{Na}$) $^+$: 296.2, Found ($\text{M} + \text{Na}$) $^+$: 296.3.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was assumed to be analogous to that obtained for (*R*)-*N*-(3-oxo-1,4-diphenylbutyl)acetamide (**1.136**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 4.0 % MeOH) analysis of β -amidoketone product 1.153:





(*R*)-*N*-(3-Oxo-1,6-diphenylhexyl)acetamide (1.154). The

product was prepared according to the general allene diboration/imine allylation procedures on pages 57 and 60. A

white solid. Mp 111-113 °C. R_f = 0.16 (1:2 hexanes:EtOAc); IR (KBr): 3351 (br, s), 3025 (br, m), 2932 (br, m), 1949 (w), 1872 (w), 1712 (s), 1646 (s), 1530 (s), 1367 (s), 1281 (s) cm^{-1} ; ^1H NMR: δ 7.10-7.35 (8H, m, ArH), 7.07 (2H, d, J = 7.6 Hz, ArH), 6.61 (1H, d, J = 7.6 Hz, NH), 5.36 (1H, q, J = 6.0 Hz, ArCHN), 3.05 (1H, dd, J = 16 Hz, J = 5.2 Hz, C(O)CH_aH_bCN), 2.83 (1H, dd, J = 16 Hz, J = 5.6 Hz, C(O)CH_aH_bCN), 2.50 (2H, t, J = 7.2 Hz, ArCH₂), 2.37 (1H, dt, J = 17 Hz, J = 10 Hz, -CH₂CH_aH_bC(O)), 2.26 (1H, dt, J = 17 Hz, J = 7.2 Hz, -CH₂CH_aH_bC(O)), 2.23 (3H, s, NC(O)CH₃), 1.79 (2H, p, J = 7.2 Hz, ArCH₂CH₂); ^{13}C NMR: δ 209.6, 169.3, 141.3, 140.8, 128.6, 128.4, 128.3, 127.4, 126.3, 125.9, 49.54, 47.29, 42.57, 34.74, 24.66, 23.36. LRMS (AP+) Calcd for C₂₀H₂₃NO₂ (M + Na)⁺: 332.2, Found (M + Na)⁺: 332.4.


Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was determined by comparison to authentic material prepared by the following sequence (see page 90 for further details). First, the β -amido methyl ester, (*R*)-methyl 3-acetamido-3-phenyl-propanoate (**1.162**), was prepared using asymmetric hydrogenation as described in the literature.⁵⁸ This compound was then converted to the Weinreb amide using a procedure employed by Toyooka and Nemoto.⁵⁹ Lastly, the amide was converted to the ketone by the addition of (3-phenylpropyl)magnesium bromide using Weinreb's conditions.⁶⁰


⁵⁸ Zhou, Y. G.; Tang, W.; Wang, W., B.; Li, W.; Zhang, X. *J. Am. Chem. Soc.* **2002**, *124*, 4952.

⁵⁹ Toyooka, N.; Okumura, M.; Nemoto, H. *J. Org. Chem.* **2002**, *67*, 6078.

⁶⁰ Nahm, S.; Weinreb, S., M. *Tetrahedron Lett.* **1981**, *22*, 3815.

Chiral SFC (AS-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 4.0 % MeOH) analysis of β -amidoketone product **1.154**:





(R)-N-(1-(Furan-2-yl)-3-oxo-6-phenylhexyl)acetamide

(1.155). The product was prepared according to the general

allene diboration/imine allylation procedures on pages 57 and 60. A cream-colored solid.

Mp 86-90 °C. R_f = 0.22 (4:1 CH₂Cl₂:Et₂O); IR (KBr): 3304 (br, s), 3025 (br, m), 2932

(br, m), 1950 (w), 1872 (w), 1713 (s), 1646 (s), 1522 (s), 1405 (s), 1367 (s), 1297 (s) cm⁻¹;

¹H NMR: δ 7.05-7.30 (6H, m, ArH), 6.54 (1H, d, J = 8.4 Hz, NH), 6.27 (1H, m, ArH),

6.12 (1H, m ArH), 5.44 (1H, m ArCHN), 3.06 (1H, dd, J = 17 Hz, J = 4.8 Hz,

C(O)CH_aH_bCN), 2.81 (1H, dd, J = 17 Hz, J = 6.0 Hz, C(O)CH_aH_bCN), 2.56 (2H, t, J =

7.6 Hz, ArCH₂), 2.38 (2H, m, -CH₂CH₂C(O)), 1.97 (3H, s, NC(O)CH₃), 1.85 (2H, p, J =


6.8 Hz, ArCH₂CH₂); ¹³C NMR: δ 209.0, 169.2, 153.3, 141.6, 141.3, 128.34, 128.30,


125.9, 110.4, 106.2, 44.78, 43.76, 42.24, 34.78, 24.76, 23.18. LRMS (AP+) Calcd for

C₁₈H₂₁NO₃ (M + Na)⁺: 322.1, Found (M + Na)⁺: 322.3.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was assumed to be analogous to that obtained for (*R*)-*N*-(3-oxo-1,6-diphenylhexyl)acetamide (**1.154**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 5.0 % MeOH) analysis of β -amidoketone product 1.155:







(*R,E*)-N-(4-Oxo-1-phenylundec-7-en-6-yl)acetamide

(1.156). The product was prepared according to the general allene diboration/imine allylation procedure on page 57. A white solid. Mp 62-64 °C. R_f = 0.25 (4:1 CH₂Cl₂:Et₂O); IR (KBr): 3293 (br, m), 2947 (br, m), 1708 (s), 1650 (s), 1530 (s), 1370 (s) cm⁻¹; ¹H NMR: δ 7.10-7.30 (5H, m, ArH), 6.28 (1H, d, J = 8.4 Hz, NH), 5.54 (1H, dt, J = 15 Hz, J = 6.8 Hz, vinyl), 5.42 (1H, dd, J = 15 Hz, J = 6.4 Hz, vinyl), 4.69 (1H, m, CH=CHCHN); 2.71 (1H, dd, J = 17 Hz, J = 4.8 Hz, C(O)CH_aH_bCN), 2.62 (1H, dd, J = 17 Hz, J = 5.6 Hz, C(O)CH_aH_bCN), 2.58 (2H, t, J = 7.6 Hz, ArCH₂), 2.39 (2H, m, -CH₂CH₂C(O)), 1.80-2.0 (7H, m, allylic CH₂ + ArCH₂CH₂ + NC(O)CH₃), 1.32 (2H, h, J = 7.2 Hz, CH₂CH₃), 0.85 (3H, t, J = 7.6 Hz, CH₂CH₃); ¹³C NMR: δ 210.2, 169.3, 141.5, 133.6, 128.7, 128.6, 128.5, 126.1, 47.78, 433.6, 128.7, 128.6, 128.5, 126.1, 47.78, 46.86, 42.73, 35.09, 34.32, 25.03, 23.62, 22.30, 13.74. LRMS (AP+) Calcd for C₁₉H₂₇NO₂ (M + Na)⁺: 324.2, Found (M + Na)⁺: 324.4.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was assumed to be analogous to that obtained for (*R*)-*N*-(3-oxo-1,6-diphenylhexyl)acetamide (**1.154**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 4.0 mL/min, 4.0 % MeOH) analysis of β -amidoketone product 1.156:







(R)-N-(4-Cyclohexyl-3-oxo-1-phenylbutyl)acetamide (1.157).

The product was prepared according to the general allene diboration/imine allylation procedures on pages 57 and 60. A white solid. Mp 90-94 °C. $R_f = 0.26$ (4:1 CH_2Cl_2 : Et_2O); IR (KBr): 3324 (br, s), 3064 (br, m), 2912 (br, s), 1949 (w), 1704 (s), 1642 (s), 1537 (s), 1413 (m), 1378 (m) cm^{-1} ; ^1H NMR: δ 7.15-7.40 (5H, m, ArH), 6.74 (1H, d, $J = 7.6$ Hz, NH), 5.36 (1H, m, CHN), 3.04 (1H, dd, $J = 16$ Hz, $J = 5.2$ Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.82 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.16 (2H, m, $\text{CyCH}_2\text{C}(\text{O})$), 1.97 (3H, s, $\text{NC}(\text{O})\text{CH}_3$), 1.42-1.75 (6H, m, C_6H_{11}), 1.0-1.35 (3H, m, C_6H_{11}), 0.70-0.90 (2H, m, C_6H_{11}); ^{13}C NMR: δ 209.9, 169.3, 140.9, 128.6, 127.4, 126.3, 51.28, 49.59, 47.63, 33.52, 33.04, 32.94, 26.04, 25.98, 23.37. LRMS (AP+) Calcd for $\text{C}_{18}\text{H}_{25}\text{NO}_2$ ($\text{M} + \text{Na}$) $^+$: 310.2, Found ($\text{M} + \text{Na}$) $^+$: 310.4.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was determined by comparison to authentic material prepared by the following sequence (see page 92 for further details). First, the β -amido methyl ester, (*R*)-methyl 3-acetamido-3-phenyl-propanoate (**1.162**), was prepared using asymmetric hydrogenation as described in the literature.⁵⁸ This compound was then converted to the Weinreb amide using a procedure employed by Toyooka and Nemoto.⁵⁹ Lastly, the amide was converted to the ketone by the addition of (cyclohexylmethyl)magnesium bromide using Weinreb's conditions.⁶⁰

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 3.0 % MeOH) analysis of β -amidoketone product **1.157**:







(*R,E*)-*N*-(1-Cyclohexyl-2-oxonon-5-en-4-yl)acetamide

(1.158). The product was prepared according to the general allene diboration/imine allylation procedure on page 57. A cream-colored solid. Mp 61-63 °C. R_f = 0.19 (4:1 CH₂Cl₂:Et₂O); IR (KBr): 3324 (br, m), 2920 (br, m), 1704 (s), 1627 (s), 1546 (s), 1370 (s), 1293 (s) cm⁻¹; ¹H NMR: δ 6.32 (1H, d, J = 8.4 Hz, NH), 5.53 (1H, dt, J = 16 Hz, J = 6.4 Hz, vinyl), 5.42 (1H, dd, J = 16 Hz, J = 6.4 Hz, vinyl), 4.68 (1H, m, CH=CHCHN), 2.72 (1H, dd, J = 17 Hz, J = 4.8 Hz, C(O)CH_aH_bCN), 2.64 (1H, dd, J = 17 Hz, J = 5.2 Hz, C(O)CH_aH_bCN), 2.23 (2H, d, J = 6.8 Hz, CyCH₂C(O)), 1.85-1.97 (5H, m, allylic CH₂ + NC(O)CH₃), 1.76 (1H, m, C₆H₁₁), 1.53-1.69 (5H, m, C₆H₁₁), 1.32 (2H, h, J = 7.2 Hz, CH₂CH₃) 1.02-1.26 (3H, m, C₆H₁₁), 0.82 (3H, t, J = 7.2 Hz, CH₂CH₃), 0.75-0.97 (2H, m, C₆H₁₁); ¹³C NMR: δ 210.3, 169.1, 132.3, 128.6, 51.17, 47.63, 47.12, 34.14, 33.74, 33.15, 33.04, 26.08, 25.99, 23.45, 22.11, 13.56. LRMS (AP+) Calcd for C₁₇H₂₉NO₂ (M + Na)⁺: 302.2, Found (M + Na)⁺: 302.4.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was determined by X-ray analysis of (*S*)-*N*-((*R*, *E*)-1-cyclohexyl-2-oxonon-5-en-4-yl)-2-methylbutanamide (**1.166**, see page 93), which was prepared by using (*S*)-(+)-2-methylbutyric anhydride (>99% ee) in place of Ac₂O in the protection step.

Chiral HPLC (Chiralcel-OD-H, Daicel, 3.0 % iPrOH in hexanes, 0.9 mL/min) analysis of β -amidoketone product 1.158:






(R)-N-(4-Cyclohexyl-1-(furan-2-yl)-3-oxobutyl)acetamide

(1.159). The product was prepared according to the general allene


diboration/imine allylation procedures on pages 57 and 60. A white solid. Mp 118-120 °C. $R_f = 0.26$ (4:1 $\text{CH}_2\text{Cl}_2:\text{Et}_2\text{O}$); IR (KBr): 3304 (br, s), 3064 (br, m), 2912 (br, s), 1693 (s), 1642 (s), 1537 (s), 1370 (s), 1285 (s) cm^{-1} ; ^1H NMR: δ 7.25 (1H, s, ArH), 6.68 (1H, d, $J = 8.4$ Hz, NH), 6.27 (1H, dd, $J = 4.0$ Hz, $J = 2.6$ Hz, ArH), 6.12 (1H, d, $J = 4.0$ Hz, ArH), 5.44 (1H, m, CHN), 3.06 (1H, dd, $J = 17$ Hz, $J = 4.8$ Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.80 (1H, dd, $J = 17$ Hz, $J = 6.0$ Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.23 (2H, d, $J = 6.9$ Hz, $\text{CyCH}_2\text{C}(\text{O})$), 1.96 (3H, s, $\text{NC}(\text{O})\text{CH}_3$), 1.50-1.85 (6H, m, C_6H_{11}), 1.0-1.35 (3H, m, C_6H_{11}), 0.75-1.0 (2H, m, C_6H_{11}); ^{13}C NMR: δ 209.3, 169.2, 153.4, 141.5, 110.4, 106.2, 50.95, 45.12, 43.83, 33.57, 33.02, 32.95, 26.04, 25.97, 23.22. LRMS (AP+) Calcd for $\text{C}_{16}\text{H}_{23}\text{NO}_3$ ($\text{M} + \text{Na}$) $^+$: 300.2, Found ($\text{M} + \text{Na}$) $^+$: 300.3.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material prepared by performing the diboration/imine allylation with tricyclohexylphosphine as the achiral ligand in the diboration step. Absolute stereochemistry was assumed to be analogous to that obtained for (*R*)-*N*-(4-cyclohexyl-3-oxo-1-phenylbutyl)acetamide (**1.157**) and (*R,E*)-*N*-(1-cyclohexyl-2-oxonon-5-en-4-yl)acetamide (**1.158**).

Chiral SFC (AS-H, Chiralpak, 150 bar, 50°C, flow = 3.0 mL/min, 3.0 % MeOH) analysis of β -amidoketone product 1.159:




Synthesis of Vinylboronate Ester **1.141**



The diboration-allylation sequence was carried out as described above when using silylaldimines (page 57) using 50.0 mg (0.430 mmol) of phenyl allene and 114 mg (0.645 mmol) of *N*-(trimethylsilyl)benzaldimine. After this was complete, 3.0 equiv of (Boc)₂O was added, followed by 0.5 mL of CHCl₃. This mixture was allowed to stir for 1 h at ambient temperature. Volatile material was then removed under reduced pressure, and silica gel chromatography (hexanes/EtOAc) of the mixture afforded 114 mg (0.254 mmol, 59%) of **1.141** as a white foam.

(*R,Z*)-tert-Butyl-1,4-diphenyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)but-3-enylcarbamate (1.141**).** *R_f* = 0.24 (CH₂Cl₂); IR (KBr): 3390 (br, m), 2975 (br, m), 1950 (w), 1717 (s), 1611 (m), 1514 (s), 1382 (s), 1344 (s), 1262 (s), 1165 (s) cm⁻¹; ¹H NMR: δ 7.41 (1H, s, vinyl **CH**), 7.12-7.38 (10H, m, Ar**H**), 5.81 (1H, br d, *J* = 5.2 Hz, **NH**), 4.71 (1H, br m, Ar**CHN**), 2.77 (1H, m, C(B)**CH_aH_bCN**), 2.54 (1H, m, C(B)**CH_aH_bCN**), 1.37 (9H, s, OC(**CH₃**)₃), 1.35 (6H, s, C(**CH₃**)₂), 1.33 (6H, s, C(**CH₃**)₂); ¹³C NMR: δ 155.3, 145.7, 144.2, 137.3, 128.9, 128.3, 128.2, 127.5, 126.7, 125.8, 84.01, 78.86, 56.33, 36.48, 28.40, 24.98, 24.68. LRMS (ES⁺) Calcd for C₂₇H₃₆BNO₄ (M + Na)⁺: 472.3, Found (M + Na)⁺: 472.3.

Synthesis of Suzuki-Miyaura Coupling Product 1.142




To 66.9 mg (0.149 mmol) of **1.141**, in a 20-mL scintillation vial with a magnetic stir-bar, was weighed 6.1 mg (0.0075 mmol) of [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane complex. The vial was capped with a septum and purged with N₂. To the vial was added 1.06 mL of dioxane, followed by 25.1 μ L (0.224 mmol) of iodobenzene, and lastly 0.11 mL of 4 M KOH. The septum was then quickly replaced with a polypropylene cap; the vial was sealed with electrical tape and heated at 80 °C for 14 h. After this time period, the mixture was transferred to a separatory funnel with 20 mL of CH₂Cl₂ and 10 mL of H₂O. The organic layer was collected, and the aqueous layer was extracted with CH₂Cl₂ (1 x 20 mL). The combined organics were washed with 15 mL of H₂O, 10 mL of brine, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure. Purification by silica gel chromatography (9:1 hexanes:EtOAc) afforded 37.2 mg (0.0931 mmol, 62 %) of (*R,E*)-*tert*-butyl-1,3,4-triphenylbut-3-enylcarbamate (**1.142**) as a waxy solid.

(*R,E*)-*tert*-Butyl-1,3,4-triphenylbut-3-enylcarbamate (1.142). R_f = 0.18 (9:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3437 (br, m), 2966 (br, m), 1953 (w), 1705 (s), 1495 (s), 1363 (s), 1246 (m), 1165 (s) cm⁻¹; ¹H NMR: δ 7.45 (2H, d, J = 7.2 Hz, ArH), 7.40 (2H, t, J = 7.2 Hz, ArH), 7.33 (3H, t, J = 7.6 Hz, ArH), 7.17-7.29 (4H, m, ArH), 7.11 (2H, d, J = 7.6 Hz, ArH), 7.00 (2H, dd, J = 7.6 Hz, J = 2.0 Hz, ArH), 6.75 (1H, s,


vinyl CH), 4.65 (2H, br s, NH + ArCHN), 3.19 (1H, m, ArCHNCH_aH_b), 3.07 (1H, m, ArCHNCH_aH_b), 1.37 (9H, s, OC(CH₃)₃); ¹³C NMR: δ 154.8, 142.1, 139.4, 137.7, 131.2, 128.8, 128.5, 128.3, 128.2, 127.4, 127.1, 126.8, 126.7, 126.2, 53.85, 37.26, 28.28. LRMS (ES+) Calcd for C₂₇H₂₉NO₂ (M + Na)⁺: 422.2, Found (M + Na)⁺: 422.4.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material, which was prepared using racemic starting material that had been synthesized as described above with tricyclohexylphosphine in place of the chiral ligand in the diboration step. Absolute configuration was assumed to be analogous to that obtained for (*R*)-*N*-(3-oxo-1,4-diphenylbutyl)acetamide (**1.136**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 0.5 mL/min, 5.0 % MeOH) analysis of the Suzuki-coupling product 1.142:




Synthesis of Vinylboronate Ester **1.143**



The diboration-allylation sequence was carried out as described above when using silylaldimines (page 57) using 50.0 mg (0.430 mmol) of phenyl allene and 114 mg (0.645 mmol) of *N*-(trimethylsilyl)benzaldimine. After this was complete, 3.0 equiv of the propionic anhydride was added, followed by 0.5 mL of CHCl_3 . This mixture was allowed to stir for 1 h at ambient temperature. Volatile material was then removed under reduced pressure, and silica gel chromatography (hexanes/EtOAc) of the mixture afforded 108 mg (0.266 mmol, 62%) of **1.143** as a waxy solid.

(*R,Z*)-*N*-(1,4-Diphenyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)but-3-enyl)propionamide (1.143). R_f = 0.27 (10:1 CH_2Cl_2 : Et_2O); IR (KBr): 3293 (br, s), 3060 (br, m), 2970 (br, m), 1949 (w), 1642 (s), 1600 (s), 1537 (s), 1355 (s), 1142 (s) cm^{-1} ; ^1H NMR: δ 7.10-7.48 (11H, m, ArH + vinyl CH), 6.36 (1H, d, J = 6.8 Hz, NH), 5.05 (1H, m, ArCHN), 2.82 (1H, m, C(B)CH_aH_bCN), 2.73 (1H, m, C(B)CH_aH_bCN), 2.12 (2H, q, J = 7.6 Hz, NC(O)CH₂CH₃), 1.32 (6H, s, C(CH₃)₂), 1.30 (6H, s, C(CH₃)₂), 1.08 (3H, t, J = 7.6 Hz, NC(O)CH₂CH₃); ^{13}C NMR: δ 172.9, 145.9, 143.2, 137.3, 128.9, 128.33, 128.31, 127.5, 126.8, 126.0, 83.94, 54.33, 35.87, 29.63, 25.18, 24.51, 9.78. LRMS (AP+) Calcd for $\text{C}_{25}\text{H}_{32}\text{BNO}_3$ ($\text{M} + \text{Na}$)⁺: 428.2, Found ($\text{M} + \text{Na}$)⁺: 428.5.

Synthesis of Protonation Product 1.144




To 85.2 mg (0.210 mmol) of **1.143**, in a 20-mL scintillation vial with a magnetic stir-bar, was added 2.1 mL of propionic acid. The vial was purged with N₂, capped with a polypropylene cap, sealed with electrical tape, and heated at 140 °C for 28 h. After this time period, the mixture was transferred to a separatory funnel with 20 mL of Et₂O and extracted with saturated aqueous NaHCO₃ (3 x 15 mL). The combined aqueous layers were then extracted with Et₂O (1 x 20 mL). The combined organics were washed with 10 mL of brine, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure to afford a brown solid. Purification using silica gel chromatography (20:1 CH₂Cl₂:Et₂O) afforded 54.2 mg (0.194 mmol, 92 %) of (*R,Z*)-*N*-(1,4-diphenylbut-3-enyl)propionamide (**1.144**) as a white solid.

(*R,Z*)-*N*-(1,4-Diphenylbut-3-enyl)propionamide (1.144). R_f = 0.22 (20:1 CH₂Cl₂:Et₂O); IR (KBr): 3328 (br, m), 3017 (br, m), 2940 (br, m), 1953 (w), 1884 (w), 1802 (w), 1642 (s), 1530 (s), 1452 (s), 1383 (s), 1274 (s), 1102 (s) cm⁻¹; ¹H NMR: δ 7.15-7.39 (10H, m, ArH), 6.51 (1H, d, J = 11.6 Hz, ArCH=CH), 5.71 (1H, d, J = 8.0 Hz, NH), 5.58 (1H, m, ArCH=CHCH₂), 5.13 (1H, q, J = 7.6 Hz, ArCHN), 2.84 (2H, t, J = 6.8 Hz, ArCH=CHCH₂), 2.15 (2H, q, J = 7.6 Hz, CH₂CH₃), 1.10 (3H, t, J = 7.6 Hz, CH₂CH₃); ¹³C NMR: δ 173.0, 141.63, 137.0, 131.5, 128.7, 128.6, 128.3, 127.7, 127.4,


126.9, 126.5, 52.97, 34.93, 29.69, 9.76. LRMS (ES⁺) Calcd for C₁₉H₂₁NO (M + Na)⁺: 302.2, Found (M + Na)⁺: 302.3.

Proof of stereochemistry. Stereochemical ratios were determined in comparison with authentic racemic material, which was prepared using racemic starting material that had been synthesized as described above with tricyclohexylphosphine in place of the chiral ligand in the diboration step. Absolute configuration was assumed to be analogous to that obtained for (*R*)-*N*-(3-oxo-1,4-diphenylbutyl)acetamide (**1.136**).

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 4.0 mL/min, 3.0 % MeOH) analysis of the protonation product 1.144:



Synthesis of Aminoboronate 1.145




An oven-dried 2-dram vial equipped with a magnetic stir-bar was charged with 9.9 mg (0.011 mmol) of tris(dibenzylideneacetone)dipalladium(0), 7.2 mg (0.0258 mmol) of PCy_3 , and 2.7 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min; then, 131 mg (0.516 mmol) of bis(pinacolato)diboron was added followed by 50.0 mg (0.430 mmol) of phenyl allene. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 14 h. After this time period, the vial was taken back into the dry-box, and 76.2 mg (0.430 mmol) of *N*-(trimethylsilyl)benzaldimine was weighed in. Next, 20.9 μL (0.516 mmol) of MeOH was added dropwise while stirring the reaction. A white precipitate formed after 1 h at ambient temperature, and the mixture was then poured into 15 mL of hexanes. The mixture was filtered using vacuum filtration, and 55 mg (0.16 mmol, 37%) of an off-white solid was collected. Recrystallization from a mixture of hexanes and EtOAc afforded X-ray quality crystals. See page 95 for crystal parameters.

(*Z*)-1,4-Diphenyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)but-3-en-1-amine

(1.145). Mp 170-173 °C. IR (KBr): 3309 (br, m), 3196 (br, m), 3067 (br, s), 2971 (br, s), 1945 (w), 1884 (w), 1806 (w), 1596 (s), 1495 (s), 1460 (s), 1383 (s), 1212 (s), 1150 (s)

cm⁻¹; ¹H NMR: δ 7.24-7.50 (5H, m, ArH), 7.11 (5H, s, ArCH), 6.85 (1H, s, vinyl CH), 3.5-5.0 (2H, br s, NH₂), 4.09 (1H, m, ArCHN), 2.93 (1H, m, ArCH(N)CH_aH_b), 2.83 (1H, m, ArCH(N)CH_aH_b), 1.20 (6H, s, C(CH₃)₂), 1.17 (6H, s, C(CH₃)₂); ¹³C NMR: δ 140.9, 139.3, 131.6, 128.7, 128.6, 128.1, 127.9, 126.9, 125.9, 80.25, 57.26, 38.84, 25.49. LRMS (ES+) Calcd for C₂₂H₂₈BNO₂ (M + H)⁺: 350.2, Found (M + H)⁺: 350.4.


Procedure for Absolute Stereochemistry Determination of **1.136** by X-ray Analysis




The diboration/allylation was carried out as described above when using silylaldimines using 69.7 mg (0.600 mmol) of phenyl allene and 70.8 mg (0.400 mmol) of *N*-(trimethylsilyl)benzalimine (**1.66**). After the allylation step was complete, 0.24 mL (1.2 mmol) of (*S*)-(+)-2-methylbutyric anhydride (**1.160**, >98% ee and commercially available from Aldrich Chemical Co.) was added in place of Ac₂O in the dry-box. After oxidation and the aqueous workup, silica gel column chromatography (30:1 CH₂Cl₂:Et₂O) afforded 71 mg (0.33 mmol, 55%) of (*S*)-2-methyl-*N*-((*R*)-3-oxo-1,4-diphenylbutyl)butanamide (**1.161**) as a white solid. Recrystallization using a mixture of hexanes and EtOAc afforded X-ray quality crystals.

(S)-2-Methyl-N-[(R)-3-oxo-1,4-diphenylbutyl]butanamide (1.161). Mp 115-118 °C. $R_f = 0.19$ (2:1 hexanes:EtOAc); IR (KBr): 3328 (br, m), 3029 (br, m), 2928 (br, m), 1945 (w), 1871 (w), 1706 (s), 1639 (s), 1526 (s), 1491 (s), 1449 (s), 1238 (s), 1172 (s) cm^{-1} ; ^1H NMR: δ 7.20-7.32 (6H, m, ArH), 7.16 (2H, d, $J = 7.2$ Hz, ArH), 7.04 (2H, d, $J = 6.8$ Hz, ArH), 6.57 (1H, d, $J = 7.6$ Hz, NH), 5.40 (1H, m, ArCHN), 3.62 (1H, d, $J = 16$ Hz, ArCH_aH_bC(O)), 3.56 (1H, d, $J = 16$ Hz, ArCH_aH_bC(O)), 3.10 (1H, dd, $J = 16$ Hz, $J = 5.6$ Hz, C(O)CH_aH_bCN), 2.89 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz, C(O)CH_aH_bCN), 2.09 (1H, h, $J = 7.2$ Hz, NC(O)CH), 1.63 (1H, m, NC(O)CHCH_aH_bCH₃), 1.40 (1H, m, NC(O)CHCH_aH_bCH₃), 1.08 (3H, d, $J = 6.8$ Hz, CHCH₃), 0.86 (3H, t, $J = 7.6$ Hz, CH₂CH₃); ^{13}C NMR: δ 207.2, 175.7, 140.9, 133.3, 129.4, 128.7, 128.6, 127.3, 127.1, 126.1, 50.69, 49.13, 46.47, 43.07, 27.18, 17.30, 11.82. LRMS (ES+) Calcd for C₂₁H₂₅NO₂ (M + Na)⁺: 346.2, Found (M + Na)⁺: 346.3.

Figure 1.6: Crystal Structure of 1.161



Procedure for Absolute Stereochemistry Determination of **1.154**



A 50 mL RBF with magnetic stir-bar was charged with 445 mg (10.6 mmol) of $\text{LiOH}\cdot\text{H}_2\text{O}$, followed by 1.17 g (5.29 mmol) of **1.162**,⁶¹ and lastly 12.6 mL of a 3:1 $\text{MeOH}:\text{H}_2\text{O}$ mixture. The flask was purged with N_2 , fitted with a reflux condenser, and heated at $80\text{ }^\circ\text{C}$ for 1 h. After cooling to room temperature, the reaction was acidified with 40 mL of 1 M HCl and extracted with EtOAc (2x, ~50 mL). The combined organics were dried with anhydrous Na_2SO_4 and concentrated into a 100 mL RBF. A magnetic stir-bar was added, and the flask purged with N_2 . Next, 36 mL of CH_2Cl_2 was added by syringe, and the mixture was cooled to $0\text{ }^\circ\text{C}$. 1.12 g (6.88 mmol) of CDI was added, the ice bath was removed, and the reaction was allowed to stir at ambient temperature for 30 min. The mixture was re-cooled to $0\text{ }^\circ\text{C}$ and 671 mg (6.88 mmol) of (*N,O*)-dimethylhydroxylamine hydrochloride was added, followed by 0.96 mL (6.9 mmol) of TEA . The ice bath was removed, and the mixture was allowed to stir at ambient temperature for 24 h. Next, 3.5 mL of Ac_2O was added and stirring was continued for an additional 1 h. The reaction was transferred to a separatory funnel and washed sequentially with saturated aqueous NaHCO_3 (2x), 1 M HCl (2x), and brine (1x). The organic layer was dried with anhydrous Na_2SO_4 , and volatile material was removed under


⁶¹ **1.162** was prepared in 52% yield and 18% ee using (*R*)-BINAP according to reference 58.

reduced pressure. Purification of the mixture using silica gel chromatography (5% MeOH in EtOAc) afforded 873 mg (3.49 mmol, 66%) of **1.163** as a white solid.

(R)-3-Acetamido-N-methoxy-N-methyl-3-phenylpropanamide (1.163). $R_f = 0.19$ (5% MeOH in EtOAc); IR (KBr): 3352 (br, s), 3033 (br, m), 2940 (br, m), 1957 (w), 1876 (w), 1670 (s), 1631 (s), 1534 (s), 1460 (s), 1367 (s), 1293 (s), 1204 (s) cm^{-1} ; ^1H NMR: δ 7.41 (1H, d, $J = 6.8$ Hz, **NH**), 7.15-7.40 (5H, m, **ArH**), 5.39 (1H, m, **ArCHN**), 3.44 (3H, s, **CH₃**), 3.12 (1H, dd, $J = 16$ Hz, $J = 5.0$ Hz, **ArCH(N)CH_aH_b**), 3.06 (3H, s, **CH₃**), 2.77 (1H, dd, $J = 16$ Hz, $J = 5.0$ Hz, **ArCH(N)CH_aH_b**), 1.99 (3H, s, **NC(O)CH₃**); ^{13}C NMR: δ 171.9, 169.3, 141.3, 128.5, 127.2, 126.2, 61.16, 49.66, 36.65, 31.73, 23.36. LRMS (ES+) Calcd for $\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_3$ ($\text{M} + \text{Na}$) $^+$: 273.1, Found ($\text{M} + \text{Na}$) $^+$: 273.2.


To 38.1 mg (0.152 mmol) of **1.163** in 1.5 mL of THF at 0 °C was added 0.60 mL of a 0.78 M solution of (3-phenylpropyl)magnesium bromide in THF dropwise. This mixture was allowed to stir for 2 h at 0 °C and then, at ambient temperature for 20 min. The reaction was quenched by the addition of 3 mL of saturated aqueous NH_4Cl . Water was added, and the mixture extracted with CH_2Cl_2 . The combined organics were washed with 1 M HCl (2x), brine, and dried with anhydrous Na_2SO_4 . After removal of volatile material under reduced pressure, silica gel chromatography (hexanes/EtOAc) of the mixture afforded 21.1 mg (0.0682 mmol, 45%) of **1.154**. Spectral data were in good agreement with **1.154** prepared by the tandem diboration/allylation sequence.

Procedure for Absolute Stereochemistry Determination of **1.157**



To 48.0 mg (0.192 mmol) of **1.163** in 1.9 mL of THF at 0 °C was added 0.89 mL of a 0.67 M solution of (cyclohexylmethyl)magnesium bromide in THF dropwise. This mixture allowed to reach ambient temperature and allowed to stir overnight. The reaction was quenched by the addition of 3 mL of saturated aqueous NH₄Cl. Water was added, and the mixture extracted with CH₂Cl₂. The combined organics were washed with 1 M HCl (2x), brine, and dried with anhydrous Na₂SO₄. After removal of volatile material under reduced pressure, silica gel chromatography (hexanes/EtOAc) of the mixture afforded 3.3 mg (0.012 mmol, 6%) of **1.157**. Spectral data were in good agreement with **1.157** prepared by the tandem diboration/allylation sequence.

Procedure for Absolute Stereochemistry Determination of **1.158** by X-ray Analysis




The diboration/allylation was carried out as described above when using silylaldimines (page 57) using 50.0 mg (0.409 mmol) of cyclohexyl allene and 104 mg (0.614 mmol) of silylaldimine **1.165**. After the allylation step was complete, 0.24 mL (1.2 mmol) of (S)-(+)-2-methylbutyric anhydride (**1.160**, >98% ee and commercially available from Aldrich Chemical Co.) was added in place of Ac_2O in the dry-box. After oxidation and the aqueous workup, silica gel column chromatography (30:1 $\text{CH}_2\text{Cl}_2:\text{Et}_2\text{O}$) afforded 81.7 mg (0.254 mmol, 62 %) of (S)-N-((R, E)-1-cyclohexyl-2-oxonon-5-en-4-yl)-2-methylbutanamide (**1.166**) as a white solid. Recrystallization using a mixture of hexanes and EtOAc afforded X-ray quality crystals.

(S)-N-[(R,E)-1-Cyclohexyl-2-oxonon-5-en-4-yl]-2-methylbutanamide (1.166). Mp 92-96 °C. R_f = 0.19 (30:1 $\text{CH}_2\text{Cl}_2:\text{Et}_2\text{O}$); IR (KBr): 3393 (br, s), 2916 (br, s), 1709 (s), 1631 (s), 1538 (s), 1383 (m), 1281 (m) cm^{-1} ; ^1H NMR: δ 6.26 (1H, d, J = 8.4 Hz, NH), 5.52 (1H, dt, J = 16 Hz, J = 6.4 Hz, vinyl), 5.43 (1H, dd, J = 16 Hz, J = 6.0 Hz, vinyl), 4.70 (1H, m, $\text{CH}=\text{CHCHN}$), 2.73 (1H, dd, J = 17 Hz, J = 4.8 Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.63 (1H, dd, J = 17 Hz, J = 5.6 Hz, $\text{C}(\text{O})\text{CH}_a\text{H}_b\text{CN}$), 2.23 (2H, d, J = 6.8 Hz, $\text{CyCH}_2\text{C}(\text{O})$), 2.04 (1H, h, J = 6.8 Hz, $\text{NC}(\text{O})\text{CH}(\text{CH}_3)\text{CH}_2-$), 1.93 (2H, q, J = 7.2 Hz, allylic CH_2), 1.54-

1.86 (7H, m, aliphatic), 1.03-1.45 (7H, m, aliphatic), 1.07 (3H, d, $J = 6.8$ Hz, NC(O)CH(CH₃)CH₂-), 0.77-0.96 (7H, m, aliphatic); ¹³C NMR: δ 210.3, 175.4, 132.2, 128.8, 51.15, 47.30, 47.26, 43.19, 34.16, 33.78, 33.15, 33.06, 27.25, 26.10, 26.00, 21.15, 17.35, 13.54, 11.81. LRMS (ES+) Calcd for C₂₀H₃₅NO₂ (M + Na)⁺: 344.3, Found (M + Na)⁺: 344.2.

Figure 1.7: Crystal Structure of 1.166



X-ray Data for 1.145

Space Group and Cell Dimensions Monoclinic, P 2₁/c
a 12.6666(10) b 19.3167(15) c 8.4376(6)
beta 105.406(6)
Volume 1990.3(3)Å³

Empirical formula : C₂₂ H₂₈ B N O₂

Cell dimensions were obtained from 732 reflections with
2Theta angle in the range 5.00 - 50.00 degrees.

Crystal dimensions : 0.30 X 0.02 X 0.02 mm

FW = 349.28 Z = 4 F(000) = 752.40

Dcalc 1.166Mg.m⁻³, mu 0.07mm⁻¹, lambda 0.71073Å,
2Theta(max) 50.0

The intensity data were collected on a Bruker SMART 1K
diffractometer, using the omega scan mode.

The h,k,l ranges used during structure solution and
refinement are :--Hmin,max -15 14; Kmin,max 0 22;
Lmin,max 0 10

No. of reflections measured 11886

No. of unique reflections 3494

No. of reflections with Inet > 2.5sigma(Inet) 1595

Merging R-value on intensities 0.037

Correction was made for absorption using SADABS

Details of the last least squares cycle
54 atoms, 235 parameters Full-matrix on Fo Counter wts
(k 0.000400)

The residuals are as follows :--

Significant reflections: 1590 Rf 0.071, Rw 0.072

All reflections: 3494 Rf 0.161, Rw 0.084

Included reflections: 1590 Rf 0.071, Rw 0.072

GoF 2.0295

where Rf = Sum(Fo-Fc)/Sum(Fo),

Rw = Sqrt[Sum(w(Fo-Fc)**2)/Sum(wFo**2)] and

GoF = Sqrt[Sum(w(Fo-Fc)**2)/(No. of reflns - No.

of
params.)]

The maximum shift/sigma ratio was 0.000.

Last D-map: Minimum density -0.310e/A**3,
Maximum density 0.430e/A**3.

Table 1. Atomic Parameters x,y,z and Biso.
E.S.Ds. refer to the last digit printed.

	x	y	z	Biso
B1	0.8037 (5)	0.2052 (3)	0.1942(7)	3.2 (3)
O2	0.85974(25)	0.17727(15)	0.0785(3)	3.30(15)
C3	0.9617 (4)	0.1482 (3)	0.1705(6)	3.9 (3)
C4	1.0499 (4)	0.2050 (3)	0.2071(7)	5.9 (3)
C5	0.9978 (5)	0.0926 (3)	0.0699(7)	5.9 (3)
C6	0.9361 (4)	0.1241 (3)	0.3282(6)	4.2 (3)
C7	1.0314 (5)	0.1169 (3)	0.4767(7)	6.3 (3)
C8	0.8732 (5)	0.0546 (3)	0.3015(7)	6.4 (4)
O9	0.8622 (3)	0.17714(16)	0.3539(3)	4.13(17)
C10	0.6724 (4)	0.2050 (3)	0.1409(5)	3.4 (3)
C11	0.6053 (4)	0.1512 (3)	0.0922(5)	3.4 (3)
C12	0.4856 (5)	0.1509 (3)	0.0330(6)	3.6 (3)
C13	0.4335 (5)	0.1100 (3)	-0.1000(7)	4.9 (3)
C14	0.3213 (6)	0.1123 (4)	-0.1668(7)	6.3 (4)
C15	0.2579 (5)	0.1555 (4)	-0.1006(10)	6.6 (4)
C16	0.3067 (6)	0.1946 (3)	0.0340(9)	5.9 (4)
C17	0.4180 (5)	0.1930 (3)	0.1006(7)	5.0 (3)
C18	0.6312 (4)	0.2789 (3)	0.1418(7)	5.8 (3)
C19	0.7165 (5)	0.3237 (3)	0.1198(7)	5.8 (3)
C20	0.7019 (4)	0.4005 (3)	0.1391(6)	3.9 (3)
C21	0.7276 (4)	0.4332 (3)	0.2881(6)	4.3 (3)
C22	0.7101 (5)	0.5028 (4)	0.3003(8)	5.9 (4)
C23	0.6637 (6)	0.5404 (3)	0.1660(11)	6.8 (5)
C24	0.6347 (6)	0.5096 (4)	0.0181(9)	8.1 (5)
C25	0.6533 (6)	0.4391 (4)	0.0022(7)	7.0 (4)
N26	0.8204 (3)	0.29206(19)	0.2056(4)	3.65(20)
H4a	1.049	0.229	0.107	6.8
H4b	1.121	0.185	0.251	6.8
H4c	1.035	0.237	0.285	6.8
H5a	0.941	0.058	0.039	6.7
H5b	1.064	0.071	0.134	6.7
H5c	1.011	0.113	-0.027	6.7
H7a	1.067	0.161	0.502	6.8
H7b	1.082	0.084	0.455	6.8

H7c	1.006	0.102	0.569	6.8
H8a	0.846	0.045	0.395	7.2
H8b	0.922	0.018	0.289	7.2
H8c	0.813	0.057	0.205	7.2
H11	0.640	0.107	0.097	4.2
H13	0.476	0.079	-0.146	5.8
H14	0.288	0.084	-0.260	7.0
H15	0.180	0.158	-0.148	7.4
H16	0.262	0.224	0.082	7.0
H17	0.450	0.221	0.195	6.1
H18a	0.564	0.285	0.058	6.9
H18b	0.619	0.289	0.247	6.9
H19	0.713	0.318	0.005	6.4
H21	0.760	0.407	0.386	5.2
H22	0.731	0.525	0.406	6.7
H23	0.651	0.589	0.175	7.9
H24	0.600	0.536	-0.078	9.3
H25	0.634	0.418	-0.104	7.9
H26a	0.839	0.307	0.318	4.6
H26b	0.878	0.306	0.158	4.6

Biso is the Mean of the Principal Axes of the Thermal Ellipsoid

Table of $u(i,j)$ or U values *100.
E.S.Ds. refer to the last digit printed

	u11(U)	u22	u33	u12	u13	u23
B1	5.3(4)	3.2(4)	3.7(3)	0.0(3)	1.4(3)	0.2(3)
O2	4.89(22)	4.01(21)	4.10(18)	1.38(17)	2.00(17)	0.46(16)
C3	4.4(4)	5.0(4)	5.5(3)	1.4(3)	1.5(3)	0.3(3)
C4	6.3(4)	6.3(4)	10.1(5)	1.0(3)	2.9(4)	1.1(4)
C5	7.4(4)	6.6(4)	8.3(4)	3.1(3)	2.0(3)	-0.5(3)
C6	6.6(4)	4.0(4)	4.7(3)	1.9(3)	0.4(3)	0.7(3)
C7	10.0(5)	5.7(4)	7.4(4)	1.5(4)	0.7(4)	1.2(3)
C8	9.1(5)	4.7(4)	10.6(5)	0.9(3)	3.1(4)	1.8(3)
O9	7.2(3)	4.50(23)	4.32(20)	1.65(20)	2.17(18)	0.83(17)
C10	5.2(4)	3.4(3)	5.1(3)	0.6(3)	2.5(3)	0.7(3)
C11	4.8(4)	4.2(3)	3.9(3)	0.6(3)	1.3(3)	0.15(25)
C12	5.1(4)	5.0(4)	4.2(3)	-0.4(3)	2.1(3)	0.3(3)
C13	5.2(5)	7.6(5)	6.4(4)	-0.4(3)	2.2(3)	-1.1(3)
C14	6.1(5)	9.4(6)	8.0(4)	-0.9(4)	1.2(4)	-1.3(4)

C15	5.2(5)	8.7(6)	11.1(6)	-0.3(4)	2.2(4)	1.0(5)
C16	5.6(5)	7.3(5)	10.8(5)	0.5(4)	4.2(4)	0.0(4)
C17	6.1(4)	5.8(4)	8.2(4)	-0.5(3)	4.1(4)	-0.9(3)
C18	5.8(4)	4.8(4)	12.4(5)	1.3(3)	4.4(4)	2.0(4)
C19	7.4(5)	4.4(4)	9.5(5)	1.7(4)	1.0(4)	-0.1(3)
C20	6.3(4)	4.0(4)	5.2(4)	1.1(3)	2.6(3)	-0.2(3)
C21	6.7(4)	4.2(4)	5.7(4)	-0.2(3)	2.2(3)	0.4(3)
C22	8.1(5)	5.4(5)	9.0(5)	0.0(4)	2.7(4)	-1.9(4)
C23	9.5(6)	4.3(5)	13.4(7)	1.2(4)	5.5(5)	0.2(5)
C24	15.3(7)	7.5(6)	9.7(6)	6.5(5)	6.1(5)	4.7(5)
C25	13.6(6)	9.1(5)	4.4(4)	5.0(5)	3.0(4)	0.0(4)
N26	4.5(3)	3.7(3)	6.2(3)	0.95(21)	2.32(22)	0.43(21)
H4a	8.6					
H4b	8.6					
H4c	8.6					
H5a	8.4					
H5b	8.4					
H5c	8.4					
H7a	8.7					
H7b	8.7					
H7c	8.7					
H8a	9.1					
H8b	9.1					
H8c	9.1					
H11	5.3					
H13	7.4					
H14	8.8					
H15	9.4					
H16	8.9					
H17	7.7					
H18a	8.7					
H18b	8.7					
H19	8.1					
H21	6.6					
H22	8.5					
H23	10.1					
H24	11.8					
H25	10.0					
H26a	5.8					
H26b	5.8					

Anisotropic Temperature Factors are of the form

$$\text{Temp} = -2\pi^2 (h^2 u_{11}^* a^{*2} + \dots + 2hk u_{12}^* a^{*2} b^{*2} + \dots)$$

DISANG -- The NRCVAX Distance and Angle Program

The Space Group is P 2₁/C Centrosymmetric
The Equivalent Positions are:

1) x y z 2) -x 1/2+y 1/2-z

The Lattice is Primitive. There are no Centering Vectors

Bond lengths (Å):

B(1)-O(2)	1.454(6)	C(12)-C(17)	1.407(8)
B(1)-O(9)	1.459(6)	C(13)-C(14)	1.384(9)
B(1)-C(10)	1.604(8)	C(14)-C(15)	1.375(11)
B(1)-N(26)	1.691(7)	C(15)-C(16)	1.366(12)
O(2)-C(3)	1.433(6)	C(16)-C(17)	1.372(10)
C(3)-C(4)	1.537(8)	C(18)-C(19)	1.435(9)
C(3)-C(5)	1.513(7)	C(19)-C(20)	1.508(8)
C(3)-C(6)	1.524(7)	C(19)-N(26)	1.458(7)
C(6)-C(7)	1.498(8)	C(20)-C(21)	1.367(8)
C(6)-C(8)	1.547(8)	C(20)-C(25)	1.374(8)
C(6)-O(9)	1.442(6)	C(21)-C(22)	1.370(9)
C(10)-C(11)	1.337(8)	C(22)-C(23)	1.344(12)
C(10)-C(18)	1.520(7)	C(23)-C(24)	1.343(13)
C(11)-C(12)	1.465(8)	C(24)-C(25)	1.395(11)
C(12)-C(13)	1.389(8)		

Bond angles (deg):

O(2)-B(1)-O(9)	105.3(4)	C(10)-C(11)-C(12)	128.6(4)
O(2)-B(1)-C(10)	117.8(4)	C(11)-C(12)-C(13)	120.3(5)
O(2)-B(1)-N(26)	109.3(4)	C(11)-C(12)-C(17)	123.1(5)
O(9)-B(1)-C(10)	119.5(4)	C(13)-C(12)-C(17)	116.6(5)
O(9)-B(1)-N(26)	106.9(4)	C(12)-C(13)-C(14)	121.8(5)
C(10)-B(1)-N(26)	97.0(4)	C(13)-C(14)-C(15)	120.1(6)
B(1)-O(2)-C(3)	108.2(3)	C(14)-C(15)-C(16)	119.2(6)
O(2)-C(3)-C(4)	109.3(4)	C(15)-C(16)-C(17)	121.2(6)
O(2)-C(3)-C(5)	109.7(4)	C(12)-C(17)-C(16)	121.1(5)
O(2)-C(3)-C(6)	102.8(4)	C(10)-C(18)-C(19)	107.2(4)
C(4)-C(3)-C(5)	107.9(4)	C(18)-C(19)-C(20)	117.5(5)
C(4)-C(3)-C(6)	110.8(4)	C(18)-C(19)-N(26)	107.1(5)
C(5)-C(3)-C(6)	116.1(4)	C(20)-C(19)-N(26)	118.7(5)
C(3)-C(6)-C(7)	116.6(5)	C(19)-C(20)-C(21)	123.2(5)
C(3)-C(6)-C(8)	111.0(4)	C(19)-C(20)-C(25)	118.8(5)
C(3)-C(6)-O(9)	102.2(4)	C(21)-C(20)-C(25)	117.8(5)
C(7)-C(6)-C(8)	108.1(4)	C(20)-C(21)-C(22)	121.3(5)
C(7)-C(6)-O(9)	110.6(4)	C(21)-C(22)-C(23)	120.7(6)

C(8)-C(6)-O(9)	108.0(4)	C(22)-C(23)-C(24)	119.6(6)
B(1)-O(9)-C(6)	108.4(3)	C(23)-C(24)-C(25)	120.7(6)
B(1)-C(10)-C(11)	128.0(4)	C(20)-C(25)-C(24)	119.9(5)
B(1)-C(10)-C(18)	109.2(4)	B(1)-N(26)-C(19)	107.8(4)
C(11)-C(10)-C(18)	122.7(5)		

Chapter 2


Development of the Transition-Metal-Catalyzed Enantioselective Conjugate

Allylation

I. Introduction

The conjugate addition of organometallic reagents to α,β -unsaturated carbonyl electrophiles is an important process for construction of carbon-carbon bonds in organic synthesis (Scheme 2.1).¹ Importantly, this process introduces a new stereocenter into the product, and therefore, development of asymmetric conjugate addition processes is an important area of research for the synthesis of non-racemic organic molecules. Not surprisingly then, many groups have been involved in this area of research, and many useful metal-catalyzed asymmetric conjugate addition processes have been developed.^{1,2}

Scheme 2.1: General Metal-Catalyzed Conjugate Addition



Surprisingly, despite intense research in this area, the use of allyl nucleophiles in asymmetric conjugate addition is still an unsolved problem. Interestingly, while most

¹ (a) Tomioka, K.; Nagaoka, Y. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: Berlin, 1999; Vol. 3, p 1105. (b) Kraus, N.; Hoffmann-Röder, A. *Synthesis* **2001**, 171.

² Reviews: (a) Christoffers, J.; Koripelly, G.; Rosiak, A.; Rössle, M. *Synthesis* **2007**, 1279. (b) Rossiter, B. E.; Swingle, N. M. *Chem. Rev.* **1992**, 92, 771. (c) Sibi, M. K.; Shankar, M. *Tetrahedron* **2000**, 56, 8033.


organocuprates³ reliably afford conjugate addition over direct 1,2-addition, allylcuprates are a notable exception to this trend. The 1,4 to 1,2 selectivity for addition reactions with diallylcuprate is highly substrate dependent and often affords the 1,2-allylation product in preference to the conjugate addition product.⁴ As depicted in Scheme 2.2, cyclohexenone (**2.1**) reacts readily with lithium diallylcuprate giving the conjugate addition product in good yield; however, isophorone (**2.4**) gave only the homoallylic alcohol product. In contrast, isophorone undergoes clean conjugate addition when using other Grignard reagents in the presence of a Cu-metal (Scheme 2.2, eq 3).⁵ Clearly, Cu-based allyl nucleophiles are less reliable in conjugate addition reactions than their vinyl, aryl, and alkyl counterparts. Other methods have been developed to overcome this shortcoming with allyl nucleophiles; however, a general catalytic asymmetric conjugate allylation is elusive.

³ Reviews: (a) Woodward, S. *Chem. Soc. Rev.* **2000**, 29, 393. (b) Nakamura, E.; Mori, S. *Angew. Chem. Int. Ed.* **2000**, 39, 3750.

⁴ (a) House, H. O.; Fischer, Jr. W. F. *J. Org. Chem.* **1969**, 34, 3615. (b) Majetich, G.; Casares, A.; Chapman, D.; Behnke, M. *J. Org. Chem.* **1986**, 51, 1745.

⁵ Kharasch, M. S.; Tawney, P. O. *J. Am. Chem. Soc.* **1941**, 63, 2308.

Scheme 2.2: Reaction of Di(allyl)cuprate with Enones



II. Background

A. Non-Enantioselective Conjugate Allylation Processes

The earliest example of a reliable conjugate allylation was reported by Hosomi and Sakurai in 1978.⁶ Using a stoichiometric amount of TiCl_4 as Lewis acid, allylic silanes undergo effective conjugate allylation with a variety of unsaturated ketones. One notable example is given in Scheme 2.3 (eq 1), where a sterically hindered angular allyl group can be installed in excellent yield. Catalytic variants of the Sakurai conjugate allylation have been developed using either InCl_3 or HNTf_2 as catalysts (Scheme 2.3, eq 2 and 3, respectively).^{7,8} For the catalytic reaction with InCl_3 , the addition of TMSCl is crucial. This phenomenon was suggested to result from trapping of an intermediate In-enolate by TMSCl to free the catalyst and drive equilibrium in this process. It should be


⁶ Hosomi, A.; Sakurai, H. *J. Am. Chem. Soc.* **1978**, 99, 1673.

⁷ (a) Lee, P. H.; Lee, K.; Sung, S. -Y.; Chang, S. *J. Org. Chem.* **2001**, 66, 8646. (b) Lee, P. H.; Seomoon, D.; Kim, S.; Nagaiah, K.; Damle, S. V.; Lee, K. *Synthesis* **2003**, 2189.

⁸ Kuhnert, N.; Peverley, J.; Robertson, J. *Tetrahedron Lett.* **1998**, 39, 3215.


noted that TiCl_4 is not a competent catalyst under these conditions and must be used in stoichiometric amounts. Amberlyst 15 is also a suitable catalyst for intramolecular conjugate allylation when tethered allylic silanes are used.⁹

Scheme 2.3: Conjugate Allylation with Allylsilanes



In addition to allylsilane, allyltrimethylstannane is also an effective nucleophile in conjugate allylation when using 1 equiv of $(\text{Et}_2\text{Al})_2\text{SO}_4$ as Lewis acid (Scheme 2.4).¹⁰ However, this method does require somewhat more forcing conditions (80 °C reaction temperature) for successful reaction.

Scheme 2.4: Conjugate Allylation with Allyltrimethylstannane



⁹ Schinzer, D.; Ringe, K. *Synlett* **1994**, 463.


¹⁰ Hosomi, A.; Iguchi, H.; Endo, M.; Sakurai, H. *Chem. Lett.* **1979**, 977.

Further study into the conjugate allylation with allylcuprates has led to improvements in the 1,4-selectivity when using these reagents. Studies by House and coworkers showed that the purity of the Cu salt used to prepare the cuprate affects the regioselectivity in the conjugate allylation.¹¹ When using freshly purified CuBr•DMS to form the allylcuprate, good yield of conjugate allylation product **2.3** was obtained (Scheme 2.5, eq 1); however, when older samples of CuBr•DMS were used, mixtures of 1,4- and 1,2-addition products were isolated. This was postulated to result from decomposition of the allylcuprate to allyllithium, facilitated by Cu(II) contaminants; allyllithium preferentially gives 1,2-addition to the enone. Furthermore, even with freshly purified Cu-salts, the regioselectivity is highly dependent on the enone used in the reaction. However, a prediction of regioselectivity can be made based on the reduction potential (E_{red}) of the enone. If the enone has E_{red} that is -2.0 V or less negative, then conjugate allylation should be preferred over 1,2-addition. However, if the E_{red} is more negative than -2.0 V, the mode of addition cannot be predicted. It should be noted that E_{red} was initially used by House for predictive purposes because the mechanism for conjugate addition was believed to occur by a SET process at that time; this view of the reaction mechanism is no longer accepted.^{3b,12} Hence, a correlation between E_{red} and reaction outcome is purely coincidental.

¹¹ House, H. O.; Wilkins, J. M. *J. Org. Chem.* **1978**, *43*, 2443.

¹² For recent spectroscopic evidence for involvement of Cu(III) intermediates in cuprate chemistry, see: Bertz, S. H.; Cope, S.; Murphy, M.; Ogle, C. A.; Taylor, B. J. *J. Am. Chem. Soc.* **2007**, *129*, 7208. (b) Gärtner, T.; Henze, W.; Gschwind, R. M. *J. Am. Chem. Soc.* **2007**, *129*, 11362.


Scheme 2.5: Effect of Cu-Salt Purity in Conjugate Allylation



These problems associated with allylcuprates can be overcome by the use of allylic copper reagents with TMSCl as an additive (Scheme 2.6).¹³ The use of (allyl)Cu·LiI or (allyl)Cu·MgBr₂, prepared using either allyllithium or allylmagnesium bromide, respectively, reliably affords the conjugate allylation product. TMSCl must be employed as an additive for successful reaction; in its absence, 1,2-addition is observed in low yield. TMSCl is known to accelerate Cu-mediated and Cu-catalyzed conjugate addition reactions.³ Furthermore, when TMSCl is used in conjunction with higher or lower order allylcuprates, only 1,2-addition is observed. Crotyl and methallyl Grignard reagents also react with good regioselectivity with respect to the crotyl reagent. For cyclohexenone derivatives, crotylation α to the organometallic is observed (eq 2), while with cyclopentenone, crotylation γ to the organometallic results (not shown).

¹³ (a) Lipshutz, B. H.; Ellsworth, E. L.; Dimcock, S. H.; Smith, R. A. *J. Am. Chem. Soc.* **1990**, *112*, 4404.
 (b) Lipshutz, B. H.; Hackmann, C. *J. Org. Chem.* **1994**, *59*, 7437.

Scheme 2.6: Use of Allylic Copper Reagents in Conjugate Allylation



Other allylic organometallic reagents that have been employed for conjugate allylation include tantalum,¹⁴ barium,¹⁵ and indium.¹⁶ Notably, in the cases of allyltantalum and allylbarium reagents, high selectivity for conjugate allylation over 1,2-addition is observed in the absence of additives. Furthermore, substituted allyltantalum reagents (i.e. crotyl or cinnamyl) give the branched products in good yield, whereas substituted allylic barium reagents afford the linear product.

B. Asymmetric Conjugate Allylation Processes

Surprisingly, despite efforts to effect reliable conjugate addition of allylic nucleophiles, the addition of these nucleophiles with absolute stereochemical control is


¹⁴ Shibata, I.; Kano, T.; Kanazawa, N.; Fukuoka, S.; Baba, A. *Angew. Chem. Int. Ed.* **2002**, *41*, 1389.

¹⁵ Yanagisawa, A.; Habaue, S.; Yasue, K.; Yamamoto, H. *J. Am. Chem. Soc.* **1994**, *116*, 6130.

¹⁶ Lee, P. H.; Ahn, H.; Lee, K.; Sung, S. -Y.; Kim, S. *Tetrahedron Lett.* **2001**, *42*, 37.

currently limited to the use of chiral auxiliaries. Wu and co-workers¹⁷ have shown that unsaturated chiral electrophiles **2.25** and **2.27**, bearing the Evans' chiral oxazolidinone auxiliary,¹⁸ participate in the Sakurai conjugate allylation with good diastereocontrol (Scheme 2.7). The cinnamyl derived substrate **2.27** is allylated with lower diastereocontrol; however, recrystallization affords the pure diastereomer **2.28** in moderate yield. Furthermore, the auxiliary can be removed using LiOH to afford the carboxylic acid derivative.

Scheme 2.7: Asymmetric Sakurai Conjugate Allylation



The Williams' group has also employed this auxiliary in asymmetric conjugate allylation using both allylic tributylstannane¹⁹ and allyl Grignard²⁰ nucleophiles (Scheme 2.8). Notably, the choice of allylic nucleophile dictates which diastereomer is formed in

¹⁷ Wu, M. -J.; Wu, C. -C.; Lee, P. -C. *Tetrahedron Lett.* **1992**, 33, 2547.


¹⁸ Reviews: (a) Evans, D. A.; Kim, A. S. In *Handbook of Reagents in Organic Synthesis: Reagents, Auxiliaries, and Catalysts for C-C Bonds*; Coates, R. M.; Denmark, S. E., Eds.; John Wiley & Sons: New York, 1999; pp 91-101. (b) Ager, D. J.; Prakash, I.; Schaad, D. R. *Aldrichimica Acta* **1997**, 30, 3.

¹⁹ Williams, D. R.; Mullins, R. J.; Miller, N. A. *Chem. Commun.* **2003**, 2220.

²⁰ Williams, D. R.; Kissel, W. S.; Li, J. J. *Tetrahedron Lett.* **1998**, 39, 8593.

the reaction (eq 1 vs eq 2) thereby allowing access to either diastereomer from the same starting material. Note that these reactions require stoichiometric amounts of Lewis acid, and in the case of Grignard reagents, 1 equiv of CuBr•DMS is required. This is not surprising on the basis of the findings of Lipshutz and co-workers¹³ who determined that only allylic copper reagents reliably afford conjugate allylation products (see page 106). Finally, crotylstannanes can also be employed when Zr is used as the Lewis acid (Scheme 2.8, eq 3). Only an *anti*-relationship between the β and γ stereocenters is observed in this case.


Scheme 2.8: Asymmetric Conjugate Allylation with Allylic Stannanes and Allylmagnesium Chloride



C. Conjugate Addition Reactions Catalyzed by Group 10 Transition Metals

Seminal work by Mole and co-workers²¹ established the utility of Ni as a catalyst for conjugate addition of AlMe₃ to enones (Scheme 2.9). In the presence of Ni(acac)₂, AlMe₃ adds to enone **2.34**, in the 1,4-sense, giving vinyl acetate **2.35** after addition of Ac₂O. Notably, in the absence of the Ni(II) salt, only 1,2-addition of AlMe₃ to **2.34** is observed. Since this initial report, Ni-catalysis has been applied to the conjugate addition of organozirconium,²² organozinc,²³ organoboron,²⁴ organostannane,²⁵ and organoindium²⁶ reagents. Despite the use of Ni(II) salts as catalysts, the active catalyst is believed to be Ni(0), which is generated *in situ* from the reduction of Ni(II) under the reaction conditions.²²

Scheme 2.9: Ni-Catalyzed Conjugate Addition of AlMe₃ to Enones



This class of conjugate addition reactions requires a Lewis acid for successful reaction. The role of the Lewis acid in this process has been examined by Kurosawa and

²¹ Jefferey, E. A.; Meisters, A.; Mole, T. *J. Organomet. Chem.* **1974**, 174, 365.

²² Schwartz, J.; Loots, M. J.; Kosugi, H. *J. Am. Chem. Soc.* **1980**, 102, 1333.

²³ (a) Greene, A. E.; Lansard, J. -P.; Luche, J. -L.; Petrier, C. *J. Org. Chem.* **1984**, 49, 931. (b) Soai, K.; Hayasaka, T.; Ugajin, S. *J. Chem. Soc., Chem. Commun.* **1989**, 516. (c) For a Pd(0)-catalyzed variant see: Marshall, J. A.; Herold, M.; Eidam, H. S.; Eidam, P. *Org. Lett.* **2006**, 8, 5505.


²⁴ (a) Hirano, K.; Yorimitsu, H.; Oshima, K. *Org. Lett.* **2007**, 9, 1541. (b) Shirakawa, E.; Yasuhara, Y.; Hayashi, T. *Chem. Lett.* **2006**, 35, 768.

²⁵ Grisso, B. A.; Johnson, J. R.; Mackenzie, P. B. *J. Am. Chem. Soc.* **1992**, 114, 5160.

²⁶ Pérez, I.; Sestelo, J. P.; Maestro, M. A.; Mouriño, A.; Sarandeses, L. A. *J. Org. Chem.* **1998**, 63, 10074.


co-workers.²⁷ When enal or enone Pd-complexes **2.37** or **2.39** are treated with various Lewis acids, complexes **2.38** or **2.40** are isolated in near quantitative yields, respectively (Scheme 2.10, eq 1 and 2). Furthermore, in the presence of 5 mol% $\text{Pd}_2(\text{dba})_3$ and 10 mol% PPh_3 , effective conjugate addition of AlMe_3 to enone **2.41** is observed (eq 3). Based on these observations, a plausible catalytic cycle for this transformation was developed by the authors, which correlates to the findings of Mackenzie²⁵ (Scheme 2.11). Activation of $(\text{PPh}_3)_2\text{Pd}(\text{enone})$ complex (**2.43**) by a Lewis acid (AlMe_3) enables oxidative addition of the metal to the bound enone affording **2.44**. Subsequent transmetalation and reductive elimination generates the aluminum enolate **2.46**, which is quenched upon workup to afford the observed product (**2.42**).

Scheme 2.10: Reaction of $\text{L}_2\text{Pd}(\text{enone})$ Complexes with Lewis Acids



²⁷ (a) Ogoshi, S.; Yoshida, T.; Nishida, T.; Morita, M.; Kurosawa, H. *J. Am. Chem. Soc.* **2001**, *123*, 1944. (b) Morita, M.; Inoue, K.; Ogoshi, S.; Kurosawa, H. *Organometallics* **2003**, *22*, 5468. (c) Ogoshi, S.; Morita, M.; Kurosawa, H. *J. Am. Chem. Soc.* **2003**, *125*, 9020.

Scheme 2.11: Mechanism of the Pd-Catalyzed Conjugate Addition of AlMe₃



This mechanism for conjugate addition catalyzed by group 10 metals, which relies on redox chemistry at the metal center, is in stark contrast to the extensively studied conjugate addition of organoboron, organosilicon, organotin, and organotitanium reagents catalyzed by Rh(I) complexes.²⁸ More recently Pd(II) complexes have also been shown to effect conjugate addition when using boronic acids²⁹ or organosiloxanes³⁰ as nucleophiles. In these systems, Rh(I)/Rh(III) or Pd(0)/Pd(II) catalytic cycles are not believed to be responsible for product formation. Rather, transmetalation of the nucleophile to the Rh(I) or Pd(II) salt occurs, followed by addition to the α,β -unsaturated

²⁸ Review: Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, *103*, 2829.

²⁹ (a) Nishikata, T.; Yamamoto, Y.; Miyaoura, N. *Angew. Chem. Int. Ed.* **2003**, *42*, 2768. (b) Lu, X.; Lin, S. *J. Org. Chem.* **2005**, *70*, 9651. (c) Nishikata, T.; Yamamoto, Y.; Miyaoura, N. *Organometallics* **2004**, *23*, 4317. (d) Gini, F.; Hessen, B.; Minaard, A. J. *Org. Lett.* **2005**, *7*, 5309. (e) Horiguchi, H.; Tsurugi, H.; Satoh, T.; Miura, M. *J. Org. Chem.* **2008**, *73*, 1590. (f) He, P.; Lu, Y.; Dong, C. –G.; Hu, Q. –S. *Org. Lett.* **2007**, *9*, 343. (g) Yamamoto, T.; Iizuka, M.; Ohta, T.; Ito, Y. *Chem. Lett.* **2006**, *35*, 198 and references therein.

³⁰ (a) Lerebours, R.; Wolf, C. *Org. Lett.* **2007**, *9*, 2737. (b) Gini, F.; Hessen, B.; Feringa, B. L.; Minaard, A. J. *Chem. Commun.* **2007**, 710. (c) Denmark, S. E.; Amishiro, N. *J. Org. Chem.* **2003**, *68*, 6997.

carbonyl electrophile giving a Rh(I) or Pd(II) enolate. Quench of the enolate by a proton source, usually water, releases the product and a Rh(I) or Pd(II) salt.


III. Discovery and Development of a Transition-Metal-Catalyzed Conjugate Allylation with Allylboronate Ester Nucleophiles³¹

A. Reaction Discovery

While developing the asymmetric allene diboration described in Chapter 1, a side-product was isolated from the reaction and was determined to be **2.49** (Scheme 2.12, eq 1). This product may arise from conjugate addition of allylboronate ester **2.48** to dibenzylidene acetone (dba), which was present in solution from the precatalyst. Furthermore, if dba was added to the diboration reaction after completion, **2.48** was not observed in the ¹H NMR spectrum of the unpurified reaction mixture, and **2.49** was isolated in good yield (Scheme 2.12, eq 2). Control experiments verified that both Pd₂(dba)₃ and PCy₃ were required for conversion to **2.49**. Mixing **2.47** with dba in toluene without both of these reagents resulted in no conversion. Additionally, the Pd-catalyzed reaction of enantiomerically pure **2.48** with dba, followed by oxidation of the vinylboronate functionality, afforded **2.50** as a single diastereomer in good yield (eq 3). However, the chirality transfer in this process was low.

³¹ (a) Sieber, J. D.; Liu, S.; Morken, J. P. *J. Am. Chem. Soc.* **2007**, *129*, 2214. (b) Sieber, J. D.; Morken, J. P. *J. Am. Chem. Soc.* **2008**, *130*, 4978.

Scheme 2.12: Conjugate Allylation with Allene Diboration Products



To study this reaction further, dba was treated with commercially available allylboronic acid pinacol ester (**2.51**) using $\text{Pd}_2(\text{dba})_3/\text{PCy}_3$ as catalyst (Scheme 2.13). This reaction resulted in the formation of conjugate allylation product **2.52** in 79% yield after only 1 h at room temperature. Again, both Pd and phosphine were required for conjugate allylation. Interestingly, only dba was reactive under these conditions. When applying this catalyst system to conjugate allylation of the α,β -unsaturated electrophiles depicted in Figure 2.1, no conjugate allylation was detected. In addition, heating these reactions to 65 °C in THF did not improve reactivity.

Scheme 2.13: Pd-Catalyzed Conjugate Allylation of Dbal





Figure 2.1: Electrophiles Tested in the Pd-Catalyzed Conjugate Allylation



B. Reaction Development

Discovery of a conjugate allylation process that proceeded in the presence of a transition metal catalyst and a phosphine ligand provided an opportunity to develop an asymmetric catalytic conjugate allylation, whereby a chiral ligand could be used to control facial selectivity in the reaction. However, because only dialkylidene ketones, such as dba, were reactive electrophiles, unsymmetrical dialkylidene ketone **2.62** was prepared and subjected to the conjugate allylation procedure (Table 2.1). For conjugate allylation of **2.62** to be synthetically useful, the catalyst for the reaction must effect chemoselective addition of the allyl nucleophile to either the β - or β' -position of the dialkylidene ketone selectively. To achieve this goal, a survey of group 10 metals, in

conjunction with phosphine ligands, were analyzed in the conjugate allylation of **2.62**.

The results of this investigation are given in Table 2.1.

Table 2.1: Catalyst Optimization for the Conjugate Allylation of 2.62

Reaction scheme: **2.62** (a 1,5-dicarbonyl compound with a phenyl group at the β' position and an *n*-pentyl group at the β position) reacts with **2.51** (an allylboronate ester) in the presence of a catalyst in THF at room temperature to yield **2.63** (the β -allylation product) and **2.64** (the β' -allylation product).

entry	metal	ligand	time (h)	β : β' ^a	% yield ^b
1	2.5% Pd ₂ (dba) ₃	6% PCy ₃	2	n/a	< 5
2	2.5% Pd ₂ (dba) ₃	6% PCy ₃	16	73:27	67
3	2.5% Pd ₂ (dba) ₃	6% P(NMe ₂) ₃	2	50:50	76
4	2.5% Pd ₂ (dba) ₃	6% PPh ₃	2	50:50	18
5	2.5% Pd ₂ (dba) ₃	6% P(OPh) ₃	2	36:64	17
6	10% Ni(cod) ₂	20% PCy ₃	4	88:12	82
7	10% Ni(cod) ₂	20% P(NMe ₂) ₃	4	71:29	81
8	10% Ni(cod) ₂	20% PPh ₃	4	70:30	49
9	10% Ni(cod) ₂	20% P(OPh) ₃	4	52:48	55

^aChemoselectivity determined by ¹H NMR analysis of the unpurified reaction mixture. ^bYield of conjugate allylation product after silica gel chromatography. Isolated as a mixture of constitutional isomers. Value is an average of at least 2 experiments.

Under Pd-catalysis, when a bulky, Lewis basic ligand was employed, the allylation was sluggish, but allylation preferentially occurred at the alkylidene site (β -position) of the substrate, giving **2.63** in a modest 73:27 ratio (entry 2). HMPT proved to be an efficient ligand in terms of reaction rate; however, no chemoselection was observed in this case (entry 3). Notably, when using Ni as the catalyst, good yield of the

conjugate allylation product could be obtained in only 4 h at room temperature (entry 6). Furthermore, the highest chemoselectivity was obtained in this case, favoring conjugate allylation to the alkylidene site of **2.62** in an 88:12 ratio. Under Ni-catalysis, a drop in both chemoselectivity and rate was observed when using phosphines of decreased Lewis basicity (entries 6-9).

C. Scope and Utility of the Ni-Catalyzed Conjugate Allylation

With identification of a Ni-catalyst that afforded chemoselective conjugate allylation to the alkylidene unit of unsymmetrical dialkylidene ketone **2.62**, analysis of the scope of this catalyst commenced. A class of unsymmetrical dialkylidene ketones was synthesized where each ketone contained a benzylidene group, for substrate activation, and a variable alkylidene group (**2.66**, Table 2.2). These ketones were most conveniently prepared using the Weinreb³² amide ketone synthesis. Those used in the conjugate allylation reaction are shown in Table 2.2. In general, the ketones were prepared in good yields, where yields were optimal when vinyl iodides were used as precursors to the vinyl lithium nucleophile. Furthermore, *n*-BuLi could be used for the Li/halogen exchange when using vinyl iodides (entries 1 and 3-6), while *t*-BuLi was required for the Li/halogen exchange when employing vinyl bromides (entries 2, 7, and 8); use of *n*-BuLi in the Li/halogen exchange step with vinyl bromides led to inferior yields. In addition, Grignard reagents were ineffective in the ketone synthesis.

³² Nahm, S.; Weinreb, S., M. *Tetrahedron Lett.* **1981**, 22, 3815.

Table 2.2: Synthesis of Unsymmetrical Dialkylidene Ketones

entry	X	alkylLi	product	% yield
1	I	<i>n</i> -BuLi		91
2	Br	<i>t</i> -BuLi		76
3	I	<i>n</i> -BuLi		96
4	I	<i>n</i> -BuLi		86
5	I	<i>n</i> -BuLi		75
6	I	<i>n</i> -BuLi		87
7	Br	<i>t</i> -BuLi		53
8	Br	<i>t</i> -BuLi		53

With access to a variety of unsymmetrical dialkylidene ketones, the substrates prepared in Table 2.2 were subjected to the conjugate allylation reaction using Ni(cod)₂/PCy₃ as catalyst. The results of this survey are shown in Table 2.3. When methyl-substituted ketone **2.67** or *t*-butyl-substituted ketone **2.69** was used in the conjugate allylation, high chemoselectivity was observed in the allylation (entries 2 and

4). Furthermore, when trisubstituted alkylidene substrates **2.72** and **2.73** were used, complete chemoselectivity was observed; however, the reaction temperature had to be increased to 65 °C to achieve useful conversions (entries 8 and 9). The result in entry 8 is notable since the reaction installs the allyl group at the more sterically encumbered carbon, allowing access to an all-carbon quaternary center. Alkoxy groups were also tolerated in the reaction (entries 5 and 6). Notably, in the presence of an allylic TBS-ether, conjugate allylation proceeded in good yield (entry 6) even though allylic ethers are known to participate in allylic substitution reactions with boronic acids under Ni-catalysis.³³ However, chemoselectivity was low for this substrate.

³³ Chung, K. G.; Miyake, Y.; Uemura, S. *J. Chem. Soc., Perkin Trans. 1* **2000**, 15.

Table 2.3: Substrate Scope in the Ni-Catalyzed Conjugate Allylation

Reaction scheme: Enone **2.74** (with substituents R, R', R'') reacts with allylboronate **2.51** (1.2 equiv) in the presence of 10% Ni(cod)₂ and 20% PCy₃ in THF at room temperature to yield β -addition product **2.75** and β' -addition product **2.76**.

entry	R	R'	R''	time (h)	β : β' ^a	% yield ^b
1	<i>n</i> -pentyl	H	H	4	88:12	82
2	Me	H	H	24	95:5	75
3	Cy	H	H	4	83:17	83
4	<i>t</i> -Bu	H	H	4	>95:5	74
5	CH ₂ OTBS	H	H	24	61:39	66
6	CH ₂ CH ₂ OTBS	H	H	7	85:15	82
7	Ph	H	H	4	n/a	74
8 ^{c,d}	Me	Me	H	24	>95:5	80
9 ^c	Me	H	Me	24	>95:5	76 ^e


^aDetermined by ¹H NMR analysis of the unpurified reaction mixture. ^bIsolated yield after silica gel chromatography. Yield represents an isomeric mixture and is an average of at least two experiments in each case. ^cReaction was performed at 65 °C. ^d2.0 equiv of **2.51** was used. ^eProduct consists of a 1.4:1 diastereomeric mixture.

To further develop the utility of this conjugate allylation protocol, a method for removal of the activating benzylidene unit was sought. A Sn-catalyzed Baeyer-Villiger reaction, developed by Shibasaki³⁴ and co-workers, furnished unsaturated ester **2.79** with complete chemo- and regioselectivity (Scheme 2.14, eq 1). Furthermore, the ester could be saponified with LiOH in excellent yield (Scheme 2.14, eq 2). In addition to cleavage of the benzylidene activating group, ring-closing metathesis was used to access 5-

³⁴ Göttlich, R.; Yamakoshi, K.; Sasai, H.; Shibasaki, M. *Synlett* **1997**, 971.

substituted cyclohexenone **2.81** (Scheme 2.14, eq 3). The Hoveyda-Grubbs³⁵ second generation catalyst was the optimal metathesis catalyst in this case.

Scheme 2.14: Transformations of the Benzyldiene Activating Group




D. Mechanistic Aspects of the Ni-Catalyzed Conjugate Allylation

A reasonable catalytic cycle for the conjugate allylation is shown in Scheme 2.15. Based off of the work of Kurosawa²⁷ and Mackenzie²⁵ on the Pd- and Ni-catalyzed conjugate addition reaction of enones (see section II.C, page 110), allylboronate ester **2.51** likely acts as a Lewis acid to facilitate oxidative addition of Ni(0) to dba, furnishing LNi(allyl) complex **2.83**. Subsequent transmetalation would afford bis(allyl)Ni-complex **2.84**. Reductive elimination from **2.84** gives the product masked as a boron enolate (**2.85**), which is quenched upon workup. In support of this hypothesis, when Pd(TFA)₂ or

³⁵ Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, *122*, 8168.

Ni(acac)₂ was used in the conjugate allylation, no products were observed, indicating that M(0) is likely the active catalyst in the reaction. When Ni(0) was generated *in situ* by treatment of Ni(acac)₂ with DIBAL,³⁶ before addition of **2.51** and **2.62**, conjugate allylation was observed, further implicating M(0) as the active catalyst.

Scheme 2.15: Proposed Catalytic Cycle for the Conjugate Allylation




Attempts to trap the boron enolate intermediate, after the conjugate allylation was complete, were successful and informative (Scheme 2.16). In this experiment, conjugate allylation was performed in the presence of 4Å MS and subsequently quenched by the addition of *N*-(trimethylsilyl)imidazole (**2.86**).³⁷ Silyl enol ether **2.87** was isolated from the reaction mixture as a single olefin isomer. NOESY analysis of **2.87** verified the *E*-

³⁶ DIBAL is known to reduce Ni(II) to Ni(0). See: Krysan, D. J.; Mackenzie, P. B. *J. Org. Chem.* **1990**, *55*, 4229 and references therein.

³⁷ For the use of *N*-(trimethylsilyl)imidazole for silylation of boron enolates, see: Hooz, J.; Oudenes, J. *Tetrahedron Lett.* **1983**, 24, 5695.

configuration of the silyl enol ether. When TMSCl was used in the trapping step, **2.87** was not observed, and only the ketone product was isolated.

Scheme 2.16: Trap of the Intermediate Boron Enolate




The enhanced reactivity of dialkylidene ketones relative to simple enones must be the result of a reduced energy barrier for one of the elementary steps in the catalytic cycle by the additional alkene unit of the DAK. While the oxidative addition or transmetalation steps may be facilitated by the additional alkene present in the substrate, we propose that acceleration of the reductive elimination step is responsible for the enhanced reactivity of dialkylidene ketones. This proposal is on the basis of the calculated reductive elimination barriers for bis(allyl)Pd-complexes (Figure 2.2).³⁸ Of the three possibilities, it is calculated that in the presence of donor ligands, reductive elimination from bis(η¹-allyl)Pd-complex **2.90** is the lowest energy pathway. While complex **2.90** is slightly higher in energy relative to **2.88** or **2.89**, **2.90** has the opportunity to undergo a [3,3']-reductive elimination pathway as depicted in Figure 2.2. Reductive elimination from **2.90** is calculated to occur via this process rather than through direct

³⁸ Mendez, M.; Cuerva, J. M.; Gomez-Bengoa, E.; Cardenas, D. J.; Echavarren, A. M. *Chem. –Eur. J.* **2002**, 8, 3620.

coupling of carbons 3 and 3'. Thus reductive elimination is significantly more facile from **2.90** relative to **2.88** or **2.89**.

Figure 2.2: Calculated Reductive Elimination Barriers for Bis(allyl)Pd-Complexes³⁸



Based on this data, an analogous mode of reductive elimination can be envisioned in the conjugate allylation from intermediate **2.91** (Scheme 2.17, eq 1). Within the proposed catalytic cycle, **2.91** is formed after oxidative addition and transmetalation. As depicted in Scheme 2.17, **2.91** may undergo reductive elimination through a [3,3']-reductive elimination pathway without formation of a higher energy bis(η^1 -allyl)Ni(PR₃)₂ complex. This process is not available to enones lacking the extra alkene present in dialkylidene ketones, which may explain their resistance to reaction in the conjugate allylation. To ascertain if this scenario was feasible, Dr. Shubin Liu³⁹ performed DFT calculations on model system **2.92** using the B3LYP⁴⁰ method (Scheme 2.17, eqs 2 and 3).³¹ The Stuttgart RSC 1997 ECP basis set⁴¹ was used for Pd, and 6-311+G*⁴² was used for other elements. These calculations verified that reductive elimination through a

³⁹ Division of Research Computing, Information Technology Services, University of North Carolina, Chapel Hill, NC 27599

⁴⁰ (a) Becke, A. D. *J. Chem. Phys.* **1993**, 98, 1372. (b) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, 37, 785.


⁴¹ Bergner, A.; Dolg, M.; Kuechle, W.; Stoll, H.; Preuss, H. *Mol. Phys.* **1993**, 80, 1431.

⁴² Krishnan, R.; Brinkley, J. S.; Seeger, R.; Pople, J. A. *J. Chem. Phys.* **1980**, 72, 650.


[3,3']-pathway (eq 2) has a remarkably low barrier (1.52 kcal/mol). Replacement of the pendant alkene with a methyl group increased the barrier significantly (Scheme 2.17, eq 3). In addition, the calculations predicted that 1,2-addition (**2.94**) and not 1,4-addition (**2.95**) should predominate in the unactivated enone. Figure 2.3 shows the calculated transition state for the reductive elimination of **2.92**. This proposal predicts that the *E*-configured boron enolate should be produced in the reaction, which correlates nicely to the observed *E*-configuration of silyl enol ether **2.87**.

Scheme 2.17: Calculated Reductive Elimination Barriers in the Conjugate

Allylation



**Figure 2.3: Calculated Transition Structure for the Reductive Elimination from
2.92**




With calculated evidence for the facilitation of the reductive elimination step via a [3,3']-reductive elimination pathway, experimental evidence to support these calculations was sought. Therefore, a dialkylidene ketone bearing an internal phosphino group (**2.96**) was prepared from **2.71** (Scheme 2.18, eq 1). It was postulated that if this internal phosphino group served as a ligand for Ni in the conjugate allylation, it should direct oxidative addition of the metal catalyst to the proximal alkene of **2.96**. If this was the case, and the [3,3']-reductive elimination proceeded as proposed above, then conjugate allylation should occur at the distal alkene of **2.96**, giving **2.97**, and ultimately furnish **2.98** (Scheme 2.18, eq 2). Subjection of ketone **2.96** to the conjugate allylation reaction, in the absence of any external phosphine ligand, gave conjugate allylation exclusively at the benzylidene site in **2.96** (Scheme 2.19, eq 1).⁴³ No evidence for conjugate allylation


⁴³ Note that in addition to the conjugate allylation product **2.98** some of the derived phosphine oxide was isolated due to oxidation of **2.98** during purification.

to the alkene proximal to the phosphine directing group was observed. Furthermore, a control experiment using dialkylidene ketone **2.100** as the substrate proceeded with little chemoselection, verifying that the phosphino group in **2.96** was responsible for the high chemoselectivity (Scheme 2.19, eq 2).

Scheme 2.18: Analysis of a Phosphino-Directing Dialkylidene Ketone




Scheme 2.19: Conjugate Allylation of a Phosphine Containing DAK



The experiments outlined in Scheme 2.19 offer further support for the proposed [3,3']-reductive elimination pathway. However, the possibility that the tether in **2.96** may be long enough to direct the Ni-catalyst to the distal alkene in the substrate could not be ruled out. If this is the case, **2.98** may be formed by oxidative addition to the distal olefin in **2.96** followed by transmetalation and direct reductive elimination. To ascertain which olefin of **2.96** may coordinate to Ni, complex **2.101** was prepared (Scheme 2.20). When mixing equimolar amounts of **2.96**, Ni(cod)₂, and PPh₃ in THF at room temperature, complex **2.101** was formed. While suitable crystals for X-ray diffraction could not be obtained, NMR analysis of the red solid, when dissolved in solution, allowed for verification of the structure of **2.101**. Proton and carbon assignments were made by ¹H, ¹³C, COSY, HSQC, and NOESY spectroscopic analyses. The hydrogen atoms at C3 and C4 in **2.101** exhibited an upfield shift of Δδ 3.01 ppm and Δδ 2.45 ppm, respectively relative to **2.96**. Likewise the carbon resonances for C3 and C4 moved to higher field by

$\Delta\delta$ 71.6 ppm and $\Delta\delta$ 60.2 ppm relative to starting material. In contrast, the hydrogen atoms at C6 and C7 of **2.101** exhibited a less substantial perturbation ($\Delta\delta$ 1.01 ppm and $\Delta\delta$ <0.05 ppm). Similarly, the carbon resonances for C6 and C7 were relatively unaffected, showing $\Delta\delta$ 0.6 ppm and $\Delta\delta$ 1.8 ppm, respectively. In addition to affecting the alkene hydrogens, the metal association also affected the chemical shift of the hydrogen atoms at C2 and C1. The proton resonance at C2 of **2.101** experienced a substantial upfield shift ($\Delta\delta$ 0.76 ppm), and the protons at C1 in the metal complex were now diastereotopic, one moving 0.72 ppm downfield and the other 0.44 ppm upfield. Collectively, these spectroscopic changes observed when **2.96** was treated with Ni(cod)₂ and PPh₃, were consistent with Ni binding to the alkene adjacent to the directing group. Other spectroscopic features were also consistent with this formulation; the ³¹P NMR showed two doublet resonances at 41.21 ppm and 28.59 ppm (J = 32 Hz).

Scheme 2.20: Synthesis of (PPh₃)Ni(**2.96**) Complex



The identity of complex **2.101** and the chemoselectivity in the conjugate allylation of **2.96** provides further support for the validity of a [3,3']-reductive elimination pathway


operating in the conjugate allylation of dialkylidene ketones. This then may explain the enhanced reactivity of DAKs relative to other enones.

To further investigate the intermediate bis(allyl)Ni-complexes formed in the conjugate allylation, the reaction was carried out using deuterium labeled allylboronic acid pinacol ester (**2.51-d₂**, Scheme 2.21). Interestingly, under Ni-catalysis (eq 1), scrambling of the deuterium label was observed in the product; however, under Pd-catalysis (eq 2), the site of the deuterium label was retained in the product. Since there is good evidence that transmetalation of allyl groups occurs via an S_E'-type mechanism,⁴⁴ transmetalation from complex **2.102** should afford **2.103** with the label at the vinylic position (Scheme 2.22). When M=Pd, [3,3']-reductive elimination from **2.103** would furnish the retained label in the product. However, when M = Ni, η^1 - η^3 - η^1 allyl isomerization, presumably through **2.104**, likely occurs faster than reductive elimination and leads to scrambling of the label. The lower yields obtained in the reactions using **2.51-d₂** were attributed to the presence of vinylboronic acid pinacol ester, which was a contaminant in **2.51-d₂** resulting from its preparation.⁴⁵


⁴⁴ (a) Hayashi, T.; Konishi, M.; Kumada, M. *J. Chem. Soc. Chem. Commun.* **1983**, 736. (b) Hatanaka, Y.; Goda, K.; Hiyama, T. *Tetrahedron Lett.* **1994**, 35, 1279. (c) Hiyama, T.; Matsushashi, H.; Fujita, A.; Tanaka, M.; Hirabayashi, K.; Shimizu, M.; Mori, A. *Organometallics* **1996**, 15, 5762. (d) Yamamoto, Y.; Takada, S.; Miyaura, N. *Chem. Lett.* **2006**, 35, 704.

⁴⁵ See page 333.

Scheme 2.21: Deuterium Labeling Experiments




Scheme 2.22: Possible Explanation for Labeling Experiments



Finally, NMR investigations of mixtures of dba, Ni(cod)₂, and PCy₃ in THF- d_8 revealed that a new complex formed that had incorporated both dba and PCy₃. Analysis of the ³¹P NMR spectrum revealed that only one phosphine was ligated to the metal

center even in the presence of excess PCy₃: when 2.0 equiv of PCy₃ was used, relative to dba and Ni(cod)₂, a new peak in the ³¹P NMR spectrum was observed at 38 ppm; in addition to this new peak, free PCy₃ was also present, with the integration between these two peaks being ~1:1. Importantly, only singlets were observed in the ³¹P NMR spectrum. No evidence for a (PCy₃)₂Ni-adduct was observed. Furthermore, integration of the ¹H NMR spectrum when using a 1:1:1 mixture of Ni:PCy₃:dba was consistent with only one olefin of dba binding to the Ni-center in this complex. These observations may be explained by invoking the existence of dimeric Ni-complex **2.106** (Figure 2.4). While concrete evidence for such a structure by X-ray crystallography could not be obtained, similar (phosphine)Ni(enone) dimers have been isolated by Kurosawa.⁴⁶ Further evidence for dimeric Ni-catalysts was found while studying the enantioselective version of this reaction. These results are discussed in section IV.C (page 143).

Figure 2.4: Proposed [(PCy₃)Ni(dba)]₂ Dimer



⁴⁶ Ogoshi, S.; Nagata, M.; Kurosawa, H. *J. Am. Chem. Soc.* **2006**, 128, 5350.

IV. Development of a Transition-Metal-Catalyzed Asymmetric Conjugate Allylation with Allylboronate Ester Nucleophiles^{31b}

A. Discovery of a Competent Chiral Catalyst

With a better understanding of this new conjugate allylation obtained from our studies of the non-enantioselective reaction, our focus changed to finding a general chiral catalyst that would affect both a chemoselective and an enantioselective conjugate allylation. Chiral TADDOL⁴⁷-derived phosphoramidites, phosphites, and phosphonites (see Figure 2.5) were initially examined as ligands in the reaction when Ni(cod)₂ was used as the metal precatalyst. The results of this study are shown in Table 2.4.

Using (TADDOL)MONOPHOS⁴⁸ (**2.107**) as the chiral ligand, or other phosphoramidites bearing the TADDOL scaffold, the conjugate allylation product was obtained in moderate yield and poor enantioselectivity (entries 1-3). Interestingly, all of the TADDOL-derived ligands examined gave preferential allylation to the benzyldiene site in **2.62** rather than the alkylidene site as was observed in the non-enantioselective reaction. Modification of the TADDOL backbone by substitution at the 3,5-positions of the phenyl ring did lead to improvements in selectivity. Xylyl-substituted TADDOL phosphoramidite **2.110** gave excellent chemoselectivity in the reaction; however, the enantioselectivity was not acceptable (entry 4). Installing larger 3,5-di-*t*-butylphenyl groups on the TADDOL backbone led to an improvement in enantioselectivity (entry 5, 69% ee). Since chemoselectivity and yield were low with this ligand, other 3,5-di-*t*-

⁴⁷ Review: Seebach, D.; Beck, A. K.; Heckel, A. *Angew. Chem. Int. Ed.* **2001**, 42, 92.

⁴⁸ (a) Keller, E.; Maurer, J.; Naasz, R.; Schader, T.; Meetsma, A.; Feringa, B. L. *Tetrahedron Asymm.* **1998**, 9, 2409. (b) Alexakis, A.; Vastra, J.; Burton, J.; Benhaim, C.; Manganey, P. *Tetrahedron Lett.* **1998**, 39, 7869.

butylphenyl TADDOL ligands were examined (entries 6-8). Gratifyingly, phosphonite **2.114** affected the conjugate allylation in excellent chemoselectivity, good yield, and high enantioselectivity (entry 8, 88% ee). Other TADDOL-derived phosphonites were inferior to **2.114** in the asymmetric allylation (entries 9 and 10).


Table 2.4: Survey of Chiral Ligands in the Conjugate Allylation

Reaction scheme: Enone **2.62** (Ph-CH=CH-C(=O)-CH=CH-n-pentyl) reacts with allylboronate **2.51** (1.2 equiv) in the presence of 10% Ni(cod)₂ and 20% ligand in THF at room temperature for 8 hours to yield (S)-**2.64** (β'-addition product).

entry	Ligand ^a	β:β' ^b	% yield ^c	% ee ^{d,e}
1	2.107	1:3.8	36	44 (75)
2	2.108	1:4.3	31	14 (26)
3	2.109	1:2.2	57	-2 (84)
4	2.110	1:≥20	53	44
5	2.111	1:4.7	29	69 (0)
6	2.112	1:2.5	24	47 (1)
7	2.113	1:16	62	-3
8	2.114	1:20	69	88
9	2.115	1:3.6	49	28 (81)
10	2.116	1:≥20	50	-15

^aSee Figure 2.5 for ligand structures. ^bValue determined by ¹H NMR analysis of the unpurified reaction mixture. ^cIsolated yield after silica gel chromatography. Isolated as a mixture of constitutional isomers. ^dValue represents that of the major isomer. ^eValue in parentheses represents that of the minor constitutional isomer.

Figure 2.5: Chiral Ligands Used in the Chiral Catalyst Survey



Upon identification of a chiral catalyst for asymmetric conjugate allylation, reaction parameters were optimized in order to reduce catalyst loading and maximize yield and selectivity. The catalyst loading could be reduced to 5% Ni(cod)₂ and 10% **2.114** when conducting the reaction at 0.5 M concentration relative to substrate. Furthermore, the enantioselectivity could be increased to 93% ee by using toluene as the reaction solvent instead of THF.

Based on the mechanistic paradigm discussed in section III.D (page 121), it is likely that the nature of the substituent on the activating alkene may affect reaction selectivity. That is, since oxidative addition of the catalyst is proposed to occur at the activating alkene, interaction between the coordination sphere of the metal and the substituents on this alkene may be important for controlling the facial selectivity of the reaction. Therefore, the alkyl substituent on the activating alkylidene unit of the substrate was modified to determine if this group affected reaction selectivity. Furthermore, smaller groups would be more efficient in terms of atom economy. The results of this investigation are shown in Table 2.5. A dramatic effect on chemoselectivity and enantioselectivity was observed. When the substituent of the alkylidene unit was small, higher chemoselectivities were obtained, but suboptimal enantioselectivities prohibited the use of these groups (entries 2 and 3). The larger cyclohexyl group increased enantioselectivity, but chemoselectivity decreased (entry 4). Thus, the *n*-pentyl group afforded the best combination of high chemoselectivity and enantioselectivity.

Table 2.5: Optimization of the Activating Alkene in the DAK Substrate

entry	R	$\beta:\beta'$ ^a	% yield ^b	% ee ^c
1	<i>n</i> -pentyl	1:17	77	93
2	Me	1: \geq 20	60	67
3	Et	1: \geq 20	82	83
4	Cy	1:7	76	96










^aValue determined by ¹H NMR (entries 2-4) or GLC (entry 1) analysis of the unpurified reaction mixture.

^bIsolated yield after silica gel chromatography. Isolated as a mixture of constitutional isomers. ^cValue represents that of the major isomer.

B. Scope and Utility in the Ni-Catalyzed Asymmetric Conjugate Allylation

With the conditions for the asymmetric conjugate allylation optimized, the scope of the reaction was analyzed. Unsymmetrical dialkylidene ketones, all having a heptylidene unit as the activating group, were prepared in good yield using the Weinreb³² amide ketone synthesis (**2.121**). Results for the syntheses of these substrates are given in Table 2.6.

Table 2.6: Synthesis of Unsymmetrical Dialkylidene Ketones

$ \begin{array}{c} \text{O} \\ \parallel \\ \text{Ar}-\text{CH}=\text{CH}-\text{N}(\text{Me})\text{OMe} \end{array} + \text{I}-\text{CH}=\text{CH}-n\text{-pentyl} \xrightarrow[\text{THF, } -78^\circ\text{C, 15 min}]{n\text{-BuLi}} \text{Ar}-\text{CH}=\text{CH}-\text{C}(=\text{O})-\text{CH}=\text{CH}-n\text{-pentyl} $		
2.119	2.120	2.121
entry	product	% yield
1	 2.122	73
2	 2.123	90
3	 2.124	89
4	 2.125	79
5	 2.126	66
6	 2.127	82
7	 2.128	87
8	 2.129	88
9	 2.130	86

The scope of the asymmetric conjugate allylation is given in Table 2.7. Notably, in all cases, high enantioselectivity was obtained regardless of the aryl substituent of the substrate. Electron-deficient arenes were efficient in terms of reaction rate; however, chemoselectivity was lower for these substrates (entries 2 and 6). Electron-rich arenes reacted slower, and therefore, required longer reaction times; however, chemoselectivities were better for these substrates (entries 3, 8, and 9). Substitution at the *ortho* position of the substrate dramatically increased the chemoselectivity, but allylation was slower with these substrates (entries 4, 5 and 7). In addition, substrates bearing functionalized aromatic groups common in natural products and drug targets were effective in the reaction (entries 8 and 9).


Table 2.7: Substrate Scope in the Asymmetric Conjugate Allylation

entry	product	time (h)	$\beta':\beta^a$	% yield ^b	% ee
1		7	17:1	77	93
2		7	6.5:1	81	93
3		12	20:1	66	91
4		12	>20:1	66	93
5		48	32:1	45	94
6		7	7.0:1	78	94
7		12	49:1	81	92
8		12	21:1	60	92
9		12	>20:1	74	92
10		12	5.1:1	52	93

^aValue determined by ¹H NMR (entries 4 and 9) or GLC (entries 1-3, 5-8, and 10) analysis of the unpurified reaction mixture. ^bIsolated yield after silica gel chromatography. Value is an average of at least 2 experiments in each case.

The activating heptylidene unit required for the asymmetric conjugate allylation could be cleaved using the Sn-catalyzed Baeyer-Villiger oxidation³⁴ described previously (Scheme 2.23, eq 1). Saponification of the resultant ester afforded enantiomerically enriched carboxylic acid **2.80** in near quantitative yield.

Scheme 2.23: Cleavage of the Alkylidene Activating Unit



To further demonstrate the utility of the asymmetric conjugate allylation reaction, this methodology was applied to the synthesis of aminoguanidine **2.148** (Scheme 2.24). Aminoguanidine **2.148** was recently identified as a potent inhibitor of integral plasma membrane proteins that control intracellular pH through Na⁺/H⁺ exchange (NHE) across the cell membrane.⁴⁹ Such targets are potentially useful drugs for the treatment of ischemia-reperfusion injuries. Starting from conjugate allylation product **2.137**, ring-closing metathesis (RCM) afforded cyclic enone **2.142** in excellent yield. The synthesis of enantiomerically enriched 5-substituted cyclohexenones, analogous to **2.142**, via this process (i.e. conjugate allylation followed by RCM) is notable since these compounds are

⁴⁹ Fukumoto, S.; Imamiya, E.; Kusumoto, K.; Fujiwara, S.; Watanabe, T.; Shiraishi, M. *J. Med. Chem.* **2002**, *45*, 3009.

useful synthetic building blocks that can be difficult to prepare by other routes.⁵⁰ Subsequent α -iodination⁵¹ furnished α -iodoketone **2.143**. An α,β -unsaturated hydrazone side-chain was then introduced via Stille-coupling⁵² between **2.143** and stannane **2.144** to produce **2.145**. Finally, a one-pot sequence⁵³ involving thermal olefin isomerization, followed by electrocyclization, and elimination of dimethylamine afforded pyridine **2.146** in modest isolated yield. In addition to **2.146**, a side-product was isolated and tentatively assigned as quinoline **2.147** from its ¹H NMR spectrum.⁵⁴ The ratio of **2.146**:**2.147** was ~2:1 as determined from the ¹H NMR spectrum of the unpurified reaction mixture. Further optimization of the reaction conditions (solvent, concentration, temperature) did not suppress the formation of this side-product. At this point, the synthesis of **2.146** constituted a formal synthesis of **2.148**. To complete the synthesis, the aminoguanidine functional group was introduced according to the literature.⁴⁹

⁵⁰ Asymmetric synthesis of 5-substituted cyclohexenones is typically achieved by kinetic resolution. See: Chen, Q.; Kuriyama, M.; Soeta, T.; Hao, X.; Yamada, K. -I.; Tomioka, K. *Org. Lett.* **2005**, 7, 4439.


⁵¹ Krafft, M. E.; Cran, J. W. *Synlett* **2005**, 8, 1263.

⁵² For Pd-catalyzed couplings to α -iodoenones, see: (a) Sirisoma, N. S.; Johnson, C. R. *Tetrahedron Lett.* **1998**, 39, 2059. (b) Johnson, C. R.; Adams, J. P.; Collins, M. A. *J. Chem. Soc. Perkin Trans. 1* **1993**, 1. (c) Johnson, C. R.; Adams, J. P.; Braun, M. P.; Senanayaka, C. B. W. *Tetrahedron Lett.* **1992**, 33, 919. (d) Ruel, F. S.; Braun, M. P.; Johnson, C. R. *Org. Synth.* **1998**, 75, 69. (e) Banks, J. C.; Van Mele, D.; Frost, C. G. *Tetrahedron Lett.* **2006**, 47, 2863. (f) Negishi, E.-I. *J. Organomet. Chem.* **1999**, 576, 179.

⁵³ (a) Staben, S. T.; Kennedy-Smith, J. J.; Huang, D.; Corkey, B. K.; LaLonde, R. L.; Toste, F. D. *Angew. Chem. Int. Ed.* **2006**, 45, 5991. (b) Hosokawa, T.; Shimo, N.; Maeda, K.; Sonoda, A.; Murahashi, S.-I.; *Tetrahedron Lett.* **1976**, 383. (c) Verboom, W.; Van Eijk, P. J. S. S.; Conti, P. G. M.; Reinhoudt, D. N. *Tetrahedron* **1989**, 45, 3131. (d) Tanaka, K.; Mori, H.; Yamamoto, M.; Katsumua, S. *J. Org. Chem.* **2001**, 66, 3099.

⁵⁴ Rigorous characterization of **2.147** was not performed.

Scheme 2.24: Synthesis of a Potent NHE Inhibitor



C. Mechanistic Aspects of the Ni-Catalyzed Asymmetric Conjugate Allylation

The effect of ligand enantiomeric purity on selectivity in the conjugate allylation was examined to determine if a linear correlation would result.⁵⁵ Significantly, a slight negative non-linear effect was observed (Figure 2.6). Furthermore, the chemoselectivity

⁵⁵ For reviews of non-linear effects in asymmetric catalysis, see: (a) Kagan, H. B. *Adv. Synth. Catal.* **2001**, 343, 227. (b) Blackmond, D. G. *Acc. Chem. Res.* **2000**, 33, 402.

of the reaction was also affected by the enantiomeric purity of the ligand. That is, when racemic ligand was employed in the conjugate allylation, the highest chemoselectivity was observed (27:1 β' : β). In addition, the reaction using racemic catalyst was slightly faster than the conjugate allylation employing ligand of >99% ee (Figure 2.7). Clearly, heterochiral complexes form in the conjugate allylation reaction when the ligand enantiomeric purity is not 100%.

Figure 2.6: Effect of Ligand Enantiomeric Purity on Reaction Selectivity






Figure 2.7: Effect of Ligand Enantiomeric Purity on Reaction Rate



Since Pd has been shown to bind two phosphines when coordinated to dba,⁵⁶ formation of a (substrate)ML₂ complex during the reaction could explain these results. However, ³¹P NMR analysis of a 1:1:1 and a 1:2:1 mixture of Ni(cod)₂, chiral ligand **2.114**, and dba in C₆D₆ showed the presence of mostly free ligand (156.8 ppm), with a small, new singlet resonance at 166.0 ppm. No evidence for a ML₂ complex was observed. Therefore, a more likely explanation is the formation of [(**2.114**)Ni(substrate)]₂ dimers (Figure 2.8) analogous to [(PCy₃)Ni(dba)]₂ (**2.106**) described in section III.D (Figure 2.4, page 132). When **2.114** is used in varying enantiomeric purities, either **2.51** induced oxidative addition or dimer dissociation occurs at different rates and with different chemoselectivities for the heterochiral versus homochiral dimeric complexes shown in Figure 2.8. This rate difference produces a non-linear effect.

⁵⁶ (a) Amatore, C.; Jutland, A. *Coord. Chem. Rev.* **1998**, 178-180, 511. (b) Amatore, C.; Jutland, A.; Meyer, G. *Inorg. Chim. Acta.* **1998**, 273, 76. (c) Amatore, C.; Jutland, A.; Khalil, F.; M'Barki, M. A.; Mottier, L. *Organometallics* **1993**, 12, 3168.

Figure 2.8: Proposed Dimeric Complexes in the Asymmetric Conjugate Allylation



The effect of the metal:ligand ratio was next analyzed for the asymmetric conjugate allylation (Table 2.8). In the absence of ligand, the reaction was quite sluggish, and substantial amounts of the 1,2-addition product was observed (entry 1). In addition, exclusive conjugate allylation to the alkylidene unit of **2.62** was observed when using Ni(cod)₂ as catalyst alone. Addition of 0.5 equivalents of ligand relative to metal was enough to favor conjugate allylation over 1,2-allylation of **2.62** (entry 2). Furthermore, the addition of ligand caused a reversal in chemoselectivity, favoring the benzylidene allylation product in a 6.6:1 ratio. Notably, pronounced ligand acceleration⁵⁷ was observed considering that high conversion and enantioselection was obtained even in the excess of free Ni(cod)₂ relative to ligand. Addition of one equivalent of ligand relative to metal increased conversion, chemoselectivity, and enantioselectivity, while addition of excess ligand relative to metal did not offer any improvements.

⁵⁷ Berrisford, D. J.; Bolm, C.; Sharpless, K. B. *Angew. Chem. Int. Ed.* **1995**, *34*, 1059.


Table 2.8: Effect of Metal:Ligand Ratio in the Conjugate Allylation

entry	mol% ligand	1,4:1,2 ^a	β' : β ^b	% conv ^c	% yield ^d	% ee ^e
1	0	1:1.4	1:>50	38	25	n/a
2	2.5	5:1	6.6:1	82	59	95 (22)
3	5	>20:1	17:1	>95	69	95 (60)
4	7.5	>20:1	17:1	>95	80	94 (86)
5	10	>20:1	17:1	>95	77	93 (87)

^aValue determined by ¹H NMR analysis of the unpurified reaction mixture. ^bValue determined by GLC analysis of the unpurified reaction mixture. ^cValue represents the ratio of **2.64**:**2.62** in the ¹H NMR spectrum. ^dCombined yield of 1,2- and 1,4-addition products after silica gel chromatography. ^eValue for the major isomer; number in parentheses represents the value for the minor constitutional isomer.

Deuterium labeling experiments were performed employing the chiral catalyst used in the asymmetric conjugate allylation and **2.51**-d₂ (Scheme 2.25). As was observed in the conjugate allylation of **2.62** with **2.51**-d₂ using Ni(cod)₂/PCy₃ as catalyst, deuterium scrambling was also observed in the asymmetric conjugate allylation. Thus η^1 - η^3 - η^1 allyl isomerization is likely faster than reductive elimination with the chiral catalyst as discussed in section III.D (Scheme 2.22, page 131).

Scheme 2.25: Deuterium Labeling Experiments



D. Model for Chemoselectivity in the Conjugate Allylation

Using the chemoselectivity data obtained for the ligand surveys in both the enantioselective and non-enantioselective conjugate allylation reaction, a working model for chemoselectivity was developed. Displayed in Table 2.9 is a summary of this data, giving the correlation between ligand basicity and chemoselectivity in the conjugate allylation. The CO stretching frequencies for the *trans*-(ligand)₂Rh(CO)Cl complex prepared from each ligand was used as a gauge of ligand basicity.⁵⁸ When achiral ligands were employed, chemoselectivity was dependent on the Lewis basicity of the achiral ligand. That is, as the Lewis basicity of the ligand decreased, the chemoselectivity decreased (entries 1-5). Use of highly Lewis basic PCy₃ gave the highest chemoselectivity for alkylidene allylation (entry 1). Furthermore, ligand size did not have a significant affect when using achiral phosphine ligands (entries 4 and 5). Tri(*ortho-tert*-butylphenyl)phosphite and triphenylphosphite gave comparable levels of chemoselectivity, yet the Tolman cone angle for P(O-2-*t*BuPh)₃ is significantly larger

⁵⁸ (a) Vastag, S.; Heil, B.; Markó, L. *J. Mol. Cat.* **1979**, *5*, 189. (b) Ohgomori, Y.; Yoshida, S.-I.; Watanabe, Y. *J. Chem. Soc., Dalton Trans.* **1987**, 2969. (c) Otto, S.; Roodt, A. *Inorg. Chim. Acta* **2004**, *357*, 1. (d) Fernández, E.; Ruiz, A.; Claver, C.; Castellón, S.; Polo, A. *Organometallics* **1998**, *17*, 2857. (e) Burks, H. E.; Liu, S.; Morken, J. P. *J. Am. Chem. Soc.* **2007**, *129*, 8766.

relative to P(OPh)₃ (175° vs 128°, respectively).⁵⁹ However, as can be noted in entries 4 and 5 (Table 2.9), these ligands still fit the ligand basicity trend. Interestingly, TADDOL-derived chiral phosphines did not fit this Lewis basicity trend (entry 6). The electronic nature of TADDOL-derived phosphoramidite **2.110** is similar to PPh₃, yet very high chemoselectivity was realized using this ligand relative to PPh₃, which was relatively non-selective (entry 3 vs 6). Presumably, there is a steric preference with the chiral ligand, which leads to this enhanced chemoselectivity. Since achiral ligands do not appear to effect chemoselectivity based on sterics, the chirality of the TADDOL backbone has a dramatic effect on chemoselectivity that cannot be explained by simple cone angle arguments.

Table 2.9: Correlation Between Ligand Basicity and Chemoselectivity

Reaction scheme: $\text{Ph}-\text{CH}=\text{CH}-\text{C}(=\text{O})-\text{CH}=\text{CH}-n\text{-pentyl}$ (2.62) reacts with ligand **2.51** (1.2 equiv) in the presence of 10% Ni(cod)₂ in THF at rt for 4 h to give $\text{Ph}-\text{CH}=\text{CH}-\text{C}(=\text{O})-\text{CH}(\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3)-\text{CH}=\text{CH}_2$ (2.63, β -addition) and $\text{Ph}-\text{CH}(\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3)-\text{CH}=\text{CH}-\text{C}(=\text{O})-\text{CH}=\text{CH}_2$ (2.64, β' -addition).


entry	ligand	ν_{CO} (cm ⁻¹) ^a	$\beta':\beta$	% yield
1	PCy ₃	1943	1:7.5	82
2	P(NMe ₂) ₃	1964	1:2.5	82
3	PPh ₃	1979	1:2.4	49
4	P(O-2- <i>t</i> BuPh) ₃	2012	1:1.5	40
5	P(OPh) ₃	2016	1:1.1	55
6	2.110	1984	>20:1	53

^aThese values represent that of the *trans*-L₂Rh(CO)Cl complex. See footnote 58.

⁵⁹ Tolman, C. A. *Chem. Rev.* **1977**, 77, 313.

The above observations can best be accommodated by considering the stability of the π -allyl complexes (**2.152** vs **2.154**, Scheme 2.26). This argument assumes that π -allyl stability is product-determining. In support of this hypothesis, electron-rich metals, where the π -allyl bearing the electron-withdrawing aryl ring should be favored (**2.154**), lead to the alkylidene allylation product via [3,3']-reductive elimination. This was the observed reaction outcome using Lewis basic PCy₃. Furthermore, steric analysis of **2.152** and **2.154** predicts that when the alkyl group is small, **2.152** should predominate to minimize steric interactions with the ligand (L). While this was not observed for achiral ligands, TADDOL-derived phosphines appear to effect chemoselectivity based on this steric preference, and therefore, afford arylidene allylation. As was observed during optimization of the substrate (Table 2.5, page 137), increasing the size of the alkyl group led to decreased chemoselectivity for arylidene allylation. This result supports a steric model when using chiral TADDOL-derived phosphonite **2.114**. Moreover, the substrate scope study for the asymmetric conjugate allylation revealed that smaller aryl groups led to decreased chemoselectivity (Table 2.7, entry 1 vs 10, page 140). In addition, *ortho*-substitution on the aryl group of the substrate led to increased chemoselectivity in the reaction (Table 2.7, entry 1 vs 4, 5, and 7). These observations are consistent with a steric argument grounded on π -allyl complex stability.

Scheme 2.26: Model for Chemoselectivity in the Conjugate Allylation



While the chemoselectivity in the asymmetric conjugate allylation always favors arylidene allylation, subtle changes in the electronic nature of the arylidene unit of the substrate affected chemoselectivity. Arguably, $d\pi\text{-}\pi^*$ donation from the metal to the aryl-ring of the substrate, similar to the dba-effect⁶⁰ recognized in cross-coupling reactions, may add to the stabilization of **2.154**. Thus, when the aryl group of the substrate is electron-withdrawing, this $d\pi\text{-}\pi^*$ donation should be enhanced, leading to more alkylidene allylation. The reverse should be true for electron-rich aryl groups, and this trend was observed in the conjugate allylation: when substrates with electron-withdrawing substituents on the aryl ring were used in the asymmetric conjugate allylation, lower chemoselectivities were obtained (i.e. more alkylidene allylation, Table 2.7, entry 1 vs 2 and 6, page 140). Likewise, substrates bearing electron-donating substituents on the aryl ring gave more arylidene allylation (Table 2.7, entry 1 vs 3 and 8).

⁶⁰ (a) Mace, Y.; Kapdi, A. R.; Fairlamb, J. S.; Jutand, A. *Organometallics* **2006**, 25, 1795. (b) Fairlamb, I. J. S.; Kapdi, A. R.; Lee, A. F. *Org. Lett.* **2004**, 6, 4435.

In summary, for achiral ligands, the Lewis basicity appears to control chemoselectivity in the allylation. However, with chiral TADDOL-derived ligands, sterics appear to be more significant in controlling chemoselectivity.

E. Extension to an Asymmetric Chemoselective Alkylidene Conjugate Allylation

The general asymmetric conjugate allylation described in Table 2.7 (page 140) provides a solution for asymmetric conjugate allylation to furnish benzylic stereocenters; however, formation of aliphatic stereocenters is also desirable. Because of the difference in reactivity observed with Pd vs Ni-catalysts, various chiral TADDOL-derived phosphine ligands were examined, using $\text{Pd}_2(\text{dba})_3$ as the precatalyst, with the goal of developing an asymmetric alkylidene conjugate allylation (Table 2.10). Modification of the amino moiety of the ligand did not improve enantioselectivity (entries 1-3); however, modification of the TADDOL-backbone led to increased enantioselectivities (entries 4 and 5). While this survey did identify a chiral catalyst that afforded the alkylidene allylation product in excellent enantiomeric purity (entry 5), no catalyst system gave useful levels of chemoselectivity for alkylidene allylation. In fact, as observed under Ni-catalysis, chiral Pd-catalysts bearing TADDOL-derived phosphine ligands generally favored arylidene allylation.

Table 2.10: Chiral Catalyst Survey with Pd₂(dba)₃ as Precatalyst

Reaction scheme: Enone **2.62** (Ph-CH=CH-C(=O)-CH=CH-n-pentyl) reacts with allylboronate ester **2.51** (1.2 equiv) in the presence of 2.5% Pd₂(dba)₃, 6% ligand, THF, rt. The reaction yields two products: **2.63** (β-addition) and **2.64** (β'-addition).

entry	ligand ^a	β:β' ^b	% yield ^c	% ee ^d
1	2.107	1:3.4	76	25 (12)
2	2.108	1:26	66	-14 (8)
3	2.109	1:3.4	63	38 (25)
4	2.110	1:2	80	79 (32)
5	2.111	1.6:1	74	96 (33)
6	2.112	1:17	68	83 (2)
7	2.113	1:7.8	82	39 (19)
8	2.114	1:11	74	46 (16)

^aSee Figure 2.5 for ligand structures (page 135). ^bValue determined by ¹H NMR analysis of the unpurified reaction mixture. ^cIsolated yield after silica gel chromatography. Isolated as a mixture of constitutional isomers ^dValue for the β-isomer, value in parentheses represents that of the β'-isomer.

To solve this chemoselectivity issue, it was reasoned that the aryl-group in **2.62** could be used to control chemoselectivity in the conjugate allylation. This strategy was based on the chemoselectivity model discussed in section IV.D (page 148). Since smaller aryl groups should increase alkylidene allylation, we reasoned that replacement of the phenyl ring in **2.62** with a proton should allow for maximum alkylidene allylation. Therefore, **2.156** was prepared (using carbonylative Stille-coupling⁶¹) and surveyed in the conjugate allylation (Table 2.11). Under Ni-catalysis, good chemoselectivity could

⁶¹ Goure, W. F.; Wright, M. E.; Davis, P. D.; Labadie, S. S.; Stille, J. K. *J. Am. Chem. Soc.* **1984**, *106*, 6417.

be obtained in some cases; however, enantioselectivity was never of acceptable quality (entries 1-5). Interestingly, the chiral catalyst developed for the asymmetric arylidene conjugate allylation gave only modest enantioselectivity (entry 1). The use of Pd as catalyst did consistently give high chemoselectivity for the desired constitutional isomer (entries 6-15). TADDOL-derived phosphoramidites proved to be the best ligand class for the reaction with **2.156** (entries 6-10). Again, substitution on the aromatic groups of the TADDOL-scaffold was required for enantioselection. Phosphoramidites with 3,5-disubstituted phenyl groups on the TADDOL backbone all gave higher enantioselectivities relative to the parent phenyl compound (compare entries 6 and 7 with entries 8-12). Furthermore, phosphonites (entries 12-14) and phosphites (entry 15) were inferior in the conjugate allylation of **2.156**. It was clear from these results that a new ligand structure was needed for successful allylation of **2.156** with high enantioselection.

Table 2.11: Chiral Catalyst Survey with 2.156 as Substrate


entry	metal	ligand ^a	$\beta:\beta'$ ^b	% yield ^c	% ee
1	5% Ni(cod) ₂	10% 2.114	14:1	21	54
2	5% Ni(cod) ₂	10% 2.115	9:1	31	45
3	5% Ni(cod) ₂	10% 2.111	2.4:1	32	1
4	5% Ni(cod) ₂	10% 2.112	1.5:1	44	0
5	5% Ni(cod) ₂	10% 2.113	0:1	n.d.	n/a
6	2.5% Pd ₂ (dba) ₃	6% 2.107	15:1	45	-7
7	2.5% Pd ₂ (dba) ₃	6% 2.109	19:1	38	9
8	2.5% Pd ₂ (dba) ₃	6% 2.110	11:1	3	-47
9	2.5% Pd ₂ (dba) ₃	6% 2.111	20:1	15	32
10	2.5% Pd ₂ (dba) ₃	6% 2.112	35:1	10	50
12	2.5% Pd ₂ (dba) ₃	6% 2.115	22:1	43	2
13	2.5% Pd ₂ (dba) ₃	6% 2.116	17:1	38	-41
14	2.5% Pd ₂ (dba) ₃	6% 2.114	17:1	22	-5
15	2.5% Pd ₂ (dba) ₃	6% 2.113	32:1	1	-8

^aSee Figure 2.5 for ligand structures (page 135). ^bValue determined by GLC analysis of the unpurified reaction mixture. ^cIsolated yield after silica gel chromatography. Isolated as a mixture of constitutional isomers.

From the data displayed in Table 2.11, it was determined that TADDOL-derived phosphoramidites, containing 3,5-disubstituted aryl rings on the TADDOL backbone, were promising ligands and should be explored further. In addition, an increase in enantioselectivity was observed with increasing size of the amino moiety of the

phosphoramidite (Table 2.11, entries 9 and 10). Therefore, the TADDOL-derived phosphoramidite ligands in Figure 2.9 were synthesized to find a competent chiral ligand for this transformation. These phosphoramidite ligands were rationally designed to possess sterically more demanding amino groups.

Figure 2.9: New Ligands for Asymmetric Alkylidene Allylation



The results for the conjugate allylation of **2.156** when using these newly prepared ligands are tabulated in Table 2.12. When increasing the size of the amino moiety on the phosphoramidite from a 6- to 7-membered heterocycle, little improvement was observed in the enantioselectivity; however, chemoselectivity improved dramatically (Table 2.12: compare entries 1 and 2, and compare Table 2.11, entry 10 with Table 2.12, entry 3).

Gratifyingly, introduction of a methyl group onto the heterocycle (**2.162**) provided a solution (Table 2.12, entry 4). Thus in the presence of phosphoramidite **2.162**, the conjugate allylation of **2.156** proceeded in excellent enantioselectivity and chemoselectivity in modest yield. Furthermore, the opposite diastereomer of ligand (**2.163**) was the mismatched case, affording the conjugate allylation product in excellent chemoselectivity, but with poor enantioselectivity (entry 5).

Table 2.12: Chiral Catalyst Survey with Newly Prepared Phosphoramidites

Reaction scheme: Enone **2.156** (with β' and β positions labeled) reacts with allylboronate **2.51** (1.2 equiv) in the presence of a catalyst in PhMe at room temperature to yield β -addition product **2.157** and β' -addition product **2.158**.

entry	metal	ligand ^a	β : β' ^b	% yield ^c	% ee
1	2.5% Pd ₂ (dba) ₃	6% 2.159	14:1	16	-29
2	2.5% Pd ₂ (dba) ₃	6% 2.160	21:1	7	-24
3	2.5% Pd ₂ (dba) ₃	6% 2.161	63:1	50	52
4	2.5% Pd ₂ (dba) ₃	6% 2.162	>50:1	39	86
5	2.5% Pd ₂ (dba) ₃	6% 2.163	>50:1	47	-44

^aSee Figure 2.9 for ligand structures (page 156). ^bValue determined by GLC analysis of the unpurified reaction mixture. ^cIsolated yield after silica gel chromatography. Isolated as a mixture of constitutional isomers.

With a chiral catalyst identified for alkylidene conjugate allylation, a short solvent screen was conducted to determine if enantioselection could be enhanced (Table 2.13). Etheral solvents were better in terms of chemical yield; however, all solvents examined

gave comparable levels of enantioselection. THF was determined to be the best choice of solvent for the reaction.

Table 2.13: Solvent Effects in the Asymmetric Conjugate Allylation of 2.156

entry	solvent	$\beta:\beta'$ ^a	% yield ^b	% ee
1	THF	>50:1	52	86
2	Et ₂ O	>50:1	48	87
3	<i>t</i> -BuOMe	>50:1	41	85
4	<i>o</i> -DCB	>50:1	42	84

^aValue determined by GLC analysis of the unpurified reaction mixture. ^bIsolated yield after purification by silica gel chromatography.


After successfully developing a catalyst for chemoselective alkylidene allylation, a method for the formation of all-carbon quaternary stereocenters was sought. Initially, **2.164** was prepared and examined in the conjugate allylation (Table 2.14). Unfortunately, when employing the ligand identified for the asymmetric allylation of **2.156** in the conjugate allylation of **2.164**, only modest enantioselectivity was obtained in conjunction with poor conversion (entry 1). Increasing the reaction temperature to 70 °C did not improve reaction efficiency (entry 2). Furthermore, the use of other phosphoramidites, under Pd-catalysis, also gave both poor conversion and

enantioselectivity (entries 3-6). Conversion was slightly better with phosphoramidites containing smaller amino moieties (entries 4 and 5); however, this led to decreased enantioselectivities. Moreover, a phosphite bearing a bulky *t*-butyl group also gave poor conversion and enantioselectivity (entry 7). Increasing the reaction temperature and/or time in these reactions did not increase reaction conversion and often gave unidentified side-products. Interestingly, use of the chiral Ni-catalyst developed for the asymmetric arylidene allylation gave high conversion, but substantial amounts of the undesired constitutional isomer (entry 8).

Table 2.14: Attempts at Formation of All-Carbon Quaternary Stereocenters

entry	metal	ligand	% conv ^a	β:β' ^a	% ee
1 ^b	2.5% Pd ₂ (dba) ₃	6% 2.162 ^c	2	>50:1	73
2	2.5% Pd ₂ (dba) ₃	6% 2.162 ^c	9	>50:1	63
3	2.5% Pd ₂ (dba) ₃	6% 2.163 ^c	6	>50:1	-33
4	2.5% Pd ₂ (dba) ₃	6% 2.110 ^d	15	>50:1	-13
5	2.5% Pd ₂ (dba) ₃	6% 2.111 ^d	18	>50:1	33
6	2.5% Pd ₂ (dba) ₃	6% 2.166	7	>50:1	51
7	2.5% Pd ₂ (dba) ₃	6% 2.167	14	>50:1	-27
8	5% Ni(cod) ₂	10% 2.114 ^d	98	1:6.7	50


^aValue determined by GLC analysis of the unpurified reaction mixture. ^bReaction run at rt for 24 h. ^cSee Figure 2.9 (page 156). ^dSee Figure 2.5 (page 135).



Further analysis of the conjugate allylation with **2.164** showed that Ni-catalysts were more active than their Pd counterparts, giving high conversion (Scheme 2.27). Unfortunately, the undesired constitutional isomer was formed as the major product in high chemoselectivity. Assuming that oxidative addition (π -allyl formation) occurs to the less hindered β' -site of **2.164**, it is likely that direct reductive elimination to the β' -


position is now competitive with [3,3']-reductive elimination to the β -position. Rate enhancement that is typically provided by [3,3']-reductive elimination is likely lost because of the steric hindrance associated with forming an all-carbon quaternary stereocenter. Supporting the hypothesis that direct reductive elimination may occur, is the observation that simple alkylboron reagents have been used as nucleophiles in the conjugate addition of acrylates under Ni(0) catalysis.^{24a}

Scheme 2.27: Reaction of 2.164 with Ni(cod)₂/PPh₃




To solve this chemoselectivity issue when using Ni as the catalyst, it was considered that reintroduction of the aryl group in the dialkylidene ketone substrate may slow down direct reductive elimination and allow for [3,3']-reductive elimination to be competitive again. Bolstering this idea, was the observation that Ni/PCy₃ gave chemoselective conjugate allylation to form an all-carbon quaternary center when **2.72** was used as substrate (Table 2.3, entry 8, page 120). Thus, **2.169** was prepared and analyzed as a substrate for asymmetric conjugate allylation (Scheme 2.28). While chemoselectivity did improve when using **2.169** as substrate, both isolated yield and enantioselectivity were poor.

Scheme 2.28: Attempts at an Asymmetric Conjugate Allylation of 2.169



An important observation in the conjugate allylation of **2.169** was that the trisubstituted olefin of the recovered starting material had isomerized under the reaction conditions. Based on our understanding of this conjugate allylation process, we had hoped that quaternary stereocenters bearing groups of similar size could be prepared in high enantioselectivity. That is to say, successful asymmetric conjugate allylation of **2.169** would formally result from differentiation of a methyl and a butyl group by a chiral catalyst. Since these groups are similar in size, such a process is difficult. However, it is possible that the chiral catalyst dictates which enantiotopic face of the intermediate π -allyl it will bind (i.e. **2.171** vs **2.172**, Scheme 2.29). If this is the case, and the [3,3']-reductive elimination is stereospecific, then the activating alkene of the substrate should dictate the enantioselectivity in the reaction and not the substituents adjacent to the carbon being allylated. If this contention is accurate, then the geometry of the trisubstituted olefin in **2.169** should control which enantiomer is formed in the conjugate allylation. *Thus, a chiral catalyst for successful asymmetric conjugate allylation of 2.169 must not isomerize the trisubstituted olefin during the conjugate allylation.*

Scheme 2.29: Possible Mode of Asymmetric Induction in the Conjugate Allylation



In an attempt to increase the rate of conjugate allylation relative to starting material isomerization, the reaction in Scheme 2.28 was performed using 3.0 equiv of **2.51** at 1 M concentration relative to **2.169**. Isomerization of recovered **2.169** was less in this case (~5:1); however, **2.170** was isolated in only 31% yield, 9:1 chemoselectivity, and 45% ee. Furthermore, improvement in conversion, as evident by ^1H NMR analysis of the unpurified reaction mixture, was not observed despite the slight increase in yield. If enantioselectivity in the reaction is controlled by the model given in Scheme 2.29, then one would predict that **2.170** should be obtained in ~86% ee. This conclusion is based on the enantiomeric purity of the alkylidene allylation product resulting from allylation of **2.62** when using $\text{Ni}(\text{cod})_2$ /**2.114** as catalyst (Table 2.8, entry 4, page 147). This is clearly not the case in the above example, even when alkene isomerization in **2.169** had been reduced. The difference in temperature between the present reaction and the asymmetric allylation of **2.62**, given in Table 2.8, may be the result for this discrepancy. However, this has yet to be determined since the allylation reaction of **2.169** does not proceed at room temperature when $\text{Ni}(\text{cod})_2$ and **2.114** are employed. Another possible explanation

is that steric interaction between the methyl group of the substrate and the allyl ligand of the metal result in lower enantioselectivity.


The formation of all-carbon quaternary stereocenters via asymmetric conjugate allylation remains an unsolved problem. From the experiments discussed above, it is likely that a catalyst system which affects conjugate allylation at room temperature may be required in order to achieve high enantioselectivities. No chiral Pd or Ni catalysts prepared from chiral TADDOL-derived phosphonites, phosphoramidites, or phosphites were found to form quaternary stereocenters at room temperature. However, because the activating olefin in the substrate can be used as a tuning element, the addition of electron withdrawing substituents on the phenyl ring in **2.169** may further lower the barrier for reductive elimination by withdrawing electron density from the metal center. This may allow the conjugate allylation to proceed at reduced temperatures. Application of this strategy has not been carried out; therefore, its relevance is pure conjecture at this point.

F. Other Related Reactions

As discussed in section III.A (page 113), typical unsaturated electrophiles (Figure 2.1, page 115) were inert to the catalytic conjugate allylation described herein. However, another class of α,β -unsaturated electrophiles that were found to be reactive were α,β -unsaturated α -ketoesters (**2.173**, Scheme 2.30). Surprisingly, these substrates gave complete 1,2-addition (eq 1). While it was not determined why these substrates reacted in a 1,2-fashion, rather than the expected 1,4-sense, it is likely that these substrates can

react analogously to the well known Tsuji-allylation (Scheme 2.30, eq 3).⁶² In the Tsuji reaction, allyl enol carbonate **2.177** is converted to α -allylcyclohexanone (**2.179**) via intermediate **2.178**. In an analogous manner, the bis(allyl)Pd complex **2.175** (eq 2), in the allylation of **2.173**, may be in equilibrium with Pd-enolate complex **2.176**, which is remarkably similar to **2.178**. Reductive elimination at the α -carbon of the extended ester enolate offers a rationale for the production of **2.174**.

Scheme 2.30: Use of an α,β -Unsaturated α -Ketoesters




Importantly, the 1,2-addition product of **2.173** was not derived from background reaction between **2.173** and **2.51**: control experiments verified that both metal and phosphine ligand were required for successful allylation. Further, performing the

⁶² (a) Tsuji, J.; Minami, I.; Shimizu, S. *Tetrahedron Lett.* **1983**, 24, 1793. (b) Minami, I.; Nisar, M.; Yuhara, M.; Shimizu, I.; Tsuji, J. *Synthesis* **1987**, 992.


allylation in the presence of chiral phosphoramidite ligand **2.110** led to formation of **2.174** in good yield with some enantiocontrol (Scheme 2.31).

Scheme 2.31: Enantioselective Allylation of 2.173




In addition to allylation, crotylation of **2.173** was also successful using Pd-catalysis (Scheme 2.32). The use of *Z*-configured crotylboronate **2.180** led to the γ -crotylation product **2.181** as a single diastereomer in good yield (eq 1). The use of the *E*-configured crotylboronate led to the same diastereomer of product; however, much longer reaction time was needed (eq 2). Interestingly, the use of THF as solvent provided a dramatic increase in rate, allowing for the crotylation to be performed at room temperature with excellent yield (eq 3). It should be noted that aliphatic α,β -unsaturated α -ketoesters were not allylated or crotylated under the reaction conditions described here. Competitive β -hydride elimination appeared to be problematic when these electrophiles were used.

Scheme 2.32: Catalytic Crotylation of **2.173**




Similar to crotylboronates, the allene diboration products participated in the catalytic allylation of **2.173** (Scheme 2.33). A one-pot tandem diboration/allylation/oxidation gave **2.182** as a single diastereomer in good yield (eq 1). Again, 1,2-addition to **2.173** was observed. Interestingly, when the same sequence was performed using the optimal chiral ligand for the asymmetric allene diboration, very little of the allylation/oxidation product was isolated (eq 2). Furthermore, no enantioinduction was observed. However, treatment of **2.173** with purified, enantiopure **2.48**, using $\text{Pd}_2(\text{dba})_3/\text{P}(\text{tPr})_3$ as the catalyst in the allylation, gave good yield of **2.182**, after oxidation, with moderate chirality transfer (eq 3). Clearly, this suggests that the chirality of the enantiomerically pure catalyst in eq 2 is mismatched with the chirality of **2.48** in the subsequent catalytic allylation. Thus, a tandem one-pot diboration/allylation process could not be developed.

Scheme 2.33: Catalytic Addition of a Diboronate Esters to 2.173



Finally, in addition to unsaturated α -ketoesters, $\alpha,\beta,\gamma,\delta$ -unsaturated enone **2.183** also underwent allylation under Ni-catalysis (Scheme 2.34). Due to the fact that the (*E,Z*)-diene (**2.184**) was isolated as the reaction product, this class of substrates arguably reacts through a [3,3']-reductive elimination pathway (**2.185**) to arrive at the observed product. Use of a chiral TADDOL-derived phosphonite afforded the allylation product in modest enantioselectivity, verifying the involvement of the chiral catalyst.

Scheme 2.34: Catalytic Allylation of an $\alpha,\beta,\gamma,\delta$ -Unsaturated Enone



V. Conclusions

The discovery and development of a transition-metal-catalyzed asymmetric conjugate allylation has been reported. The following insights have been gained from this research:

1. Allylboronate nucleophiles readily add to the β -position of dialkylidene ketones when group 10 transition metals are used as catalysts. Furthermore, this reactivity is unique to dialkylidene ketones. This enhancement of reactivity is postulated to occur due to a facile [3,3']-reductive elimination pathway that can operate for this class of substrates, which is not accessible to other α,β -unsaturated electrophiles.
2. Under Ni(0)-catalysis, the use of Lewis basic ligands leads to chemoselective conjugate allylation to the alkylidene site in unsymmetrical dialkylidene ketones,

while the use of chiral TADDOL-derived ligands affords chemoselective and enantioselective arylidene allylation.

3. The alkyl and aryl groups on the activating alkene in unsymmetrical dialkylidene ketones can be used as a control element to affect both chemoselectivity and enantioselectivity.
4. A catalytic asymmetric conjugate allylation that affords enantiomerically enriched conjugate allylation products from arylidene or alkylidene allylation has been described.
5. The conjugate allylation products can be further manipulated into valuable organic synthons.
6. Formation of all-carbon quaternary stereocenters via conjugate allylation is feasible; however, a competent chiral catalyst has not been found.
7. An α,β -unsaturated α -ketoester and an $\alpha,\beta,\gamma,\delta$ -unsaturated enone also participate in allylation reactions with allylboronates under group 10 metal catalysis. These electrophiles afford 1,2-allylation products rather than 1,4-allylation products.

VI. Experimental Procedures

A. General. Melting points were determined using a Mel-Temp II melting point apparatus and are uncorrected. ^1H NMR spectra were recorded on Inova-500 (500 MHz), Gemini-400 (400 MHz), Unity-300 (300 MHz), or Bruker DRX 300 or 400 MHz spectrometers. Chemical shifts are reported in ppm from tetramethylsilane with the residual solvent resonance as an internal standard (CHCl_3 : 7.24 ppm, C_6HD_5 : 7.15 ppm). Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, h = hextet, br = broad, m = multiplet), and coupling constants (Hz). ^{13}C NMR spectra were recorded on a Bruker DRX 400 (100 MHz) instrument, a Gemini-400 (100 MHz) instrument, or an Inova-500 (125 MHz) instrument with complete proton decoupling. Chemical shifts are reported in ppm from tetramethylsilane with the solvent as the internal standard (CDCl_3 : 77.0 ppm, C_6D_6 : 128.39 ppm). ^{19}F NMR spectra were recorded on an Inova-500 (470 MHz) instrument. Chemical shifts are reported in ppm and are referenced to a 0.05% solution of C_6F_6 (-63.72 ppm) in C_6D_6 . ^{31}P NMR spectra were recorded on a Unity-300 (121 MHz), a Bruker 400 MHz (161 MHz), or an Inova-500 (202 MHz) instrument with complete proton decoupling. Chemical shifts are reported in ppm and are referenced to 85% H_3PO_4 (0.0 ppm). Low-resolution mass spectrometry was performed by the University of North Carolina, Department of Chemistry Mass Spectrometry Facility and by the Boston College, Department of Chemistry Mass Spectrometry Facility. Infrared (IR) spectra were obtained on Nicolet 210 or 560 spectrophotometer. Optical rotations were measured using a Rudolf Research Analytical Autopol IV Polarimeter.

Liquid chromatography was performed using forced flow (flash chromatography) on silica gel (SiO_2 , 40-63 μm) purchased from Silicycle. Thin layer chromatography (TLC) was performed on 250 μm silica gel plates from Silicycle or 200 μm neutral alumina TLC plates from Sorbent Technologies. Visualization was achieved using UV light, phosphomolybdic acid in ethanol, potassium permanganate in water, or cerium(IV) sulfate and ammonium molybdate in sulfuric acid, each followed by heating.

Analytical chiral gas-liquid chromatography (GLC) was performed on a Hewlett-Packard 6890 series chromatograph equipped with a CTC Analysis Combi Pal autosampler by Leap Technologies (Carrboro, NC), a split mode capillary injection system, a flame ionization detector, and a Supelco β -dex 120 column with helium as the carrier gas. The inlet pressure and split ratio were 25 psi and 35, respectively, in all cases unless otherwise noted. Analytical achiral GLC was performed on a Hewlett-Packard 6890 series chromatograph equipped with a split mode capillary injection system, a flame ionization detector, and a Hewlett-Packard Ultra 1 capillary column (0.33 μm film thickness, 25 m length, 0.2mm ID) with helium as the carrier gas. The inlet pressure and split ratio were 25 psi and 15, respectively, in all cases unless otherwise noted. Analytical chiral supercritical fluid chromatography (SFC) was performed on a Berger Instruments supercritical chromatograph equipped with an Alcott autosampler and a Knauer UV detector. Analytical chiral high performance liquid chromatography (HPLC) was performed on a Shimadzu instrument with UV detector.

All reactions were conducted in oven- or flame-dried glassware under an inert atmosphere of nitrogen or argon. Toluene, toluene- d_8 , and benzene- d_6 were distilled over

CaH₂ and degassed by freeze-pump-thaw cycles prior to use. Anhydrous tetrahydrofuran (THF), used in reactions performed outside a dry-box, was freshly distilled from Na metal and benzophenone or purified using a Pure Solv MD-4 solvent purification system from Innovative Technology Inc. by passing the solvent through two activated alumina columns after being purged with Ar. Anhydrous THF, used in reactions that were prepared in a dry-box, was purchased from Aldrich Chemical Company. Methylene chloride and diethyl ether were purified using a Pure Solv MD-4 solvent purification system from Innovative Technology Inc. by passing the solvent through two activated alumina columns after being purged with Ar.

Tris(dibenzylideneacetone)dipalladium(0) and bis(1,5-cyclooctadiene)nickel(0) were purchased from Strem Chemical Company. Allylboronic acid pinacol ester (**2.51**) was purchased from Aldrich Chemical Company and distilled through a 6 inch Vigreux column (58-62 °C at 20 torr) and stored in the freezer under Ar. Allenes and allene diboration products were prepared according to the methods described in Chapter 1 (see page 47). 1-Iododheptene was synthesized from 1-heptyne via hydroalumination with DIBAL followed by iodination.¹ 2-Iodovinylcyclohexane and 1-iodo-3,3-dimethyl-1-butene were synthesized via hydroboration of cyclohexylacetylene or *t*-butylacetylene, respectively, followed by iodination.² Vinyl iodides bearing pendant TBS-protected alcohols were synthesized via hydrozirconation of the TBS-protected propargyl or homo

¹ (a) Stille, J. K.; Simpson, J. H. *J. Am. Chem. Soc.* **1987**, *109*, 2138. (b) Trost, B. M.; Rudd, M. T. *Org. Lett.* **2003**, *5*, 4599.


² (a) Brown, H. C.; Hamaoka, T. Ravindran, N.; Subrahmanyam, C.; Somayaji, V.; Bhat, N. G. *J. Org. Chem.* **1989**, *54*, 6075. (b) Gagnon, D.; Lauzon, S.; Godbout, C.; Spino, C. *Org. Lett.* **2005**, *7*, 4769.

propargyl alcohol with the Schwartz reagent, followed by iodination.³ Vinyl bromides were purchased from Aldrich Chemical Company and used without further purification. 5-Fluoro-2-methylbenzaldehyde was purchased from Oakwood Chemicals and used without further purification. All other reagents were purchased from either Fisher or Aldrich Chemical Companies and used directly.

B. Experimental Procedures.

1. Weinreb Amide Syntheses


General procedure for synthesis of Weinreb amides:



To 11.7 g (79.0 mmol) of *trans*-cinnamic acid (**2.186**) and 11.6 g (119 mmol) of (*N,O*)-dimethylhydroxylamine hydrochloride in 316 mL of CH₂Cl₂ in a 1 L round-bottom flask with magnetic stir-bar was added 1.93 g (15.8 mmol) of 4-(dimethylamino)pyridine (DMAP) followed by 22.8 g (119 mmol) of *N*-(3-dimethylaminopropyl)-*N*′-ethylcarbodiimide hydrochloride (EDCI), all in one portion. This mixture was put under N₂ and 15.7 mL (12.0 g, 119 mmol) of triethylamine was added dropwise over a 10 min period. After addition, the mixture was allowed to stir overnight at ambient temperature. The final mixture was transferred to a separatory funnel, washed with 3 M HCl (2x) followed by saturated aqueous NaHCO₃. The organic layer was dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. Silica gel chromatography


³ Germain, J.; Deslongchamps, P. *J. Org. Chem.* **2002**, 67, 5269.

(hexanes:EtOAc) afforded 13.1 g (68.5 mmol, 87%) of Weinreb amide **2.65** as a white solid. Spectral data were consistent with the literature.⁴



(E)-3-(3-Chlorophenyl)-N-methoxy-N-methylacrylamide (2.187).


Prepared according to the general procedure on page 174 in 85% yield. A white solid. Mp 48-52 °C. R_f = 0.29 (SiO₂, 2:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3069 (m), 2974 (m), 2930 (m), 2823 (m), 1659 (s), 1615 (s), 1564 (s), 1469 (s), 1425 (s), 1389 (s), 1199 (s) (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.62 (1H, d, J = 16 Hz), 7.52 (1H, s), 7.34-7.42 (1H, m), 7.22-7.32 (2H, m), 6.99 (1H, d, J = 16 Hz), 3.74 (3H, s), 3.27 (3H, s); ¹³C NMR (CDCl₃): δ 166.3, 141.7, 136.9, 134.6, 129.9, 129.6, 127.4, 126.4, 117.1, 61.91, 32.45. LRMS (ESI+) Calcd for C₁₁H₁₂ClNO₂ (M)⁺: 225.1, Found (M)⁺: 225.6.



(E)-N-Methoxy-N-methyl-3-*o*-tolylacrylamide (2.188).

Prepared according to the general procedure on page 174 in 90% yield. A colorless oil. R_f = 0.35 (SiO₂, 1:1 hexanes:EtOAc); IR (neat): 3062 (w), 2968 (m), 2936 (m), 1659 (s), 1615 (s), 1464 (m), 1413 (m), 1388 (s), 1180 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.99 (1H, d, J = 16 Hz), 7.57 (1H, d, J = 8.0 Hz), 7.12-7.26 (3H, m), 6.92 (1H, d, J = 16 Hz), 3.72 (3H, s), 3.28 (3H, s), 2.41 (3H, s); ¹³C NMR (CDCl₃): δ 166.8, 141.0, 137.5, 134.0, 130.5, 129.4, 126.1, 126.0, 116.8, 61.75, 32.39, 19.76. LRMS (ESI+) Calcd for C₁₂H₁₅NO₂ (M)⁺: 205.1, Found (M)⁺: 205.6.


⁴ Hiyama, T.; Reddy, G. B.; Minami, T.; Hanamoto, T. *Bull. Chem. Soc. Jpn.* **1995**, 68, 350.



(E)-N-Methoxy-N-methyl-3-[2-

(trifluoromethyl)phenyl]acrylamide (2.189). Prepared according


to the general procedure on page 174 in 48% yield. A colorless oil. $R_f = 0.15$ (SiO₂, 3:1 hexanes:EtOAc); IR (neat): 3075 (w), 2974 (m), 2936 (m), 2823 (w), 1841 (w), 1658 (s), 1627 (s), 1488 (s), 1381 (s), 1312 (s), 1287 (s), 1161 (s), 1123 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 8.06 (1H, app dq, $J = 16$ Hz, $J = 2.4$ Hz), 7.70 (1H, d, $J = 8.0$ Hz), 7.65 (1H, d, $J = 8.0$ Hz), 7.53 (1H, t, $J = 8.0$ Hz), 7.42 (1H, d, $J = 8.0$ Hz), 6.96 (1H, d, $J = 16$ Hz), 3.72 (3H, s), 3.28 (3H, s); ¹³C NMR (CDCl₃): δ 165.9, 138.9, 134.3, 131.9, 129.0, 128.7 (q, $^2J_{CF} = 30$ Hz), 127.9, 126.0 (q, $^3J_{CF} = 5.5$ Hz), 123.9 (q, $^1J_{CF} = 272$ Hz), 120.3, 61.88, 32.42. LRMS (APPI) Calcd for C₁₂H₁₂F₃NO₂ (M + H)⁺: 260.1, Found (M + H)⁺: 260.1.



(E)-3-(Benzo[d][1,3]dioxol-5-yl)-N-methoxy-N-


methylacrylamide (2.190). Prepared according to the general

procedure on page 174 in 57% yield. A white solid. Mp 107-110 °C. $R_f = 0.21$ (SiO₂, 1:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 2974 (m), 2936 (m), 2905 (m), 1841 (w), 1652 (s), 1614 (s), 1501 (s), 1444 (s), 1375 (s), 1255 (s), 1180 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.59 (1H, d, $J = 16$ Hz), 7.03 (1H, s), 6.99 (1H, d, $J = 8.0$ Hz), 6.81 (1H, d, $J = 16$ Hz), 6.75 (1H, d, $J = 8$ Hz), 5.94 (2H, s), 3.70 (3H, s), 3.25 (3H, s); ¹³C NMR (CDCl₃): δ 166.9, 149.0, 148.0, 142.9, 129.4, 124.0, 113.6, 108.3, 106.4, 101.3, 61.72, 32.44. LRMS (ESI⁺) Calcd for C₁₂H₁₃NO₄ (M)⁺: 235.1, Found (M)⁺: 235.6.



(E)-3-(1-Benzyl-1H-indol-3-yl)-N-methoxy-N-methylacrylamide (2.191). The title compound was prepared by

Horner-Wadsworth-Emmons olefination between diethyl(*N*-methoxy-*N*-methylcarbamoylmethyl)phosphonate and 1-benzyl-1H-indole-3-carbaldehyde⁵ in 86% yield.⁶ A white solid. Mp 42-46 °C. R_f = 0.14 (SiO₂, 2:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3106 (m), 3055 (m), 2968 (m), 2936 (m), 1651 (s), 1608 (s), 1526 (m), 1469 (s), 1381 (s), 1167 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.96 (1H, d, J = 16 Hz), 7.90-8.0 (1H, m), 7.40 (1H, s), 7.20-7.34 (6H, m), 7.10 (2H, d, J = 6.4 Hz), 7.05 (1H, d, J = 16 Hz), 5.26 (2H, s), 3.79 (3H, s), 3.31 (3H, s); ¹³C NMR (CDCl₃): δ 168.1, 137.4, 136.4, 136.2, 132.1, 128.7, 127.8, 126.7, 126.2, 122.8, 121.1, 120.4, 113.0, 110.7, 110.3, 61.59, 50.20, 32.51. LRMS (APPI) Calcd for C₂₀H₂₀N₂O₂ (M + H)⁺: 321.2, Found (M + H)⁺: 321.2.



(E)-3-(5-Fluoro-2-methylphenyl)-N-methoxy-N-methylacrylamide (2.192). The title compound was prepared by

Horner-Wadsworth-Emmons olefination between diethyl(*N*-methoxy-*N*-methylcarbamoylmethyl)phosphonate and 5-fluoro-2-methylbenzaldehyde in 78% yield.⁶⁸ A white solid. Mp 68-72 °C. R_f = 0.14 (SiO₂, 3:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3075 (w), 2973 (m), 2936 (m), 1658 (s), 1620 (s), 1589 (m), 1495 (s), 1412 (m), 1381 (s), 1262 (m), 1179 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.88 (1H, dd, J = 16 Hz, J = 1.6 Hz), 7.23 (1H, dd, J = 9.6 Hz, J = 2.4 Hz), 7.10 (1H, dd, J = 8.4 Hz, J = 5.6 Hz),


⁵ Ottoni, O.; Cruz, R.; Alves, R. *Tetrahedron* **1998**, 54, 13915.

⁶ Netz, D. F.; Seidel, J. L. *Tetrahedron Lett.* **1992**, 33, 1957.

6.90 (1H, dt, $J = 8.4$ Hz, $J = 2.4$ Hz), 6.88 (1H, d, $J = 16$ Hz), 3.73 (3H, s), 3.27 (3H, s), 2.35 (3H, s); ^{13}C NMR (CDCl_3): δ 166.4, 161.1 (d, $^1J_{\text{CF}} = 242$ Hz), 139.9, 135.6 (d, $^3J_{\text{CF}} = 7.0$ Hz), 133.1, 131.9 (d, $^3J_{\text{CF}} = 7.8$ Hz), 117.9, 116.2 (d, $^2J_{\text{CF}} = 20$ Hz), 112.4 (d, $^2J_{\text{CF}} = 22$ Hz), 61.84, 32.41. LRMS (APPI) Calcd for $\text{C}_{12}\text{H}_{14}\text{FNO}_2$ ($\text{M} + \text{H}$) $^+$: 224.1, Found ($\text{M} + \text{H}$) $^+$: 224.1.


2. Dialkylidene Ketone Syntheses

Synthesis of **2.62**




To 2.25 g (10.0 mmol) of (*E*)-1-iodoheptene (**2.120**) in 10 mL of THF at $-78\text{ }^\circ\text{C}$ was added 4.2 mL (10 mmol) of a 2.4 M solution of $n\text{-BuLi}$ in hexane dropwise. This solution was allowed to stir for 30 min at $-78\text{ }^\circ\text{C}$ and then transferred dropwise by canula to a solution of 0.965g (5.00 mmol) of **2.65** in 50 mL of THF at $-78\text{ }^\circ\text{C}$. After complete addition (10 min), TLC analysis showed complete consumption of **2.65** after 15 min at $-78\text{ }^\circ\text{C}$, so the reaction was subsequently quenched with saturated aqueous NH_4Cl . The unpurified mixture was transferred to a separatory funnel with 1 M HCl and CH_2Cl_2 . The organic layer was collected after shaking, and the aqueous layer was extracted with CH_2Cl_2 (1x). The combined organics were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. Silica gel chromatography (hexanes/EtOAc) of the

mixture afforded 1.0 g (4.4 mmol, 88%) of (1*E*, 4*E*)-1-phenyldeca-1,4-dien-3-one (**2.62**) as a yellow oil. Spectral data were consistent with the literature.⁷



(1*E*,4*E*)-1-Cyclohexyl-5-phenylpenta-1,4-dien-3-one (2.68).


Prepared in 96% yield according to the procedure used to synthesize **2.62** on page 178. A yellow solid. Mp 42-46 °C. R_f = 0.20 (12:1 hexanes:EtOAc); IR (KBr): 2924 (m), 2847 (m), 1806 (w), 1670 (s), 1627 (s), 1592 (s), 1441 (s), 1332 (s), 1278 (m), 1184 (m) cm^{-1} ; ^1H NMR: δ 7.62 (1H, d, J = 16 Hz), 7.55 (2H, m), 6.97 (1H, d, J = 164 Hz), 6.93 (1H, dd, J = 16 Hz, J = 6.8 Hz), 6.37 (1H, dd, J = 14 Hz, J = 1.2 Hz), 2.19 (1H, m), 1.61-1.88 (5H, m) 1.10-1.40 (5H, m); ^{13}C NMR: δ 189.6, 153.2, 142.8, 134.8, 130.3, 128.9, 128.2, 126.8, 124.8, 40.87, 31.80, 25.90, 25.71. LRMS (ESI+) Calcd for $\text{C}_{17}\text{H}_{10}\text{O}$ ($\text{M} + \text{Na}$)⁺: 263.1, Found ($\text{M} + \text{Na}$)⁺: 263.2.



(1*E*,4*E*)-6,6-Dimethyl-1-phenylhepta-1,4-dien-3-one (2.69).

Prepared in 86% yield according to the procedure used to synthesize **2.62** on page 178. An off-white solid. Mp 60-64 °C. R_f = 0.30 (12:1 hexanes:EtOAc); IR (KBr): 3060 (w), 3017 (w), 2948 (m), 2862 (w), 1950 (w), 1880 (w), 1658 (s), 1588 (s), 1449 (m), 1324 (s), 1208 (s), 1165 (s) cm^{-1} ; ^1H NMR: δ 7.63 (1H, d, J = 16 Hz), 7.56 (2H, m), 7.37 (3H, m), 6.71 (1H, d, J = 16 Hz), 6.32 (1H, d, J = 16 Hz), 1.12 (9H, s); ^{13}C NMR: δ 189.7, 157.9, 142.9, 134.8, 130.3, 128.9, 128.3, 124.9, 124.5, 33.97, 28.72. LRMS (ESI+) Calcd for $\text{C}_{15}\text{H}_{18}\text{O}$ ($\text{M} + \text{Na}$)⁺: 237.1, Found ($\text{M} + \text{Na}$)⁺: 237.1.


⁷ Tsuge, O.; Kanemasa, S.; Nakagawa, N.; Suga, H. *Bull. Chem. Soc. Jpn.* **1987**, 69, 4091.



(1E,4E)-6-(*t*-Butyldimethylsilyloxy)-1-phenylhexa-1,4-dien-3-

one (2.70). Prepared in 60% yield according to the procedure


used to synthesize **2.62** on page 178. A yellow oil. $R_f = 0.20$ (12:1 Hexanes:EtOAc); IR (neat): 3060 (w), 2955 (s), 2932 (s), 2854 (s), 1953 (w), 1806 (w), 1666 (s), 1634 (s), 1592 (s), 1449 (s), 1332 (s), 1254 (s), 1138 (s) cm^{-1} ; ^1H NMR: δ 7.64 (1H, d, $J = 16$ Hz), 7.56 (2H, m), 7.38 (3H, m), 7.02 (1H, dt, $J = 15$ Hz, $J = 3.6$ Hz), 6.95 (1H, d, $J = 16$ Hz), 6.73 (1H, dt, $J = 15$ Hz, $J = 2.0$ Hz), 4.41 (2H, m), 0.94 (9H, s), 0.10 (6H, s); ^{13}C NMR: δ 189.1, 146.1, 143.4, 134.7, 130.4, 128.9, 128.3, 126.5, 125.3, 62.52, 25.86, 18.37, -5.41. LRMS (ESI+) Calcd for $\text{C}_{18}\text{H}_{26}\text{O}_2\text{Si}$ ($\text{M} + \text{H}$) $^+$: 303.2, Found ($\text{M} + \text{H}$) $^+$: 303.2.




(1E,4E)-7-(*t*-Butyldimethylsilyloxy)-1-phenylhepta-1,4-dien-

3-one (2.71). Prepared in 87% yield according to the procedure used to synthesize **2.62** on page 178. A yellow oil. $R_f = 0.19$ (12:1 hexanes:EtOAc); IR (neat): 3029 (w), 2952 (s), 2854 (s), 1950 (w), 1802 (w), 1662 (s), 1631 (s), 1596 (s), 1472 (m), 1336 (m), 1254 (s), 1185 (m) cm^{-1} ; ^1H NMR: δ 7.63 (1H, d, $J = 16$ Hz), 7.55 (2H, m), 7.38 (3H, m), 6.98 (1H, dt, $J = 16$ Hz, $J = 8.6$ Hz), 6.96 (1H, d, $J = 16$ Hz), 3.61 (1H, dt, $J = 16$ Hz, $J = 1.2$ Hz), 3.76 (2H, t, $J = 6.4$ Hz) 3.08 (2H, dq, $J = 7.2$ Hz, $J = 1.6$ Hz), 0.88 (9H, s), 0.050 (3H, s), 0.042 (3H, s); ^{13}C NMR: δ 189.1, 144.7, 143.1, 134.8, 130.8, 130.3, 128.9, 128.2, 124.6, 61.60, 36.15, 25.86, 18.28, -5.33. LRMS (ESI+) Calcd for $\text{C}_{19}\text{H}_{28}\text{O}_2\text{Si}$ ($\text{M} + \text{Na}$) $^+$: 339.2, Found ($\text{M} + \text{Na}$) $^+$: 339.2.

Synthesis of **2.67**



To 1.05 g (8.68 mmol) of (*E*)-1-bromopropene in 16 mL of THF at $-78\text{ }^{\circ}\text{C}$ was added 10.8 mL (17.2 mmol) of a 1.6 M solution of *t*-BuLi in pentane dropwise over a 15 min period. This solution was allowed to stir at this temperature for 30 min and then transferred at ~ 1 drop/s by canula to a solution of 0.832 g (4.35 mmol) of **2.65** in 33 mL of THF at $-78\text{ }^{\circ}\text{C}$ (addition of the vinylolithium at faster rates gave lower yields). After complete addition, TLC analysis showed complete consumption of the starting material after 15 min at $-78\text{ }^{\circ}\text{C}$, so the reaction was subsequently quenched with saturated aqueous NH_4Cl . The unpurified mixture was transferred to a separatory funnel with 1 M HCl and Et_2O . The organic layer was collected after shaking, and the aqueous layer was extracted with Et_2O (1x). The combined organics were washed with H_2O , then brine, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. Silica gel chromatography (hexanes/ EtOAc) of the mixture afforded 0.571 g (3.32 mmol, 76%) of (1*E*, 4*E*)-1-phenylhexa-1,4-dien-3-one (**2.67**) as a yellow oil. Spectral data were consistent with the literature.⁸




(1*E*,4*E*)-4-Methyl-1-phenylhexa-1,4-dien-3-one (2.73). Prepared in 53% yield according to the procedure used to synthesize **2.67** on page

181. A yellow oil. $R_f = 0.19$ (14:1 hexanes: EtOAc); IR (neat): 3060 (m), 2920 (m), 1953


⁸ Kubota, K.; Isaka, M.; Nakamura, E. *Heterocycles* **1996**, 42, 565.

(w), 1887 (w), 1654 (s), 1600 (s), 1449 (s), 1328 (s), 1300 (s), 1223 (s) cm^{-1} ; ^1H NMR: δ 7.59 (1H, d, $J = 16$ Hz), 7.55 (2H, m), 7.37 (3H, m), 7.28 (1H, d, $J = 16$ Hz), 6.82 (1H, q, $J = 6.0$ Hz), 1.89 (3H, d, $J = 6.0$ Hz), 1.88 (3H, s); ^{13}C NMR: δ 191.7, 142.5, 139.3, 137.4, 135.2, 129.9, 128.8, 128.1, 121.7, 14.80, 11.55. LRMS (ESI+) Calcd for $\text{C}_{13}\text{H}_{14}\text{O}$ ($\text{M} + \text{Na}$) $^+$: 209.1, Found ($\text{M} + \text{Na}$) $^+$: 209.1.



***trans*-1-Phenyl-5-methyl-1,4-hexadien-3-one (2.72).** Dialkylidene ketone **2.72** was prepared in 53% yield according to the procedure used to synthesize **2.67** on page 181. This compound has been previously reported.⁹

Synthesis of Ketone 2.122




To 0.426 g (1.90 mmol) of (*E*)-1-iodoheptene in 2 mL of THF at -78 $^{\circ}\text{C}$ was added 0.76 mL (1.9 mmol) of a 2.5 M solution of *n*-BuLi in hexane dropwise over a 5 min period. This solution was allowed to stir at this temperature for 30 min and then transferred by canula to a solution of 0.215 g (0.951 mmol) of **2.187** in 10 mL of THF at -78 $^{\circ}\text{C}$. After complete addition (5 min), TLC analysis showed complete consumption of the starting material after 15 min at -78 $^{\circ}\text{C}$, so the reaction was subsequently quenched with saturated aqueous NH_4Cl . The unpurified mixture was transferred to a separatory

⁹ Inaba, S. -I.; Rieke, R. D. *J. Org. Chem.* **1985**, 50, 1373.

funnel with 1 M HCl and Et₂O. The organic layer was collected after shaking, and the aqueous layer was extracted with Et₂O (1x). The combined organics were washed with H₂O, then brine, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure. Silica gel chromatography (hexanes/EtOAc) of the mixture afforded 0.203 g (0.773 mmol, 81%) of (1*E*, 4*E*)-1-(3-chlorophenyl)deca-1,4-dien-3-one (**2.122**) as a light-yellow solid.


(1*E*,4*E*)-1-(3-Chlorophenyl)deca-1,4-dien-3-one, (2.122). Mp 38-40 °C. *R_f* = 0.19 (SiO₂, 15:1 Hexanes:EtOAc); IR (CH₂Cl₂ solution): 3031, (w), 2955 (s), 2924 (s), 2854 (s), 2817 (w), 1948 (w), 1665 (s), 1602 (s), 1463 (m), 1419 (m), 1293 (m), 1199 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.52 (1H, d, *J* = 16 Hz), 7.52 (1H, m), 7.39 (1H, dt, *J* = 6.8 Hz, *J* = 2.0 Hz), 7.25-7.34 (2H, m), 6.99 (1H, dd, *J* = 16 Hz, *J* = 7.2 Hz), 6.94 (1H, d, *J* = 16 Hz), 6.74 (1H, dt, *J* = 16 Hz, *J* = 1.6 Hz), 2.25 (2H, q, *J* = 7.2 Hz), 1.48 (2H, p, *J* = 7.2 Hz), 1.24-1.38 (4H, m) 0.871 (3H, t, *J* = 7.2 Hz); ¹³C NMR (CDCl₃): δ 188.7, 148.8, 141.0, 136.6, 134.8, 130.03, 130.00, 129.3, 127.7, 126.5, 125.7, 32.700, 31.38, 27.81, 22.44, 13.96. LRMS (APPI) Calcd for C₁₆H₁₉ClO (M + H)⁺: 263.1, Found (M + H)⁺: 263.1.



(1*E*,4*E*)-1-(4-Methoxyphenyl)deca-1,4-dien-3-one (2.123). Prepared in 90% yield according to the procedure


used to synthesize **2.122** on page 182. An oil. *R_f* = 0.11 (SiO₂, 10:1 hexanes:EtOAc); IR (neat): 3006 (w), 2955 (m), 2930 (m), 2861 (m), 1659 (s), 1627 (s), 1590 (s), 1508 (s),

1420 (m), 1306 (m), 1256 (s), 1168 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.56 (1H, d, $J = 16$ Hz), 7.46 (2H, d, $J = 8.8$ Hz), 6.93 (1H, dt, $J = 16$ Hz, $J = 7.2$ Hz), 6.84 (2H, d, $J = 8.8$ Hz), 6.80 (1H, d, $J = 16$ Hz), 6.36 (1H, d, $J = 16$ Hz), 3.76 (3H, s), 2.20 (2H, q, $J = 7.2$ Hz), 1.44 (2H, p, $J = 7.2$ Hz), 1.20-1.36 (4H, m), 0.847 (3H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 188.9, 161.3, 147.6, 142.5, 129.8, 129.1, 127.3, 122.5, 114.2, 55.17, 32.52, 31.27, 27.78, 22.34, 13.87. LRMS (APPI) Calcd for $\text{C}_{17}\text{H}_{22}\text{O}_2$ ($\text{M} + \text{H}$) $^+$: 259.2, Found ($\text{M} + \text{H}$) $^+$: 259.2.



(1E,4E)-1-o-Tolyldeca-1,4-dien-3-one (2.124). Prepared in 89% yield according to the procedure used to synthesize **2.122**


on page 182. An oil. $R_f = 0.25$ (SiO_2 , 15:1 hexanes:EtOAc); IR (neat): 3025 (w), 2955 (s), 2930 (s), 2854 (m), 1658 (s), 1627 (s), 1596 (s), 1457 (m), 1319 (m), 1099 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.92 (1H, d, $J = 16$ Hz), 7.58 (1H, d, $J = 7.6$ Hz), 7.10-7.30 (3H, m), 6.99 (1H, dt, $J = 16$ Hz, $J = 7.2$ Hz), 6.89 (1H, d, $J = 16$ Hz), 6.39 (1H, dt, $J = 16$ Hz, $J = 1.6$ Hz), 2.41 (3H, s), 2.25 (2H, q, 7.2 Hz), 1.48 (2H, p, $J = 7.6$ Hz), 1.20-1.40 (4H, m), 0.88 (3H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 189.0, 148.2, 140.3, 137.9, 133.6, 130.6, 129.9, 129.5, 126.1, 125.5, 32.61, 31.31, 27.78, 22.38, 19.71, 13.90. LRMS (ESI $^+$) Calcd for $\text{C}_{17}\text{H}_{22}\text{O}$ (M) $^+$: 242.2, Found (M) $^+$: 242.7.



(1E,4E)-1-[2-(Trifluoromethyl)phenyl]deca-1,4-dien-3-one (2.125). Prepared in 79% yield according to the procedure used

to synthesize **2.122** on page 182. A yellow oil. $R_f = 0.22$ (SiO_2 , 16:1 hexanes:EtOAc);


IR (neat): 2955 (s), 2930 (s), 2867 (m), 1659 (s), 1633 (s), 1602 (s), 1489 (m), 1313 (s), 1162 (s), 1124 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.93 (1H, d, $J = 16$ Hz), 7.71 (1H, d, $J = 8.0$ Hz), 7.65 (1H, d, $J = 8.0$ Hz), 7.53 (1H, t, $J = 8.0$ Hz), 7.43 (1H, t, $J = 8.0$ Hz), 6.98 (1H, dt, $J = 16$ Hz, $J = 7.2$ Hz), 6.86 (1H, d, $J = 16$ Hz), 6.39 (1H, dt, $J = 16$ Hz, $J = 1.6$ Hz), 2.24 (2H, q, $J = 6.8$ Hz), 1.47 (2H, p, $J = 7.2$ Hz), 1.22-1.36 (4H, m), 0.86 (3H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 188.9, 149.2, 138.1, 133.8, 132.0, 129.4, 128.9, 128.85 (q, $^2J_{\text{CF}} = 30$ Hz), 128.6, 127.7, 126.0 (q, $^3J_{\text{CF}} = 5.4$ Hz), 123.9 (q, $^1J_{\text{CF}} = 272$ Hz), 32.68, 31.31, 27.76, 22.39, 13.88. LRMS (ESI+) Calcd for $\text{C}_{17}\text{H}_{19}\text{F}_3\text{O}$ (M^+): 296.1, Found (M^+): 296.7.



(1E,4E)-1-[4-(Trifluoromethyl)phenyl]deca-1,4-dien-3-

one (2.126). Prepared in 66% yield according to the


procedure used to synthesize **2.122** on page 182. A yellow oil. $R_f = 0.23$ (SiO_2 , 16:1 hexanes:EtOAc); IR (neat): 3043 (w), 2961 (s), 2930 (s), 2861 (m), 1923 (w), 1665 (s), 1633 (s), 1469 (m), 1419 (m), 1319 (s), 1167 (s), 1135 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.65 (1H, d, $J = 8.7$ Hz), 7.63 (1H, d, $J = 8.7$ Hz), 7.62 (1H, d, $J = 16$ Hz), 7.02 (1H, d, $J = 16$ Hz), 7.02 (1H, dt, $J = 16$ Hz, $J = 6.6$ Hz), 6.41 (1H, d, $J = 16$ Hz), 2.28 (2H, q, $J = 6.6$ Hz), 1.50 (2H, p, $J = 7.5$ Hz), 1.24-1.40 (4H, m), 0.89 (3H, t, $J = 6.6$ Hz); ^{13}C NMR (CDCl_3): δ 188.7, 149.2, 140.8, 138.2, 131.6 (q, $^2J_{\text{CF}} = 32$ Hz), 129.2, 128.3, 126.8, 125.8 (q, $^3J_{\text{CF}} = 3.1$ Hz), 123.8 (q, $^1J_{\text{CF}} = 270$ Hz), 32.76, 31.42, 27.85, 22.48, 13.93. LRMS (ESI+) Calcd for $\text{C}_{17}\text{H}_{19}\text{F}_3\text{O}$ (M^+): 296.1, Found (M^+): 296.7.



(1E,4E)-1-(5-Fluoro-2-methylphenyl)deca-1,4-dien-3-one

(2.127). Prepared in 82% yield according to the procedure

used to synthesize **2.122** on page 182. A yellow oil. R_f = 0.11 (SiO₂, 20:1 hexanes:EtOAc); IR (neat): 2955 (m), 2930 (s), 2861 (m), 1659 (s), 1627 (s), 1489 (s), 1338 (m), 1237 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.83 (1H, dd, J = 16 Hz, J = 1.6 Hz), 7.26 (1H, dd, J = 9.6 Hz, J = 2.4 Hz), 7.12 (1H, dd, J = 8.4 Hz, J = 5.6 Hz), 6.99 (1H, dt, J = 16 Hz, J = 6.8 Hz), 6.95 (1H, dt, J = 8.4 Hz, J = 2.4 Hz), 6.86 (1H, d, J = 16 Hz), 6.36 (1H, dt, J = 16 Hz, J = 1.6 Hz), 2.37 (3H, s), 2.26 (2H, q, J = 6.8 Hz), 1.49 (2H, p, J = 7.2 Hz), 1.21-1.37 (4H, m), 0.88 (3H, t, J = 6.8 Hz); ¹³C NMR (CDCl₃): δ 188.8, 161.2 (d, $^1J_{CF}$ = 242 Hz), 148.8, 139.2, 135.4 (d, $^3J_{CF}$ = 7.8 Hz), 133.6, 132.0 (d, $^3J_{CF}$ = 7.8 Hz), 129.6, 126.3, 116.7 (d, $^2J_{CF}$ = 21 Hz), 112.4 (d, $^2J_{CF}$ = 22 Hz), 32.72, 31.38, 27.83, 22.45, 19.03, 13.96; ¹⁹F NMR (CDCl₃): δ 93.28 (m). LRMS (APPI) Calcd for C₁₇H₂₁FO (M + H)⁺: 261.2, Found (M + H)⁺: 261.2.




(1E,4E)-1-(Benzo[d][1,3]dioxol-5-yl)deca-1,4-dien-3-one

(2.128). Prepared in 87% yield according to the procedure


used to synthesize **2.122** on page 182. An off-white solid. Mp = 40-44 °C. R_f = 0.23 (SiO₂, 9:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 2961 (s), 2930 (s), 2861 (s), 1658 (s), 1626 (s), 1589 (s), 1488 (s), 1444 (s), 1362 (m), 1255 (s), 1205 (m), 1098 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.51 (1H, d, J = 16 Hz), 7.04 (1H, s), 7.01 (1H, d, J = 8.4 Hz), 6.94 (1H, dt, J = 16 Hz, J = 7.2 Hz), 6.77 (1H, d, J = 16 Hz), 6.77 (1H, d, J = 8.4 Hz), 6.36 (1H, dt, J = 16 Hz, J = 1.6 Hz), 5.96 (2H, s), 2.22 (2H, q, J = 7.2 Hz), 1.46 (2H, p, J = 7.2

Hz), 1.20-1.37 (4H, m) 0.86 (3H, t, $J = 6.4$ Hz); ^{13}C NMR (CDCl_3): δ 188.9, 149.6, 148.2, 147.9, 142.6, 129.25, 129.16, 124.8, 122.8, 108.5, 106.4, 101.5, 32.61, 31.35, 27.84, 22.41, 13.94. LRMS (APPI) Calcd for $\text{C}_{17}\text{H}_{20}\text{O}_3$ ($\text{M} + \text{H}$) $^+$: 273.1, Found ($\text{M} + \text{H}$) $^+$: 273.1.



(1E,4E)-1-(1-benzyl-1H-indol-3-yl)deca-1,4-dien-3-one (2.129). Prepared in 88% yield according to the procedure


used to synthesize **2.122** on page 182. A yellow solid. Mp 84-88 °C. $R_f = 0.21$ (SiO_2 , 5:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3106 (m), 3037 (w), 2961 (m), 2930 (m), 2854 (m), 1658 (m), 1627 (s), 1576 (s), 1526 (s), 1463 (m), 1387 (s), 1350 (m), 1281 (m), 1173 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.94-8.00 (1H, m), 7.91 (1H, d, $J = 16$ Hz), 7.44 (1H, s), 7.16-7.26 (6H, m), 7.13 (2H, dd, $J = 7.6$ Hz, $J = 2$ Hz), 6.99 (1H, dt, $J = 16$ Hz, $J = 8.0$ Hz), 6.99 (1H, d, $J = 16$ Hz), 6.44 (1H, dt, $J = 16$ Hz, $J = 1.6$ Hz), 5.30 (2H, s), 2.26 (2H, q, $J = 8.0$ Hz), 1.51 (2H, p, $J = 7.2$ Hz) 1.23-1.41 (4H, m), 0.90 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 189.2, 146.7, 137.7, 136.5, 136.0, 133.1, 133.0, 129.5, 128.9, 128.0, 126.9, 126.3, 123.1, 121.5, 120.7, 113.2, 110.5, 50.46, 32.65, 31.44, 27.98, 22.49, 14.01. LRMS (APPI) Calcd for $\text{C}_{25}\text{H}_{27}\text{NO}$ ($\text{M} + \text{H}$) $^+$: 358.2, Found ($\text{M} + \text{H}$) $^+$: 358.2.




(1E,4E)-1-(Furan-2-yl)deca-1,4-dien-3-one (2.130). Prepared in 86% yield according to the procedure used to synthesize **2.122**

on page 182. A yellow oil. $R_f = 0.15$ (SiO_2 , 15:1 hexanes:EtOAc); IR (neat): 3124 (w), 3037 (w), 2955 (s), 2930 (s), 2861 (m), 1658 (s), 1633 (s), 1595 (s), 1482 (m), 1306 (m),

1217 (m), 1098 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.44 (1H, s), 7.36 (1H, d, $J = 16$ Hz), 6.93 (1H, dt, $J = 16$ Hz, $J = 6.8$ Hz), 6.85 (1H, d, $J = 16$ Hz), 6.60 (1H, d, $J = 3.2$ Hz), 6.42 (1H, m), 6.29 (1H, d, $J = 16$ Hz), 2.20 (2H, q, $J = 6.8$ Hz), 1.44 (2H, p, $J = 6.8$ Hz), 1.17-1.36 (4H, m), 0.85 (3H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 188.6, 151.4, 148.0, 144.6, 129.7, 128.9, 121.8, 115.4, 112.4, 32.60, 31.31, 27.79, 22.39, 13.90. LRMS (APPI) Calcd for $\text{C}_{14}\text{H}_{18}\text{O}_2$ ($\text{M} + \text{H}$) $^+$: 219.1, Found ($\text{M} + \text{H}$) $^+$: 219.1.

 **(1E,4E)-1-Phenylhepta-1,4-dien-3-one (2.193).** Prepared in 90% yield according to the procedure used to synthesize **2.62** on page 178. A yellow oil. $R_f = 0.29$ (SiO_2 , 10:1 hexanes:EtOAc); IR (neat): 3031 (w), 2968 (m), 2936 (m), 2879 (w), 1659 (s), 1633 (s), 1602 (s), 1450 (m), 1349 (m), 1199 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.63 (1H, d, $J = 16$ Hz), 7.56 (2H, m), 7.38 (3H, m), 7.05 (1H, dt, $J = 16$ Hz, $J = 6.4$ Hz), 6.97 (1H, d, $J = 16$ Hz), 6.42 (1H, d, $J = 16$ Hz), 2.30 (2H, p, $J = 7.2$ Hz), 1.11 (3H, t, $J = 7.2$ Hz) 1.10-1.40 (5H, m); ^{13}C NMR (CDCl_3): δ 189.4, 149.7, 143.0, 134.7, 130.2, 128.8, 128.23, 128.19, 124.6, 25.77, 12.30. LRMS (APPI) Calcd for $\text{C}_{13}\text{H}_{14}\text{O}$ ($\text{M} + \text{H}$) $^+$: 187.1, Found ($\text{M} + \text{H}$) $^+$: 187.1.

Synthesis of 2.194




To a solution of 0.763 g (2.41 mmol) of **2.71** in 12 mL of 3:1 THF:H₂O was added 0.458 g (2.41 mmol) of *p*-toluenesulfonic acid monohydrate under a N₂

atmosphere. This mixture was allowed to stir at room temperature and monitored by TLC. The starting material was consumed after 1h as evident by TLC analysis, and H₂O was then added, followed by extraction with Et₂O. The combined organics were dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. Silica gel chromatography (hexanes/EtOAc) of the mixture afforded 0.479 g (2.37 mmol, 98%) of **2.194** as a yellow oil.

(1E,4E)-7-Hydroxy-1-phenylhepta-1,4-dien-3-one (2.194). $R_f = 0.30$ (SiO₂, 1:2 hexanes:EtOAc); IR (neat): 3415 (s, br), 3058 (w), 2939 (m), 2878 (m), 1958 (w), 1659 (s), 1628 (s), 1598 (s), 1494 (m), 1449 (m), 1333 (s), 1308 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.61 (1H, d, $J = 16$ Hz), 7.48-7.54 (2H, m), 7.31-7.37 (3H, m), 6.98 (1H, dt, $J = 16$ Hz, $J = 7.0$ Hz), 6.93 (1H, d, $J = 16$ Hz), 6.50 (1H, dt, $J = 16$ Hz, $J = 1.6$ Hz), 3.78 (2H, t, $J = 6.2$ Hz), 2.62-2.92 (1H, br s), 2.51 (2H, dq, $J = 7.0$ Hz, $J = 1.6$ Hz); ¹³C NMR (CDCl₃): δ 189.1, 144.4, 143.4, 134.5, 130.9, 130.4, 128.8, 128.2, 124.5, 60.84, 35.87. LRMS (ESI+) Calcd for C₁₃H₁₄O₂ (M + H)⁺: 203.1, Found (M + H)⁺: 203.1.

Synthesis of 2.96




To 0.200 g (0.968 mmol) of **2.194** in 1.9 mL of CH₂Cl₂ was added 0.386 g (1.26 mmol) of *o*-diphenylphosphinobenzoic acid, 0.260 g (1.26 mmol) of N,N'-dicyclohexylcarbodiimide (DCC), and 11.9 mg (0.0968 mmol) of 4-

(dimethylamino)pyridine (DMAP) sequentially under N₂. This mixture was then allowed to stir at ambient temperature for 2h, and the mixture was subsequently filtered through celite using CH₂Cl₂. Volatile material was removed under reduced pressure, and the resultant oil purified by silica gel chromatography (hexanes/EtOAc) to afford 0.348 g (0.709 mmol, 73%) of **2.96** as a light-yellow solid.

(3E,6E)-5-Oxo-7-phenylhepta-3,6-dienyl 2-(diphenylphosphino)benzoate (2.96). Mp 140-145 °C. R_f = 0.20 (SiO₂, 4:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3411 (br, m), 3058 (m), 2957 (m), 1967 (w), 1891 (w), 1717 (s), 1662 (s), 1628 (s), 1601 (s), 1433 (s), 1339 (m), 1275 (s), 1189 (s), 1122 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 8.00-8.07 (1H, m), 7.63 (1H, d, *J* = 16 Hz), 7.51-7.59 (2H, m), 7.20-7.42 (16H, m), 6.96 (1H, d, *J* = 16 Hz), 6.89 (1H, dt, *J* = 16 Hz, *J* = 6.8 Hz), 6.46 (1H, d, *J* = 16 Hz), 4.32 (2H, t, *J* = 6.4 Hz), 2.55 (2H, q, *J* = 6.4 Hz); ¹H NMR (C₆D₆): δ 8.04-8.10 (1H, m), 7.70 (1H, d, *J* = 16 Hz), 7.32-7.46 (4H, m), 6.85-7.25 (14H, m), 6.81 (1H, d, *J* = 16 Hz), 6.73 (1H, dt, *J* = 16 Hz, *J* = 6.8 Hz), 6.16 (1H, d, *J* = 16 Hz), 3.94 (2H, t, *J* = 6.4 Hz), 2.00 (2H, q, *J* = 6.4 Hz); ¹³C NMR (CDCl₃): δ 188.7, 166.6, 143.3, 142.6, 140.3, 140.1, 137.8, 137.7, 134.7, 134.3, 133.9, 133.7, 132.0, 131.0, 130.6, 130.4, 128.8, 128.6, 128.5, 128.4, 128.3, 124.7, 63.11, 31.74. ¹³C NMR (C₆D₆): δ 187.9, 167.0, 143.1, 142.5, 139.3, 139.2, 138.8, 135.2, 134.8, 134.6, 132.4, 131.9, 131.3, 130.5, 129.3, 129.1, 128.9, 128.8, 125.8, 63.67, 32.22. ³¹P NMR (C₆D₆): δ -2.86. LRMS (ESI+) Calcd for C₃₂H₂₇O₃P (M + H)⁺: 491.2, Found (M + H)⁺: 491.1.


Synthesis of **2.99**



To a solution of 96.7 mg (0.478 mmol) of **2.194** in 1.9 mL of CH₂Cl₂ at 0 °C was added 0.10 mL of pyridine, 12 mg of DMAP, and 0.14 mL of benzoyl chloride, sequentially. Reaction progress was monitored by TLC, and after 1h at 0 °C, 0.05 mL of pyridine and 0.07 mL of benzoyl chloride were added. After an additional 30 min at 0°C, starting material had been consumed as evident by TLC analysis. Saturated aqueous NaHCO₃ was added, and the aqueous layer extracted with Et₂O (1x). The organic layer was washed with 1M HCl (2x) and dried with anhydrous Na₂SO₄. After removal of volatile material under reduced pressure, the mixture was purified by silica gel chromatography (hexanes/EtOAc) to afford 124 mg (0.405 mmol, 85%) of **2.99** as an off-white solid.

(3E,6E)-5-Oxo-7-phenylhepta-3,6-dienyl benzoate (2.99). Mp 60-66 °C. *R_f* = 0.13 (SiO₂, 6:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3054 (m), 3025 (m), 2919 (m), 2848 (m), 1720 (s), 1661 (s), 1602 (s), 1450 (m), 1273 (s), 1188 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.96-8.06 (1H, m), 7.63 (1H, d, *J* = 16 Hz), 7.49-7.59 (3H, m), 7.33-7.47 (5H, m), 7.02 (1H, dt, *J* = 16 Hz, *J* = 6.8 Hz), 6.94 (1H, d, *J* = 16 Hz), 6.57 (1H, d, *J* = 16 Hz), 4.48 (2H, t, *J* = 6.4 Hz), 2.75 (2H, q, *J* = 6.4 Hz); ¹³C NMR (CDCl₃): δ 188.7, 166.3, 143.5, 142.6, 134.6, 133.0, 131.0, 130.4, 129.9, 129.5, 128.9, 128.35, 128.26, 124.8, 62.84, 32.03. LRMS (ES⁺) Calcd for C₂₀H₁₈O₃ (M + H)⁺: 307.1, Found (M + H)⁺: 307.1.

Synthesis of **2.156**




A pressure tube with a magnetic stir-bar was charged with 62.7 mg (0.0892 mmol) of *trans*-bis(triphenylphosphino)palladium dichloride, 4.5 mL of anhydrous THF, 1.00 g (4.46 mmol) of colorless (*E*)-1-iodoheptene (**2.120**), and 1.57 mL (1.70 g, 5.35 mmol) of tri-*n*-butyl(vinyl)tin (**2.195**). The tube was then charged with 50 psig of carbon monoxide gas, and after stirring for 1 min, the pressure was released. This was repeated two more times, and the tube was subsequently charged with 50 psig of CO and immersed in an oil bath at 40 °C and heated overnight. After cooling the reaction to room temperature, the reaction was vented and concentrated under reduced pressure. The residue was filtered through neutral alumina using 100 mL of 9:1 hexanes:EtOAc and then concentrated under reduced pressure. To this mixture was added 65 mL of Et₂O followed by 65 mL of 50% saturated aqueous KF, and the resultant biphasic mixture was allowed to stir rapidly for 2 h. This mixture was then filtered through celite, and the organic layer was collected, dried with anhydrous Na₂SO₄, and concentrated under reduced pressure. Purification of the material using silica gel chromatography (hexanes/EtOAc) afforded 462 mg (3.03 mmol, 68%) of **2.156**.

(*E*)-Deca-1,4-dien-3-one (2.156). A yellow oil. $R_f = 0.23$ (SiO₂, 25:1 hexanes:EtOAc); IR (neat): 3021 (w), 2962 (s), 2932 (s), 2865 (s), 1662 (s), 1632 (s), 1611 (s), 1468 (m), 1396 (s), 1341 (w), 1282 (w), 1105 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 6.88 (1H, dt, $J = 16$

Hz, $J = 6.8$ Hz), 6.55 (1H, dd, $J = 17$ Hz, $J = 10$ Hz), 6.29 (1H, dt, $J = 16$ Hz, $J = 1.6$ Hz), 6.21 (1H, dd, $J = 17$ Hz, $J = 1.6$ Hz), 5.74 (1H, dd, $J = 10$ Hz, $J = 1.6$ Hz), 2.18 (2H, q, $J = 7.2$ Hz), 1.42 (2H, p, $J = 7.2$ Hz), 1.18-1.34 (4H, m), 0.83 (3H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 189.5, 148.9, 134.8, 128.04, 127.98, 32.58, 31.29, 27.73, 22.36, 13.87. LRMS (ESI⁺) Calcd for $\text{C}_{10}\text{H}_{16}\text{O}$ ($\text{M} + \text{H}$)⁺: 153.1, Found ($\text{M} + \text{H}$)⁺: 153.1.

Synthesis of 2.165




A pressure tube with a magnetic stir-bar was charged with 70.2 mg (0.100 mmol) of *trans*-bis(triphenylphosphino)palladium dichloride, 5.0 mL of anhydrous THF, 1.12 g (5.00 mmol) of colorless **2.196**,¹⁰ and 1.75 mL (1.90 g, 6.00 mmol) of tri-*n*-butyl(vinyl)tin (**2.195**). The tube was then charged with 50 psig of carbon monoxide gas, and after stirring for 1 min, the pressure was released. This was repeated two more times, and the tube was subsequently charged with 50 psig of CO and immersed in an oil bath at 50 °C. After ~1 h the pressure had dropped to 30 psig, and the system was repressurized to 50 psig. After an additional 2 h at 50 °C, the pressure again had dropped to 30 psig, so the system was repressurized to 45 psig and heated for an additional 2 h. The reaction was then cooled to room temperature, vented, and concentrated under reduced pressure. The residue was filtered through neutral alumina using 100 mL of 9:1 hexanes:EtOAc and concentrated under reduced pressure. To this mixture was added 100 mL of Et_2O

¹⁰ (a) Morill, C.; Beutner, G. L.; Grubbs, R. H. *J. Org. Chem.* **2006**, *71*, 7813. (b) Negishi, E.; Van Horn, D. E.; Yoshida, T. *J. Am. Chem. Soc.* **1985**, *107*, 6639.

followed by 100 mL of 50% saturated aqueous KF, and the resultant biphasic mixture was allowed to stir rapidly for 2 h. This mixture was then filtered through celite, and the organic layer was collected, dried with anhydrous Na₂SO₄, and concentrated under reduced pressure. Purification of the material using silica gel chromatography (hexanes/EtOAc) afforded 493 mg (3.24 mmol, 65%) of **2.163**.

(E)-5-Methylnona-1,4-dien-3-one (2.163). A yellow oil. R_f = 0.18 (SiO₂, 30:1 hexanes:EtOAc); IR (neat): 3095 (w), 3018 (w), 2954 (s), 2930 (s), 2872 (m), 2860 (m), 1677 (s), 1662 (s), 1625 (s), 1598 (s), 1458 (m), 1400 (s), 1232 (m), 1132 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 6.38 (1H, dd, J = 18 Hz, J = 11 Hz), 6.23 (1H, s), 6.18 (1H, dd, J = 18 Hz, J = 1.2 Hz), 5.71 (1H, dd, J = 11 Hz, J = 1.2 Hz), 2.15 (2H, t, J = 7.6 Hz), 2.13 (3H, s), 1.45 (2H, p, J = 7.6 Hz), 1.30 (2H, h, J = 7.6 Hz), 0.89 (3H, t, J = 7.6 Hz); ¹³C NMR (CDCl₃): δ 190.4, 160.9, 138.3, 126.8, 121.3, 41.18, 29.72, 22.39, 19.65, 13.90. LRMS (ESI+) Calcd for C₁₀H₁₆O (M + H)⁺: 153.1, Found (M + H)⁺: 153.1.

Synthesis of 2.169



To 0.383 g (1.71 mmol) of **2.196** in 1.8 mL of THF at $-78\text{ }^{\circ}\text{C}$ was added 2.6 mL (3.4 mmol) of a 1.3 M solution of $t\text{-BuLi}$ in pentane dropwise over 5 min. This solution was allowed to stir at this temperature for 30 min and then transferred by canula to a solution of 0.273 g (1.43 mmol) of **2.65** (dried by azeotropic removal of water with

benzene) in 14 mL of THF at -78°C . After complete addition (10 min), the reaction was allowed to stir for 15 min at -78°C . To the reaction was then added 0.75 mL of MeOH, the dry-ice acetone bath was removed, and water was added. After reaching room temperature, the mixture was transferred to a separatory funnel with water and Et_2O . The organic layer was collected after shaking, and the aqueous layer was extracted with Et_2O (1x). The combined organics were sequentially washed with saturated aqueous NaHCO_3 , saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$, and again with saturated aqueous NaHCO_3 . The organic layer was finally dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. Silica gel chromatography (1% TEA in hexanes/ EtOAc) of the mixture afforded 0.232 g (1.02 mmol, 71%) of **2.169** as a yellow oil.


(1E,4E)-5-Methyl-1-phenylnona-1,4-dien-3-one (2.169). $R_f = 0.23$ (SiO_2 , 1% TEA in 19:1 Hexanes: EtOAc); IR (neat): 3028 (w), 2957 (m), 2930 (m), 2872 (m), 1674 (s), 1650 (s), 1622 (s), 1576 (m), 1494 (m), 1451 (m), 1385 (w), 1327 (m), 1332 (m), 1202 (m), 1168 (w) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.55 (1H, d, $J = 16$ Hz), 7.49-7.56 (2H, m), 7.31-7.40 (3H, m), 6.78 (1H, d, $J = 16$ Hz), 6.30 (1H, q, $J = 1.2$ Hz), 2.14-2.24 (5H, m), 1.44-1.54 (2H, m), 1.34 (2H, h, $J = 7.6$ Hz), 0.92 (3H, t, $J = 7.6$ Hz); ^{13}C NMR (CDCl_3): δ 190.1, 160.2, 141.7, 134.9, 130.0, 128.8, 128.3, 128.1, 122.9, 41.17, 29.77, 22.41, 19.66, 13.94. LRMS (ESI+) Calcd for $\text{C}_{16}\text{H}_{20}\text{O}$ ($\text{M} + \text{H}$) $^{+}$: 229.2, Found ($\text{M} + \text{H}$) $^{+}$: 229.1.

3. Chiral Ligand Syntheses

TADDOL syntheses:


The requisite TADDOLs were prepared via Grignard addition to the acetonide protected methyl ester of enantiomerically pure tartaric acid using the known route.⁴⁷ An example is given below for the preparation of 3,5-di-*t*-butylphenylTADDOL (**2.199**).

Synthesis of Diol **2.199**



To a flame-dried 100 mL 2-neck round-bottom flask with magnetic stir-bar and a reflux condenser was added 0.487 g (20.3 mmol) of freshly cracked Mg turnings. The Mg was then flame dried followed by the addition of 1 crystal of I₂. An addition funnel was fitted, and 5.02 g (18.6 mmol) of **2.198** in 19 mL of THF was added dropwise. After complete addition, the reaction had reached reflux, and an oil bath was applied to keep the reaction at a gentle reflux for 2 h. The resultant mixture was cooled to 0 °C, and 0.737 g (3.38 mmol) of **2.197** in 6 mL of THF was added. After complete addition (5 min), the mixture was heated to reflux overnight, and subsequently, quenched at 0 °C by the addition of 15 mL of saturated aqueous NH₄Cl, followed by addition of 15 mL of water. This mixture was then allowed to stir at room temperature for ~1 h or until all the

excess Mg metal had been consumed. The mixture was transferred to a separatory funnel and extracted with EtOAc (3x). The combined organic layers were dried with anhydrous Na₂SO₄, and volatile material was removed under reduced pressure. Silica gel chromatography of the mixture afforded 2.65 g (2.89 mmol, 86%) of **2.199** as an off-white solid.



[(4*R*,5*R*)-2,2-dimethyl-1,3-dioxolane-4,5-diyl]bis[bis(3,5-dimethoxyphenyl)methanol] (2.200**). Prepared in 78% yield**

according to the procedure used to synthesize **2.199** on page 196.


Note that 7.0 equiv of 1-bromo-3,5-dimethoxybenzene and 7.0 equiv of Mg metal was used. A white solid. $R_f = 0.11$ (SiO₂, 2:1

hexanes:EtOAc); IR (CH₂Cl₂ solution): 3329 (s, br), 2991 (s), 2940 (s), 2835 (s), 1599 (s), 1459 (s), 1421 (s), 1375 (m), 1345 (s), 1307 (s), 1202 (s), 1160 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 6.73 (4H, s), 6.50 (4H, s), 6.35 (2H, s), 6.26 (2H, s), 4.50 (2H, s), 4.49 (2H, s), 3.70 (12H, s), 3.63 (12H, s), 1.15 (6H, s); ¹³C NMR (CDCl₃): δ 160.1, 159.7, 147.4, 144.8, 109.0, 106.7, 106.0, 99.30, 99.00, 81.46, 77.97, 55.22, 55.09, 27.28. LRMS (ESI+) Calcd for C₃₉H₄₆O₁₂ (M – 2H₂O + H)⁺: 671.3, Found (M – 2H₂O + H)⁺: 671.3.

Phosphoramidite ligand syntheses:

Phosphoramidites were prepared by reaction of PCl_3 with the appropriate TADDOL and amine in the presence of TEA. An example is given below for **2.112**. **2.107**,^{48,11} **2.108**,¹² **2.109**,^{48a,73} **2.110**,¹³ and **2.112**¹⁴ have been prepared previously.

Synthesis of **2.112**



To 0.179 g (0.195 mmol) of **2.199** in an oven-dried 2-dram vial with stir-bar, under N_2 , was added 0.80 mL of THF. The mixture was cooled to 0 °C, and 0.092 mL (0.66 mmol) of TEA was added, followed by dropwise addition of 18.7 μL (0.215 mmol) of PCl_3 . A white precipitate immediately formed, and the reaction was warmed to room temperature and allowed to stir for 1 h. After cooling to 0 °C, 0.14 mL (1.4 mmol) of piperidine was added, and the reaction was allowed to warm to room temperature and allowed to stir for 3 h. The reaction was then diluted with Et_2O and filtered through a pad of celite. Removal of volatile material under reduced pressure, followed by

¹¹ Alexakis, A.; Burton, J.; Vastra, J.; Benhaim, C.; Fournioux, X.; Huevel, A. V. D.; Levêque, J. M.; Mazé, M.; Rosset, S. *Eur. J. Org. Chem.* **2000**, 4011.

¹² Yu, R. T.; Rovis, T. *J. Am. Chem. Soc.* **2006**, 128, 12370.

¹³ Boele, M. D. K.; Kamer, P. C. J.; Lutz, M.; Spek, A. L.; de Vries, J. G.; van Leeuwen, P. W. M. N.; van Strijdonck, G. P. F. *Chem. Eur. J.* **2004**, 10, 6232.

¹⁴ Woodward, A. R.; Burks, H. E.; Chan, L. M.; Morken, J. P. *Org. Lett.* **2005**, 7, 5505.



purification using silica gel chromatography (hexanes/EtOAc) afforded 0.163 g (0.158 mmol, 81%) of **2.112** as a white foamy solid.

[3,5-(^tBu)₂TADDOL]PNC₅H₁₀ (2.112). Mp 192-200 °C (sealed capillary). $R_f = 0.33$ (SiO₂, 30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3075 (w), 2966 (s), 2906 (m), 2865 (m), 1784 (w), 1599 (m), 1450 (m), 1358 (m), 1252 (m), 1201 (m), 1164 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.66 (4H, s), 7.44 (2H, br s), 7.27 (1H, s), 7.21-7.25 (3H, m), 7.17 (2H, s), 5.26 (1H, dd, $J = 6.8$ Hz, $J = 1.6$ Hz), 4.70 (1H, d, $J = 6.8$ Hz), 3.32-3.49 (4H, m), 1.60-1.75 (6H, m), 1.46 (3H, s), 1.31 (18H, s), 1.30 (18H, s), 1.28 (18H, s), 1.27 (18H, s), 0.09 (3H, s); ¹³C NMR (CDCl₃): δ 149.5, 149.1, 148.7, 148.6, 146.5, 146.19, 146.17, 142.1, 141.2, 123.7, 122.0, 121.6, 120.5, 120.1, 120.0, 119.8, 109.7, 83.79, 83.62, 82.43, 82.35, 81.06, 45.10 (d, $^2J_{CP} = 20$ Hz), 35.01, 34.94, 34.83, 31.56, 31.52, 28.02, 27.53 (d, $^3J_{CP} = 4.7$ Hz), 25.50, 24.04. ³¹P NMR (CDCl₃): δ 137.8. LRMS (ESI+) Calcd for C₆₈H₁₀₂NO₄P (M)⁺: 1027.8, Found (M)⁺: 1027.8. $[\alpha]_D^{20} = -40^\circ$ ($c = 3.0$, CHCl₃).

js3-216column

expt stdh

SAMPLE		DEC. & VT	
date	Nov 30 2007	dfreq	0
solvent	CDCl3	dn	30
file	exp	dpwr	0
ACQUISITION			
sfreq	400.023	ddp	mm
in	3.000	def	200
at	35992	PROCESSING	
nu	5998.8	wfile	ft
sw	3400	proc	not used
fb	4	fn	
bs	63		
tdwr	7.1	werf	
pw	4.000	wexp	
dl	0	wds	
tof	16	wnt	
nt	8		
ct	not used		
alock	n		
gain	not used		
FLAGS			
il	n		
in	n		
dp	y		
DISPLAY			
sp	-50.1		
wp	3255.1		
vs	370		
sc	0		
wc	250		
hzmm	13.02		
is	17778.91		
rfl	3902.0		
rtp	2895.2		
th	20		
ins	1.000		
nm			



js3-216column

exp2 std13c

SAMPLE DEC. & VT

date Nov 30 2007 dfrq 400.029

solvent CDCl3 dn H1

file /export/home/~dpwr 45

jsm/js3-216col-00f 0

dm 35:11d yyy

ACQUISITION 100.599 dmf 8617

sfreq 100.599 dmf 8617

tn C13

at 0.640 lb file

np 32876 wfile

sw 25683.4 proc

fb 14200 tn

ds 4

tpwr 57 wert

pw 8.7 wexp

dl 4.000 wbs

tof 2271.7 wnt

nt 16400

ct 100

alock not used

gain not used

flags

il 9.470

in 0

dp 0

display 0

sp 17245.6

wp 1834

vs 0

sec 250

vc 68.96


hzmm 590.00


ts 8824.2

rf1 7745.2

th 100.000

ins no ph





[3,5-xylyl]TADDOL]PNC₅H₁₀ (2.159). Prepared in 75% yield according to the procedure used to synthesize **2.112** on page 198.

A white solid. $R_f = 0.15$ (SiO₂, 30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3050 (m), 2991 (s), 2928 (s), 2845 (s), 1755 (w), 1607 (s), 1442 (s), 1379 (s), 1332 (m), 1253 (s), 1219 (s), 1160 (s) cm⁻¹;

¹H NMR (CDCl₃): δ 7.47 (2H, s), 7.28 (2H, s), 7.12 (2H, s), 7.11 (2H, s), 6.90 (2H, s), 6.89 (1H, s), 6.86 (1H, s), 5.11 (1H, dd, $J = 8.5$ Hz, $J = 3.5$ Hz), 4.76 (1H, d, $J = 8.5$ Hz), 3.32-3.45 (2H, m), 3.14-3.25 (2H, m), 2.330 (6H, s), 2.323 (6H, s), 2.316 (6H, s), 2.307 (6H, s), 1.55-1.75 (6H, m), 1.44 (3H, s), 0.322 (3H, s); ¹³C NMR (CDCl₃): δ 147.2, 146.77, 146.75, 141.95, 141.94, 141.91, 137.0, 136.6, 136.4, 136.1, 128.9, 128.7, 128.6, 128.5, 126.9, 126.6, 126.5, 125.0, 111.3, 83.09, 83.06, 82.71, 82.54, 81.21, 81.07, 81.01, 44.98 (d, $^2J_{CP} = 20$ Hz), 34.65, 27.69, 27.00 (d, $^3J_{CP} = 3.6$ Hz), 25.45, 25.26, 21.66, 21.59, 21.54. ³¹P NMR (CDCl₃): δ 138.2. LRMS (ESI+) Calcd for C₄₄H₅₄NO₄P (M + H)⁺: 692.4, Found (M + H)⁺: 692.4. $[\alpha]_D^{20} = -113^\circ$ ($c = 2.3$, CHCl₃).

jss-270column



exp1 s2pu1

SAMPLE DEC. 8 VT
date Mar 1 2008 dfrq 499.774
solvent 1 CDCl3 dn HL
file /export/home/~dpr 30
jpm/jss-270column~dpr 0
nm


ACQUISITION
sfrq 499.774 dfrq 499.774
tn H1 dseq 200
at 5.000 dres 1.0
np 70058 homo n
sw 7005.9 DEC2 0
fb 4000 dfrq2 0
bs 4 dn2 1
tpwr 57 dpr2 1
pw 4.6 dot2 0
d1 4.000 dm2 n
tot 497.0 dm2 c
nt 8 dm2 200
ct 8 dseq2 1.0
atock n dres2 n
gain not used homo PROCESSING n

FLAGS
i1 n wfile
in n proc
dp n fn
hs n match
ft
not used
f

DISPLAY
sp -59.3 werr
wp 3949.4 wepp
vs 162 wbs
sc 0 wnt
wc 250
hzmm 15.80
ts 317.63
rfi 4147.1
rfp 3618.3
th 7
rms 1.000
ph



* Impurities



[3,5-xylyl]TADDOL]PNC₆H₁₂ (2.160). Prepared in 59% yield according to the procedure used to synthesize **2.112** on page 198.



A white solid. $R_f = 0.15$ (SiO₂, 30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3000 (m), 2920 (s), 2852 (s), 1603 (s), 1459 (s), 1379 (s), 1244 (m), 1215 (s), 1164 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.48


(2H, s), 7.27 (2H, s), 7.11 (2H, s), 7.10 (2H, s), 6.89 (2H, s), 6.87 (1H, s), 6.85 (1H, s), 5.12 (1H, dd, $J = 8.5$ Hz, $J = 3.5$ Hz), 4.72 (1H, d, $J = 8.5$ Hz), 3.37-3.47 (2H, m), 3.25-3.35 (2H, m), 2.318 (6H, s), 2.308 (6H, s), 2.304 (6H, s), 2.296 (6H, s), 1.62-1.87 (8H, m), 1.46 (3H, s), 0.29 (3H, s); ¹³C NMR (CDCl₃): δ 147.3, 146.82, 146.80, 137.0, 136.6, 136.5, 136.1, 128.9, 128.7, 128.6, 128.5, 127.0, 126.51, 126.48, 125.08, 125.06, 111.2, 83.27, 83.24, 82.70, 82.53, 81.19, 81.12, 80.84, 46.83 (d, $^2J_{CP} = 21$ Hz), 34.65, 31.30 (d, $^3J_{CP} = 3.3$ Hz), 27.74, 27.03, 25.34, 25.26, 21.65, 21.60, 21.55, 21.53. ³¹P NMR (CDCl₃): δ 140.9. LRMS (ESI+) Calcd for C₄₅H₅₆NO₄P (M + H)⁺: 706.4, Found (M + H)⁺: 706.5. $[\alpha]_D^{20} = -112^\circ$ ($c = 2.6$, CHCl₃).


js5-271column

expl s2pu1

SAMPLE Mar 1 2008 dfrq 499.774 DEC. & VT
 solvent CDC13 dn H1
 title /export/home/~ dpwr 30
 jpm/js5-271col-1h.- dof 0
 ACQUISITION f1d dm nnn
 sfrq 499.774 dnt 200 C
 at 5.000 dres 1.0
 nd 7005.8 homo 1.0
 sw 7005.9 dfrq2 0
 fb 4000 dn2 1
 bs 57 dpwr2 0
 tpwr 4.6 dot2 0
 pw 4.000 dm2 0
 dt 497.0 dm2 0
 nt 8 dm2 200
 ct 8 dseq2 1.0
 alock n dres2
 gain not used homo2
 PROCESSING
 f1 n wtfile ft
 in n proc y
 dp n math not used f
 ns n
 DISPLAY -86.8 wft
 SP 4023.0 weft
 VS 162 wbs
 SC 0
 WC 250
 hzmm 16.09
 f1 394.02
 f1 4147.1
 f1p 3618.3
 th 1.000
 ins ph







[3,5-(MeO)₂TADDOL]PNMe₂ (2.201). Prepared in 75% yield according to the procedure used to synthesize **2.112** on page 198.



A white solid. $R_f = 0.19$ (SiO₂, 2:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3093 (w), 3000 (m), 2937 (m), 2831 (m), 2797 (w), 1594 (s), 1459 (s), 1421 (s), 1379 (m), 1350 (s), 1312 (s), 1286

(s), 1201 (s), 1151 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 6.96 (2H, d, $J = 2.0$ Hz), 6.80 (2H, d, $J = 2.5$ Hz), 6.69 (2H, d, $J = 2.5$ Hz), 6.67 (2H, d, $J = 2.5$ Hz), 6.30-6.36 (3H, m), 6.29 (1H, t, $J = 2.0$ Hz), 5.11 (1H, dd, $J = 8.5$ Hz, $J = 3.5$ Hz), 4.67 (1H, d, $J = 8.5$ Hz), 3.73 (6H, s), 3.719 (6H, s), 3.717 (6H, s), 3.708 (6H, s), 2.79 (6H, d, $J = 11$ Hz), 1.40 (3H, s), 0.38 (3H, s); ¹³C NMR (CDCl₃): δ 160.3, 159.96, 159.91, 159.59, 149.1, 148.4, 143.8, 143.6, 111.3, 107.8, 107.15, 107.12, 105.74, 105.66, 99.34, 98.85, 98.50, 98.43, 83.02, 82.99, 82.69, 82.52, 81.13, 81.08, 81.00, 55.28, 55.20, 55.14, 35.37 (d, $^2J_{CP} = 20$ Hz), 27.71, 25.30. ³¹P NMR (CDCl₃): δ 140.6. LRMS (ESI⁺) Calcd for C₄₁H₅₀NO₁₂P (M + H)⁺: 780.3, Found (M + H)⁺: 780.3. $[\alpha]_D^{20} = -95^\circ$ ($c = 2.6$, CHCl₃).

js4-204column



expl s2pul


SAMPLE DEC. & VT
 date Mar 1 2008 dfrq 499.774
 solvent CDCl3 dn H1
 file exp dpr 30
 ACQUISITION dpr 30
 sfrq 499.774 H1 dm mnc
 nu 5.000 ddf 200
 at 7005.8 dseg
 sw 7005.9 dres 1.0
 fb 4000 homo n
 bs 4 DEC2 0
 tdwr 57 dfrq2
 pw 4.6 dn2 1
 dl 4.000 dpr2 0
 tot 497.0 dot2 n
 nt 8 dm2 n
 ct 8 dam2 c
 atlock n dseg2 200
 gain not used dres2 1.0
 il . FLAGS n
 in n homo2 PROCESSING n
 dn n y vitfile
 bs n y proc not used f
 DISPLAY nm ft
 sp -31.6 math
 wd 3756.0 werr
 vs 119 wexp
 sc 0 wnt
 wc 250 wnt
 hzmm 15.02 wft
 fs 81.58
 rf1 4147.1
 rfp 3618.3
 th 1.000
 nm ph



exp2 szpu1

SAMPLE	date	DEC.	& VI
1	Mar 1 2008	dfrq	499.774
solvent	H ₂ O	dn	HL1
file	cddc3	exp	42
ACQUISITION	cdci3	dot	0
sfrq	125.690	dm	yvyv
tn	C13	dmm	w
at	3.000	dnt	78200
sw	200168	dses	n
sp	N00168	dm2	1.0
b	33561.1	nm0	n
bs	16000	nm0	n
tpwr	55	dfrq2	DECE2
pw	7.0	dm2	0
di	2.000	dprw2	1
tof	818.6	doft2	0
nt	1600	dm2	n
ct	48	dmn2	C
a	dm2	dm2	100000
alock	n	dsest	1.0
gain	not used	dres2	n
1	FLAGS	nm02	PROCESSING
dp	n	y	+ 1.000
ns	nm	fb/file	f
DISPLAY	29.6	PROC	not used
wd	2478.4	mth	f
ss	162	verr	159.964
sc	0	verr	159.911
wc	250	verpx	159.991
hzmm	99.97	wbs	159.964
15	500.00	wnt	159.964
rfl	13636.2		159.964
rffp	9676.4		159.964
th	8		159.964
nm	100.000		159.964
cdc	ph		159.964





[3,5-(*t*Bu)₂TADDOL]PNC₆H₁₂ (2.161). Prepared in 88% yield according to the procedure used to synthesize **2.112** on page 198.



A white solid. $R_f = 0.38$ (SiO₂, 55:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3084 (w), 2958 (s), 2907 (s), 2860 (m), 1599 (m), 1480 (m), 1451 (m), 1396 (m), 1362 (m), 1244 (m), 1198 (m) cm⁻¹; ¹H

NMR (CDCl₃): δ 7.63 (2H, d, $J = 1.5$ Hz), 7.61 (2H, d, $J = 1.5$ Hz), 7.37 (2H, br s), 7.22 (1H, t, $J = 2.0$ Hz), 7.19-7.21 (2H, m), 7.18 (1H, t, $J = 2.0$ Hz), 7.10 (2H, d, $J = 1.5$ Hz), 5.28 (1H, dd, $J = 9.0$ Hz, $J = 2.5$ Hz), 4.64 (1H, d, $J = 9.0$ Hz), 3.36-3.49 (4H, m), 1.70-1.82 (4H, m), 1.62-1.69 (4H, m), 1.47 (3H, s), 1.27 (18H, s), 1.26 (18H, s), 1.24 (18H, s), 1.22 (18H, s), 0.027 (3H, s); ¹³C NMR (CDCl₃): δ 149.5, 149.1, 148.7, 148.6, 146.6, 146.3, 142.2, 141.2, 123.7, 122.0, 121.5, 120.6, 120.12, 119.99, 119.88, 109.6, 83.92, 83.89, 83.78, 83.64, 82.45, 82.39, 80.75, 80.71, 46.76 (d, $^2J_{CP} = 21$ Hz), 34.96, 34.88, 34.75, 31.59 (d, $^3J_{CP} = 4.0$ Hz), 31.59, 31.43, 28.02, 27.10, 23.77. ³¹P NMR (CDCl₃): δ 141.5. LRMS (ESI+) Calcd for C₆₉H₁₀₄NO₄P (M + H)⁺: 1042.8, Found (M + H)⁺: 1043.3. $[\alpha]_D^{20} = -37^\circ$ ($c = 2.0$, CHCl₃).

jss-269column



exp4 szpu1


SAMPLE DEC: & VI
date Mar 1 2008 dfrq 499.774
solvent CDCl3 dn H1
file /export/home/~ dpwr 30
jpw/jss-269col-1h-~ dof 0
nmh
nm
ACQUISITION
sfrq 499.774 dm 208
in H1 dseq 1.0
at 5.000 homo 1.0
np 7005.9
fb 4000 dfrq2 0
bs 4 dn2 0
tpwr 5.7 dpwr2 1
pw 4.6 dof2 0
dl 4.000 dm2 0
tof 497.0 dmf2 0
nt 8 dseq2 200
ct 8 dres2 1.0
alock n
gain not used
FLAGS not used
PROCESSING
11 n wtitle ft
in n proc not used
dp y tn math f
hs nn
DISPLAY
sp -132.8 werr
wp 412.4 wexp
wz 500 wbp
wc 250 wnt
hzm 16.50
is 420.30
rfl 528.8
rfp 0
th 3
ins 1.000
nm ph



exp2 szpu1

SAMPLE	date	Mar	1	2008	dfrq	DEC.	& VI
solvent	CDCI3	dn			499	774	
title	exp	dpr			42		
ACQUISITION	125,660	dm			0		
strg	125,660	dm			yvvv		
infr	C13	dmf			w		
at	3,000	dms			72000		
ppm	28.105	dseq			n		
cpd	328.105	nmr			1.0		
fb	35000	nmr					
bs	18000	homo					
tprur	5	dfrq2			DECE		
dw	7.0	dm2			0		
pwl	2.000	dprw2			1		
to	818.6	dot2			0		
tof	1600	dm2			n		
nt	124	dm2			100000		
ct	124	dm2			C		
a	gain	not used					
flacs	not used	dseq2					
fl	FLACS	dm2			1.0		
dn	n	homo2			n		
dp	n	ib			PROCESSING		
ns	y	wfile			1.000		
hs	nm	proc			f		
DISPLAY	-55.1	fn:			not used		
wd	22650.4	mch			f		
ss	486						
sc	0	verr					
wc	250	wexp					
hzmm	90.48	whs					
ts	500.00	wp					
rftl	13653.1	g					
rtp	9676.4	g					
th	8						
nm	100.000	cdc					
ph							





[3,5-(*t*Bu)₂TADDOL]PNC₆H₁₂ (2.162). Prepared in 86% yield

according to the procedure used to synthesize **2.112** on page 198

using 1.5 equiv of (*S*)-(+)-2-methylpiperidine. A white solid. *R_f*

= 0.42 (SiO₂, 30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3076

(w), 2962 (s), 2907 (s), 2861 (m), 1599 (m), 1472 (m), 1388 (m),

1362 (m), 1248 (m), 1206 (m), 1160 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.65 (2H, d, *J* = 2.0

Hz), 7.63 (2H, d, *J* = 1.5 Hz), 7.34-7.46 (2H, m), 7.24 (1H, t, *J* = 1.5 Hz), 7.17-7.22 (3H,

m), 7.13 (2H, d, *J* = 2.0 Hz), 5.26 (1H, dd, *J* = 8.5 Hz, *J* = 2.5 Hz), 4.63 (1H, d, *J* = 8.5

Hz), 4.01-4.15 (1H, m), 3.45-3.57 (1H, m), 3.32-3.44 (1H, m), 1.72-1.85 (2H, m), 1.56-

1.72 (3H, m), 1.49 (4H, s), 1.29 (18H, s), 1.28 (18H, s), 1.25 (18H, s), 1.23 (18H, s),

0.042 (3H, s); ¹³C NMR (CDCl₃): δ 149.5, 149.1, 148.66, 148.61, 146.6, 146.3, 142.3,

141.4, 123.7, 122.0, 121.6, 120.5, 120.1, 119.9, 119.8, 109.4, 83.90, 83.87, 83.70, 83.56,



82.52, 82.45, 80.59, 80.56, 47.62 (d, ²*J*_{CP} = 20 Hz), 39.90 (d, ²*J*_{CP} = 19 Hz), 34.96, 34.88,


34.76, 34.75, 32.69, 31.53, 31.49, 31.58, 31.44, 28.04, 27.58, 23.80, 20.75, 18.68, 18.60.


³¹P NMR (CDCl₃): δ 137.6. LRMS (ESI+) Calcd for C₆₉H₁₀₄NO₄P (M + H)⁺: 1042.8,

Found (M + H)⁺: 1043.3. [α]_D²⁰ = -23° (*c* = 2.3, CHCl₃).

exp4 szput

[illegible]





[3,5-(*t*Bu)₂TADDOL]PNC₆H₁₂ (2.163). Prepared in 96% yield

according to the procedure used to synthesize **2.112** on page 198

using 1.5 equiv of (*S*)-(+)-2-methylpiperidine. A white solid. *R_f*

= 0.47 (SiO₂, 30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3075

(w), 2962 (s), 2911 (s), 2865 (m), 1599 (m), 1480 (m), 1396 (m),

1379 (m), 1362 (s), 1252 (s), 1201 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.71 (2H, d, *J* = 2.0

Hz), 7.67 (2H, d, *J* = 1.5 Hz), 7.48 (2H, br s), 7.22-7.28 (4H, m), 7.18 (2H, d, *J* = 2.0 Hz),

5.26 (1H, dd, *J* = 8.5 Hz, *J* = 2.5 Hz), 4.63 (1H, d, *J* = 8.5 Hz), 4.28-4.38 (1H, m), 3.49-

3.60 (1H, m), 3.31 (1H, app q, *J* = 12 Hz), 1.91-2.01 (1H, m), 1.72-1.85 (1H, m), 1.63-

1.72 (2H, m), 1.50-1.63 (5H, m), 1.321 (18H, s), 1.318 (18H, s), 1.299 (18H, s), 1.277

(18H, s), 0.078 (3H, s); ¹³C NMR (CDCl₃): δ 149.4, 149.1, 148.7, 146.7, 146.59, 146.47,

142.3, 141.1, 123.7, 122.0, 121.5, 120.5, 120.1, 119.9, 109.5, 84.12, 84.09, 84.04, 83.90,

82.21, 82.14, 80.49, 80.46, 46.64 (d, ²*J*_{CP} = 15 Hz), 39.96 (d, ²*J*_{CP} = 27 Hz), 34.98, 34.88,

34.79, 34.77, 32.21, 31.51, 31.45, 28.10, 27.69, 23.76, 20.36, 18.09, 18.05. ³¹P NMR



(CDCl₃): δ 138.0. LRMS (ESI+) Calcd for C₆₉H₁₀₄NO₄P (M + H)⁺: 1042.8, Found (M +

H)⁺: 1043.4. [α]_D²⁰ = +29° (*c* = 3.4, CHCl₃).


STANDARD PROTON PARAMETERS

exp5 szpu1

SAMPLE		date		DEC. & VI	
solvent	CDCl3	dn	dfrq	499.774	
file	exp	dot	dm		
ACQUISITION					
frq	499.774	dm			
in	H1	dm			
at	5.000	amt			
sp	2.000	dsqr			
sd	7085.5	dm2			
fb	4000	homo			
bs	4				
lpwr	57	dfrq2		DEC2	
pw	4.6	dm2			
dl	4.000	dfrq2			
tof	497.0	dot2			
nt	8	dm2			
ct	8	dm2			
alock	n	dm2			
gain	not used	dsqr2			
flags	n	dm2			
in	n	homo2			
dp	y	wfline			
nm	nm	proc			
PROCESSING					
display	-160.4	math			
wd	4115.0				
ss	4115.0	verr			
sc	0	wexp			
wc	250	wbs			
h2am	16.46	wnt			
is	922.50				
rf1	4147.0				
rfp	3618.3				
th	7				
ins	1.000				
im	ph				



219



[3,5-(*t*Bu)₂TADDOL]PN(Me)*i*Pr (2.166). Prepared in 48% yield according to the procedure used to synthesize **2.112** on page 198.



A white solid. $R_f = 0.43$ (SiO₂, 50:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3080 (w), 2962 (s), 2907 (s), 2865 (m), 1599 (m), 1476 (m), 1455 (m), 1438 (m), 1388 (m), 1358 (s), 1248 (s), 1201 (m),

1164 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.66 (2H, d, $J = 2.0$ Hz), 7.64 (2H, d, $J = 1.5$ Hz), 7.44 (2H, br s), 7.25 (1H, t, $J = 1.5$ Hz), 7.23 (2H, t, $J = 1.5$ Hz), 7.21 (1H, t, $J = 2.0$ Hz), 5.27 (1H, dd, $J = 8.5$ Hz, $J = 3.0$ Hz), 4.65 (1H, d, $J = 8.5$ Hz), 4.10-4.22 (1H, m), 2.84 (3H, d, $J = 9.5$ Hz), 1.52 (3H, s), 1.23-1.34 (78H, m), 0.069 (3H, s); ¹³C NMR (CDCl₃): δ 149.5, 149.1, 148.67, 148.64, 146.6, 146.39, 146.36, 142.3, 141.2, 123.67, 123.65, 122.0, 121.5, 120.5, 120.1, 119.89, 119.85, 109.6, 83.96, 83.93, 83.88, 83.75, 82.35, 82.29, 80.56, 80.53, 46.57 (d, $^2J_{CP} = 26$ Hz), 34.97, 34.88, 34.77, 31.50, 31.45, 30.33, 28.06, 25.16 (d, $^2J_{CP} = 19$ Hz), 23.81, 21.55 (d, $^3J_{CP} = 7.0$ Hz), 21.24, 21.22. ³¹P NMR (CDCl₃): δ 141.4. LRMS (ESI+) Calcd for C₆₇H₁₀₂NO₄P (M + H)⁺: 1016.8, Found (M + H)⁺: 1017.3. $[\alpha]_D^{20} = -27^\circ$ ($c = 3.0$, CHCl₃).

jss6-95scolunn



exp4 szpu1

SAMPLE DEC. & VT
date Mar 1 2008 dfrq 499.774
solvent CDC13 dn H1
file /export/home/~dof 30
jpm/jss6-95scol-1h-1~dof 0
nmn
c
ACQUISITION
sfreq 499.774
t1 H1
t2 H1
at 5.000
nd 70058
sw 7005.9
f2 4000
bs 4
tpr 57
pw 4.6
d1 4.000
tof 497.0
nt 8
ct 8
a1ock n
gain not used
ph 1.000
PROCESSING
11 n
in n
dp n
ns n
SD DISPLAY-132.9
wp 4105.9
vs 628
sc 0
wc 250
hzm 16.42
is 469.38
rf1 4147.2
rfp 3618.3
th 7
ins 1.000
nm



js6-95colunn
exp2 szpul


SAMPLE 1 2008 DEC. & VT
date Mar 1 2008 499.774
solvent CDCl3 H1
f1 ACQUISITION exp 42
sfrq 125.680 dm 0
in C13 dm yyy
at 3.000 dmf 7200
np 200168 dseq
sw 33361.1 dres 1.0
fb 18000 homo n
bs 4
lpwr 55 dfrq2 DEC2 0
pw 7.0 dn2 1
dl 2.000 dpr2 1
tof 818.6 dot2 0
nt 1800 dm2 n
ct 132 dm2 c
clock 10000
gain not used dseq2 1.0
j1 . FLAGS n homo2 1.0
in n y lb PROCESSING 1.00
dp n y wtfile ft
hs DISPLAY -56.6 fn math not used f
sp wp 22620.4 math
vs 283
sc 0 werr
wcmm 250 wexp
f1 90.48 wds
f2 13824.7 wnt
f3 13824.7
tbp 9876.4
ins 100.000
nm cdc ph



Phosphonite ligand syntheses:

Phosphonites were prepared by the addition of dichlorophenylphosphine to the appropriate TADDOL derivative with TEA as a base. Ligand **2.115**⁷³ has been prepared previously.

Synthesis of **2.114**



To a mixture of 0.898 g (0.981 mmol) of **2.199** and 4 Å molecular sieves in 3.9 mL of THF at 0 °C was added 0.32 ml (2.3 mmol) of TEA. Next, 150 µL (1.08 mmol) of dichlorophenylphosphine was added dropwise. The reaction was allowed to warm to room temperature and allowed to stir for 2 h. The reaction was diluted with Et₂O, filtered through celite, and concentrated under reduced pressure. Column chromatography (SiO₂, hexanes:EtOAc) afforded 0.726 g (0.711 mmol, 72%) of **2.114** as a white solid.



[3,5-(*t*Bu)₂TADDOL]PPh (2.114**).** Mp 240-260 °C (decomp.). *R_f* = 0.40 (SiO₂, 30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3075 (m), 2968 (s), 2873 (s), 1797 (w), 1595 (s), 1482 (s), 1369 (s), 1249 (s), 1205 (s), 1167 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.96-8.30 (2H, m), 7.67 (2H, s), 7.55 (2H, s), 7.38-7.53 (5H, m), 7.26 (2H, d, *J* = 6.5 Hz), 7.21 (2H, d, *J*

= 7.5 Hz), 7.14 (2H, s), 5.60 (1H, dd, $J = 8.8$ Hz, $J = 3.5$ Hz), 4.89 (1H, d, $J = 8.8$ Hz), 1.56 (3H, s) 1.20-1.33 (72H, m), 0.11 (3H, s); ^{13}C NMR (CDCl_3): δ 149.8, 149.5, 148.9, 148.8, 146.3, 145.6, 145.5, 142.6, 142.5, 141.6, 140.6, 130.5, 130.1, 129.9, 128.2, 128.1, 123.8, 122.3, 121.6, 120.7, 120.43, 120.38, 120.2, 84.39, 84.36, 84.24, 84.18, 84.09, 83.94, 83.17, 83.13, 35.02, 34.88, 34.77, 34.75, 31.53, 31.48, 31.45, 31.41, 28.09, 23.68. ^{31}P NMR (CDCl_3): δ 154.3. LRMS (ESI+) Calcd for $\text{C}_{69}\text{H}_{97}\text{O}_4\text{P}$ ($\text{M} + \text{Na} + \text{H}$) $^+$: 1044.7, Found ($\text{M} + \text{Na} + \text{H}$) $^+$: 1044.6. $[\alpha]_{\text{D}}^{20} = -44^\circ$ ($c = 0.5$, CHCl_3).

js4-287column

expi szpul



SAMPLE DEC. 8 VT
 date Apr 21 2007 dfreq 499.784
 solvent CDC13 dn H1
 file exp dpr 30
 ACQUISITION
 sfreq 499.784 dm
 tn H1 dnm
 at 5.000 dmf
 ap 70058 dseg 200
 sp 70059 dres 1.0
 fb 4000 homo n
 bs 4 temp 28.0
 tpwr 56
 pw 4.6 dfreq2 DEC2 0
 dl 4.000 dn2
 tof 497.0 dprf2 1
 nt 8 dof2 0
 ct 8 dma2 n
 alock n dma2 c
 gain not used dmf2 200
 . FLAGS dseg2
 11 n dres2 1.0
 in n homo2
 dp y wtfile PROCESSING
 ns n y wtfile ft
 DISPLAY -111.9 fnc not used
 4234.7 math
 91
 0 werr
 250 wexp
 16.94 wbs
 152.32 wnt
 4144.7
 3618.4
 7
 3.000
 nm
 ph




js4-287column

exp3 s2pu1

SAMPLE DEC. 8 VT
 date Apr 21 2007 dfreq 499.764
 solvent CDCl3 gn H1
 file /export/home/~gbrw 31
 jsm/js4-287col-13C-dof 0
 ACQUISITION: f1d 0
 srfq 125.643 dres 11905
 tn -C13 dres 1.0
 at 3.000 homo 1.0
 np 200168 homo 1.0
 sw 33361.1 temp 26.0
 fb 16000
 bs 4 dfreq2 0
 tpwr 54 dn2 1
 pw 7.0 dpwr2 1
 dl 2.000 dot2 1
 tof 818.6 dm2 1
 nt 1600 dm2 1
 ct 944 dm2 10000
 atlock n dres2 1.0
 gain not used homo2 1.0
 f1 n PROCESSING 1.0
 in n lb 1.0
 dp n y wfile ft
 hs n prgc fn
 DISPLAY -228.0 math not used f
 sp 25601.4 werr
 vp 132 wexp
 sc 0 wds
 hzmm 102.41 wft
 is 308.00
 f1 13621.1
 f2 3676.4
 tps 100.000
 nm cdc ph







[3,5-(Me)₂TADDOL]PPh (2.116). Prepared in 62% yield according to the procedure used to synthesize **2.114** on page 223.

A white solid. Mp 116-130 °C (sealed capillary, decomp.). R_f = 0.35 (SiO₂, 15:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 2991 (m), 2920 (s), 2865 (m), 1776 (w), 1599 (s), 1463 (s), 1379 (m), 1252

(m), 1214 (s), 1155 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.85-7.94 (2H, m), 7.54 (5H, s), 7.31 (2H, s), 7.12 (2H, s), 7.09 (2H, s), 6.96 (1H, s), 6.91 (2H, s), 6.86 (1H, s), 5.59 (1H, dd, J = 8.5 Hz, J = 5.5 Hz), 4.81 (1H, d, J = 8.5 Hz), 2.35 (6H, s), 2.32 (12H, s), 2.30 (6H, s), 1.62 (3H, s) 0.26 (3H, s); ¹³C NMR (CDCl₃): δ 146.7, 146.1, 141.6, 141.3, 137.1, 136.8, 136.7, 136.2, 130.3, 129.9, 129.7, 129.1, 128.9, 128.7, 128.3, 128.2, 127.2, 126.3, 125.3, 125.2, 111.2, 84.14, 83.23, 83.16, 82.79, 82.55, 82.14, 27.97, 25.08, 21.67, 21.55. ³¹P NMR (CDCl₃): δ 155.7. LRMS (ESI+) Calcd for C₄₅H₄₉O₄P (M + Na)⁺: 707.3, Found (M + Na)⁺: 706.7. $[\alpha]_D^{20}$ = -80° (c = 3.0, CHCl₃).


SpinWorks 2.5: Js4-79 column





Phosphite ligand syntheses:

Synthesis of **2.113**



To 0.179 g (0.195 mmol) of **2.199** in an oven-dried 2-dram vial with stir-bar, under N_2 , was added 0.80 mL of THF. The mixture was cooled to 0 °C, and 0.092 mL (0.66 mmol) of TEA was added, followed by dropwise addition of 18.7 μL (0.215 mmol) of PCl_3 . A white precipitate immediately formed, and the reaction was allowed to warm to room temperature and allowed to stir for 1 h. After cooling to 0 °C, 0.080 mL (1.4 mmol) of EtOH was added, and the reaction was allowed to warm to room temperature and allowed to stir for 3 h. The mixture was then diluted with Et_2O and filtered through a pad of celite. Removal of volatile material under reduced pressure, followed by purification using silica gel chromatography (hexanes/EtOAc) afforded 0.154 g (0.156 mmol, 80%) of **2.113** as a white foamy solid.



[3,5-(*t*Bu)₂TADDOL]POEt (2.113**).** Mp 72-100 °C (sealed capillary). R_f = 0.36 (SiO_2 , 30:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3075 (w), 2961 (s), 2902 (m), 2868 (m), 1788 (w), 1599 (m), 1476 (m), 1451 (m), 1392 (m), 1358 (m), 1248 (m), 1202 (m), 1168 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.55 (2H, s), 7.49 (2H, s), 7.36 (2H, s), 7.21-7.32 (6H,

m), 5.20 (1H, d, $J = 8.4$ Hz), 5.07 (1H, d, $J = 8.4$ Hz), 4.30-4.42 (1H, m), 3.97-4.10 (1H, m), 1.18-1.36 (75H, m), 1.09 (3H, s), 0.31 (3H, s); ^{13}C NMR (CDCl_3): δ 149.7, 149.2, 148.9, 148.7, 145.7, 141.3, 140.9, 123.6, 123.3, 123.0, 121.6, 121.4, 120.7, 120.3, 120.2, 120.1, 110.9, 84.21, 83.50, 83.27, 83.13, 82.51, 58.59, 34.92, 34.87, 34.83, 31.53, 31.27, 27.18, 25.17, 22.71, 17.10, 14.15. ^{31}P NMR (CDCl_3): δ 133.5. LRMS (ESI+) Calcd for $\text{C}_{65}\text{H}_{97}\text{O}_5\text{P}$ ($\text{M} + \text{Na}$) $^+$: 1011.7, Found ($\text{M} + \text{Na}$) $^+$: 1011.3. $[\alpha]_{\text{D}}^{20} = -68^\circ$ ($c = 3.0$, CHCl_3).

j56-132column



exp2 std1h


SAMPLE DEC. & VT 0
 date Dec 10 2007 dfrq
 solvent CDCl3 dn
 file /export/home/~dfr
 j56-132column~dot
 JPM/DBS/j56-132column
 ACQUISITION 400.029 dfr
 sfrq 400.029 dfr
 tn H1
 at 3.006 wfile
 np 35932 proc
 sw 5998.8 fn
 tb 3408 not used
 bs 4 weff
 tpwr 63 wexp
 pw 7.1 wbs
 dl 4.000 wnt
 tof 0
 nt 8
 ct 8
 alock not used
 gain not used
 flags
 i1 n
 i2 n
 in n
 dp y
 DISPLAY -20.2
 SD 3216.1
 WD 585
 VS 0
 SC 250
 WC 12.86
 hzmm 7397.44
 is 3901.7
 rfi 2896.2
 th 20
 ins 1.000
 nm
 ph



j56-132column
exp3 std13c

SAMPLE DEC. & VI
date Dec 10 2007 dfrq 400.029
solvent CDCl3 dn H1
file exp dpwr 45
sfrq ACQUISITION 100.539 dm 0
tn 0.313 dnm yyy
at 0.313 dnt 8617
nd 32836 1b wfile
fb 25683.4 1b wfile
bs 14200 4 proc
tpwr 5.7 fn not used
pw 8.7
dl 4.000 werr
tof 2271.7 wexp
nt 1600 wbs
ct 100 wnt
a lock n
gain not used
flags not used
i1 n
i2 n
in n
dp y
DISPLAY -267.8
SP 17414.1
WD 579
VS 0
SC 0
WC 250
hzm 69.66
ts 500.00
rf1 8882.7
rf2 7743.2
th 100.000
tms no ph





[3,5-(*t*Bu)₂TADDOL]PO^{*t*}Bu (2.167). Prepared in 19% (48%

brsm) yield according to the procedure used to synthesize **2.113** on page 230. A white solid. $R_f = 0.38$ (SiO₂, 30:1 hexanes:EtOAc);



IR (CH₂Cl₂ solution): 3080 (w), 2958 (s), 2906 (s), 2861 (s), 1771 (w), 1687 (w), 1603 (s), 1476 (s), 1451 (m), 1392 (m), 1248 (s),

1206 (s), 1168 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.38 (2H, d, $J = 1.5$ Hz), 7.37 (2H, d, $J = 2.0$ Hz), 7.30 (2H, d, $J = 2.0$ Hz), 7.24 (2H, s), 7.23 (1H, t, $J = 2.0$ Hz), 6.22 (1H, t, $J = 1.5$ Hz), 7.20 (1H, t, $J = 1.5$ Hz), 7.14 (1H, t, $J = 2.0$ Hz), 5.54 (1H, d, $J = 8.5$ Hz), 5.00 (1H, d, $J = 8.5$ Hz), 1.26 (18H, s), 1.25 (18H, s), 1.24 (18H, s), 1.06 (9H, s), 0.68 (3H, s), 0.58 (3H, s); ¹³C NMR (CDCl₃): δ 149.1, 149.0, 148.6, 148.5, 146.4, 142.0, 141.9, 123.8, 123.5, 122.5, 122.3, 121.9, 121.6, 120.4, 120.04, 120.00, 119.9, 110.8, 85.08, 85.02, 84.06, 84.03, 83.36, 83.26, 81.26, 81.22, 77.88, 77.85, 34.82, 34.80, 34.77, 34.73, 31.52, 31.46, 30.68, 30.62, 30.31, 26.19, 26.08. ³¹P NMR (CDCl₃): δ 133.9. $[\alpha]_D^{20} = -63^\circ$ ($c = 2.0$, CHCl₃).

js6-89column



exp4 szpu1

SAMPLE DEC. & VT
date Mar 1 2008 dfrq 499.774
solvent CDCl3 dn H1
file /export/home/~ dpu 30
jpm/js6-89col-1n-1~ dof 0
ACQUISITION id dm
sfrq 499.774 dm 200
f1 499.774 dm 200
at 5.000 dres 1.0
nd 7005.9 homo n
sw 7005.9 dfrq2 DEC2 0
fb 4000 dn2 1
bs 4 dn2 1
tpwr 57 dpu2 1
pw 4.6 dot2 0
dl 4.000 dm2 n
tof 497.0 dm2 c
nt 8 dm2 200
ct 8 dseq2 1.0
alock n dres2
gain not used homoz n
PROCESSING
f1 n wfile ft
f2 n proc ft
f3 n math not used
f4 n math
f5 n math
f6 n math
f7 n math
f8 n math
f9 n math
f10 n math
f11 n math
f12 n math
f13 n math
f14 n math
f15 n math
f16 n math
f17 n math
f18 n math
f19 n math
f20 n math
f21 n math
f22 n math
f23 n math
f24 n math
f25 n math
f26 n math
f27 n math
f28 n math
f29 n math
f30 n math
f31 n math
f32 n math
f33 n math
f34 n math
f35 n math
f36 n math
f37 n math
f38 n math
f39 n math
f40 n math
f41 n math
f42 n math
f43 n math
f44 n math
f45 n math
f46 n math
f47 n math
f48 n math
f49 n math
f50 n math
f51 n math
f52 n math
f53 n math
f54 n math
f55 n math
f56 n math
f57 n math
f58 n math
f59 n math
f60 n math
f61 n math
f62 n math
f63 n math
f64 n math
f65 n math
f66 n math
f67 n math
f68 n math
f69 n math
f70 n math
f71 n math
f72 n math
f73 n math
f74 n math
f75 n math
f76 n math
f77 n math
f78 n math
f79 n math
f80 n math
f81 n math
f82 n math
f83 n math
f84 n math
f85 n math
f86 n math
f87 n math
f88 n math
f89 n math
f90 n math
f91 n math
f92 n math
f93 n math
f94 n math
f95 n math
f96 n math
f97 n math
f98 n math
f99 n math
f100 n math
f101 n math
f102 n math
f103 n math
f104 n math
f105 n math
f106 n math
f107 n math
f108 n math
f109 n math
f110 n math
f111 n math
f112 n math
f113 n math
f114 n math
f115 n math
f116 n math
f117 n math
f118 n math
f119 n math
f120 n math
f121 n math
f122 n math
f123 n math
f124 n math
f125 n math
f126 n math
f127 n math
f128 n math
f129 n math
f130 n math
f131 n math
f132 n math
f133 n math
f134 n math
f135 n math
f136 n math
f137 n math
f138 n math
f139 n math
f140 n math
f141 n math
f142 n math
f143 n math
f144 n math
f145 n math
f146 n math
f147 n math
f148 n math
f149 n math
f150 n math
f151 n math
f152 n math
f153 n math
f154 n math
f155 n math
f156 n math
f157 n math
f158 n math
f159 n math
f160 n math
f161 n math
f162 n math
f163 n math
f164 n math
f165 n math
f166 n math
f167 n math
f168 n math
f169 n math
f170 n math
f171 n math
f172 n math
f173 n math
f174 n math
f175 n math
f176 n math
f177 n math
f178 n math
f179 n math
f180 n math
f181 n math
f182 n math
f183 n math
f184 n math
f185 n math
f186 n math
f187 n math
f188 n math
f189 n math
f190 n math
f191 n math
f192 n math
f193 n math
f194 n math
f195 n math
f196 n math
f197 n math
f198 n math
f199 n math
f200 n math
f201 n math
f202 n math
f203 n math
f204 n math
f205 n math
f206 n math
f207 n math
f208 n math
f209 n math
f210 n math
f211 n math
f212 n math
f213 n math
f214 n math
f215 n math
f216 n math
f217 n math
f218 n math
f219 n math
f220 n math
f221 n math
f222 n math
f223 n math
f224 n math
f225 n math
f226 n math
f227 n math
f228 n math
f229 n math
f230 n math
f231 n math
f232 n math
f233 n math
f234 n math
f235 n math
f236 n math
f237 n math
f238 n math
f239 n math
f240 n math
f241 n math
f242 n math
f243 n math
f244 n math
f245 n math
f246 n math
f247 n math
f248 n math
f249 n math
f250 n math
f251 n math
f252 n math
f253 n math
f254 n math
f255 n math
f256 n math
f257 n math
f258 n math
f259 n math
f260 n math
f261 n math
f262 n math
f263 n math
f264 n math
f265 n math
f266 n math
f267 n math
f268 n math
f269 n math
f270 n math
f271 n math
f272 n math
f273 n math
f274 n math
f275 n math
f276 n math
f277 n math
f278 n math
f279 n math
f280 n math
f281 n math
f282 n math
f283 n math
f284 n math
f285 n math
f286 n math
f287 n math
f288 n math
f289 n math
f290 n math
f291 n math
f292 n math
f293 n math
f294 n math
f295 n math
f296 n math
f297 n math
f298 n math
f299 n math
f300 n math
f301 n math
f302 n math
f303 n math
f304 n math
f305 n math
f306 n math
f307 n math
f308 n math
f309 n math
f310 n math
f311 n math
f312 n math
f313 n math
f314 n math
f315 n math
f316 n math
f317 n math
f318 n math
f319 n math
f320 n math
f321 n math
f322 n math
f323 n math
f324 n math
f325 n math
f326 n math
f327 n math
f328 n math
f329 n math
f330 n math
f331 n math
f332 n math
f333 n math
f334 n math
f335 n math
f336 n math
f337 n math
f338 n math
f339 n math
f340 n math
f341 n math
f342 n math
f343 n math
f344 n math
f345 n math
f346 n math
f347 n math
f348 n math
f349 n math
f350 n math
f351 n math
f352 n math
f353 n math
f354 n math
f355 n math
f356 n math
f357 n math
f358 n math
f359 n math
f360 n math
f361 n math
f362 n math
f363 n math
f364 n math
f365 n math
f366 n math
f367 n math
f368 n math
f369 n math
f370 n math
f371 n math
f372 n math
f373 n math
f374 n math
f375 n math
f376 n math
f377 n math
f378 n math
f379 n math
f380 n math
f381 n math
f382 n math
f383 n math
f384 n math
f385 n math
f386 n math
f387 n math
f388 n math
f389 n math
f390 n math
f391 n math
f392 n math
f393 n math
f394 n math
f395 n math
f396 n math
f397 n math
f398 n math
f399 n math
f400 n math
f401 n math
f402 n math
f403 n math
f404 n math
f405 n math
f406 n math
f407 n math
f408 n math
f409 n math
f410 n math
f411 n math
f412 n math
f413 n math
f414 n math
f415 n math
f416 n math
f417 n math
f418 n math
f419 n math
f420 n math
f421 n math
f422 n math
f423 n math
f424 n math
f425 n math
f426 n math
f427 n math
f428 n math
f429 n math
f430 n math
f431 n math
f432 n math
f433 n math
f434 n math
f435 n math
f436 n math
f437 n math
f438 n math
f439 n math
f440 n math
f441 n math
f442 n math
f443 n math
f444 n math
f445 n math
f446 n math
f447 n math
f448 n math
f449 n math
f450 n math
f451 n math
f452 n math
f453 n math
f454 n math
f455 n math
f456 n math
f457 n math
f458 n math
f459 n math
f460 n math
f461 n math
f462 n math
f463 n math
f464 n math
f465 n math
f466 n math
f467 n math
f468 n math
f469 n math
f470 n math
f471 n math
f472 n math
f473 n math
f474 n math
f475 n math
f476 n math
f477 n math
f478 n math
f479 n math
f480 n math
f481 n math
f482 n math
f483 n math
f484 n math
f485 n math
f486 n math
f487 n math
f488 n math
f489 n math
f490 n math
f491 n math
f492 n math
f493 n math
f494 n math
f495 n math
f496 n math
f497 n math
f498 n math
f499 n math
f500 n math
f501 n math
f502 n math
f503 n math
f504 n math
f505 n math
f506 n math
f507 n math
f508 n math
f509 n math
f510 n math
f511 n math
f512 n math
f513 n math
f514 n math
f515 n math
f516 n math
f517 n math
f518 n math
f519 n math
f520 n math
f521 n math
f522 n math
f523 n math
f524 n math
f525 n math
f526 n math
f527 n math
f528 n math
f529 n math
f530 n math
f531 n math
f532 n math
f533 n math
f534 n math
f535 n math
f536 n math
f537 n math
f538 n math
f539 n math
f540 n math
f541 n math
f542 n math
f543 n math
f544 n math
f545 n math
f546 n math
f547 n math
f548 n math
f549 n math
f550 n math
f551 n math
f552 n math
f553 n math
f554 n math
f555 n math
f556 n math
f557 n math
f558 n math
f559 n math
f560 n math
f561 n math
f562 n math
f563 n math
f564 n math
f565 n math
f566 n math
f567 n math
f568 n math
f569 n math
f570 n math
f571 n math
f572 n math
f573 n math
f574 n math
f575 n math
f576 n math
f577 n math
f578 n math
f579 n math
f580 n math
f581 n math
f582 n math
f583 n math
f584 n math
f585 n math
f586 n math
f587 n math
f588 n math
f589 n math
f590 n math
f591 n math
f592 n math
f593 n math
f594 n math
f595 n math
f596 n math
f597 n math
f598 n math
f599 n math
f600 n math
f601 n math
f602 n math
f603 n math
f604 n math
f605 n math
f606 n math
f607 n math
f608 n math
f609 n math
f610 n math
f611 n math
f612 n math
f613 n math
f614 n math
f615 n math
f616 n math
f617 n math
f618 n math
f619 n math
f620 n math
f621 n math
f622 n math
f623 n math
f624 n math
f625 n math
f626 n math
f627 n math
f628 n math
f629 n math
f630 n math
f631 n math
f632 n math
f633 n math
f634 n math
f635 n math
f636 n math
f637 n math
f638 n math
f639 n math
f640 n math
f641 n math
f642 n math
f643 n math
f644 n math
f645 n math
f646 n math
f647 n math
f648 n math
f649 n math
f650 n math
f651 n math
f652 n math
f653 n math
f654 n math
f655 n math
f656 n math
f657 n math
f658 n math
f659 n math
f660 n math
f661 n math
f662 n math
f663 n math
f664 n math
f665 n math
f666 n math
f667 n math
f668 n math
f669 n math
f670 n math
f671 n math
f672 n math
f673 n math
f674 n math
f675 n math
f676 n math
f677 n math
f678 n math
f679 n math
f680 n math
f681 n math
f682 n math
f683 n math
f684 n math
f685 n math
f686 n math
f687 n math
f688 n math
f689 n math
f690 n math
f691 n math
f692 n math
f693 n math
f694 n math
f695 n math
f696 n math
f697 n math
f698 n math
f699 n math
f700 n math
f701 n math
f702 n math
f703 n math
f704 n math
f705 n math
f706 n math
f707 n math
f708 n math
f709 n math
f710 n math
f711 n math
f712 n math
f713 n math
f714 n math
f715 n math
f716 n math
f717 n math
f718 n math
f719 n math
f720 n math
f721 n math
f722 n math
f723 n math
f724 n math
f725 n math
f726 n math
f727 n math
f728 n math
f729 n math
f730 n math
f731 n math
f732 n math
f733 n math
f734 n math
f735 n math
f736 n math
f737 n math
f738 n math
f739 n math
f740 n math
f741 n math
f742 n math
f743 n math
f744 n math
f745 n math
f746 n math
f747 n math
f748 n math
f749 n math
f750 n math
f751 n math
f752 n math
f753 n math
f754 n math
f755 n math
f756 n math
f757 n math
f758 n math
f759 n math
f760 n math
f761 n math
f762 n math
f763 n math
f764 n math
f765 n math
f766 n math
f767 n math
f768 n math
f769 n math
f770 n math
f771 n math
f772 n math
f773 n math
f774 n math
f775 n math
f776 n math
f777 n math
f778 n math
f779 n math
f780 n math
f781 n math
f782 n math
f783 n math
f784 n math
f785 n math
f786 n math
f787 n math
f788 n math
f789 n math
f790 n math
f791 n math
f792 n math
f793 n math
f794 n math
f795 n math
f796 n math
f797 n math
f798 n math
f799 n math
f800 n math
f801 n math
f802 n math
f803 n math
f804 n math
f805 n math
f806 n math
f807 n math
f808 n math
f809 n math
f810 n math
f811 n math
f812 n math
f813 n math
f814 n math
f815 n math
f816 n math
f817 n math
f818 n math
f819 n math
f820 n math
f821 n math
f822 n math
f823 n math
f824 n math
f825 n math
f826 n math
f827 n math
f828 n math
f829 n math
f830 n math
f831 n math
f832 n math
f833 n math
f834 n math
f835 n math
f836 n math
f837 n math
f838 n math
f839 n math
f840 n math
f841 n math
f842 n math
f843 n math
f844 n math
f845 n math
f846 n math
f847 n math
f848 n math
f849 n math
f850 n math
f851 n math
f852 n math
f853 n math
f854 n math
f855 n math
f856 n math
f857 n math
f858 n math
f859 n math
f860 n math
f861 n math
f862 n math
f863 n math
f864 n math
f865 n math
f866 n math
f867 n math
f868 n math
f869 n math
f870 n math
f871 n math
f872 n math
f873 n math
f874 n math
f875 n math
f876 n math
f877 n math
f878 n math
f879 n math
f880 n math
f881 n math
f882 n math
f883 n math
f884 n math
f885 n math
f886 n math
f887 n math
f888 n math
f889 n math
f890 n math
f891 n math
f892 n math
f893 n math
f894 n math
f895 n math
f896 n math
f897 n math
f898 n math
f899 n math
f900 n math
f901 n math
f902 n math
f903 n math
f904 n math
f905 n math
f906 n math
f907 n math
f908 n math
f909 n math
f910 n math
f911 n math
f912 n math
f913 n math
f914 n math
f915 n math
f916 n math
f917 n math
f918 n math
f919 n math
f920 n math
f921 n math
f922 n math
f923 n math
f924 n math
f925 n math
f926 n math
f927 n math
f928 n math
f929 n math
f930 n math
f931 n math
f932 n math
f933 n math
f934 n math
f935 n math
f936 n math
f937 n math
f938 n math
f939 n math
f940 n math
f941 n math
f942 n math
f943 n math
f944 n math
f945 n math
f946 n math
f947 n math
f948 n math
f949 n math
f950 n math
f951 n math
f952 n math
f953 n math
f954 n math
f955 n math
f956 n math
f957 n math
f958 n math
f959 n math
f960 n math
f961 n math
f962 n math
f963 n math
f964 n math
f965 n math
f966 n math
f967 n math
f968 n math
f969 n math
f970 n math
f971 n math
f972 n math
f973 n math
f974 n math
f975 n math
f976 n math
f977 n math
f978 n math
f979 n math
f980 n math
f981 n math
f982 n math
f983 n math
f984 n math
f985 n math
f986 n math
f987 n math
f988 n math
f989 n math
f990 n math
f991 n math
f992 n math
f993 n math
f994 n math
f995 n math
f996 n math
f997 n math
f998 n math
f999 n math
f1000 n math




```
exp7 szpu1
```

date	SAMPLE	DEC. & VI
Mar 1 2008	DDC3	499.774
solvent	exp	42
title	exp	41
ACQUISITION	exp	41
sfrq	125,660	0
tfreq	13	0
at	3,000	72000
np	200168	W
nr	33581.1	1.0
fb	1800.4	1.0
bs	55	DEC2
tpwr	7.0	0
pw	2.000	1
dl	2.000	1
tof	818.6	0
nt	1600	0
ct	188	0
clock	not used	100000
gain	not used	1.0
FLAGS	not used	1.0
11	n	PROCESSING
1n	n	1.0000
dp	y	1b
hs	nm	file
DISPLAY	-10.5	proc
wd	2256.6	math
ss	532	vert
sc	0	vert
zc	250	vert
hzm	30.31	148.614
15	500.00	148.614
rf1	13692.6	148.498
rfp	9676.4	148.498
th	8	148.498
ins	100.000	148.498
cdc	ph	148.498



4. Allylation Procedures

Synthesis of **2.49**



An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 4.9 mg (0.0054 mmol) of tris(dibenzylideneacetone)dipalladium(0), 3.6 mg (0.013 mmol) of tricyclohexylphosphine, and 0.43 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min; then, 65.5 mg (0.258 mmol) of bis(pinacolato)diboron [$B_2(\text{pin})_2$] was added, followed by 25.0 mg (0.215 mmol) of phenyl allene (**2.47**). The vial was capped and allowed to stir at ambient temperature for 4 h. Next, 0.57 mL of toluene was added, followed by 41.5 mg (0.177 mmol) of dba. This mixture was allowed to stir overnight, and water was then added. The mixture was extracted with CH_2Cl_2 , and the combined organic layers dried with anhydrous Na_2SO_4 . Volatile material was removed under reduced pressure. Silica gel chromatography of the mixture (hexanes/EtOAc) afforded 73.1 mg (0.153 mmol, 79%) of **2.49** as an off-white solid.



(E)-1,5,6-Triphenyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)octa-1,7-dien-3-one (2.49). $R_f = 0.15$ (10:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3064 (m), 3028 (m), 2979 (s), 2921 (s), 2860 (m), 1955 (w), 1881 (w), 1811 (w), 1686 (s), 1665 (s), 1607 (s), 1577 (m), 1494 (s), 1451 (s), 1305 (s), 1281 (m), 1141 (s) cm^{-1} ; 1H NMR: δ 7.50 (2H, d, $J = 7.6$ Hz), 7.13-7.42 (13H, m), 7.08 (1H, t, $J = 6.8$ Hz), 6.37 (1H, d, $J = 16$ Hz), 5.59

(1H, d, $J = 3.2$ Hz), 5.55 (1H, d, $J = 3.2$), 4.23 (1H, dt, $J = 11$ Hz, $J = 3.2$ Hz), 3.82 (1H, d, $J = 11$ Hz), 2.85 (1H, dd, $J = 16$ Hz, $J = 11$ Hz), 2.71 (1H, dd, $J = 16$ Hz, $J = 3.2$ Hz), 1.15 (6H, s), 1.13 (6H, s); ^{13}C NMR: δ 199.3, 143.5, 143.3, 141.9, 134.6, 130.8, 130.1, 128.8, 128.73, 128.65, 128.5, 128.1, 128.0, 126.5, 126.4, 126.1, 83.30, 58.29, 47.06, 44.27, 24.81, 24.49. LRMS (ESI+) Calcd for $\text{C}_{32}\text{H}_{35}\text{BO}_3$ ($\text{M} + \text{H}$) $^+$: 479.3, Found ($\text{M} + \text{H}$) $^+$: 479.3.



js6-170column

exp3 stdlh


SAMPLE DEC. & VT 0
 date Mar 11 2008 dfrq 30
 solvent CDCl3 dn 30
 file /export/home/~dvr 0
 jpm/jds/js6-170col~dot 0
 -1h.tid dm mn
 ACQUISITION 400.029 dm 200
 stfrq 400.029 dm
 in 1h1 wfile ft
 at 3.000 proc not used
 sp 5898.16 fn
 bs 3400
 tpwr 4 werr 63 wexp
 pw 7.1 wbs
 dl 4.000 wnt
 tof 0
 nt 16
 ct 8
 alock n
 gain not used
 flags n
 in n
 dp y
 DISPLAY 10.6
 sp 3083.16
 wp 5.7
 vc 0
 wc 250
 hzmm 12.36
 ts 2938.55
 rfi 3902.4
 rfp 2896.2
 th 20
 ins 1.000
 nm ph



SpinWorks 2.5: Js255column second spot (19 mg)
6 1 6 2 9 9 7 7



Synthesis of **2.50**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 1.6 mg (0.0017 mmol) of tris(dibenzylideneacetone)dipalladium(0), 1.1 mg (0.0041 mmol) of tricyclohexylphosphine, and 0.13 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 0.13 mL of THF, 0.15 mL (0.075 mmol) of a 0.50 M stock solution of (S)-**2.48** in THF, and 0.10 mL (0.068 mmol) of a 0.68 M stock solution of dba in THF were added sequentially. This mixture was allowed to stir at ambient temperature for 16 h, and then, 0.29 mL of pH 7 buffer and 0.12 mL of aqueous 30% H_2O_2 were added. After stirring this mixture for 6 h, the reaction was quenched by slow addition of saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ and extracted with CH_2Cl_2 . The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Purification using silica gel chromatography (hexanes/EtOAc) afforded 23.4 mg (0.0635 mmol, 93%) of **2.50** as a white solid.

(E)-3,4,8-Triphenyloct-7-ene-2,6-dione (2.50). Mp = 176-183 °C R_f = 0.22 (4:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3063 (m), 3025 (m), 2928 (m), 1953 (w), 1894 (w), 1805 (w), 1712 (s), 1645 (s), 1607 (s), 1497 (s), 1451 (s), 1421 (m), 1354 (s), 1253 (s) cm^{-1} ; ^1H NMR: δ 7.10-7.45 (16H, m), 6.42 (1H, d, J = 16 Hz), 4.10-4.25 (2H, m),

2.81 (1H, dd, $J = 16$ Hz, $J = 10$ Hz), 2.64 (1H, dd, $J = 16$ Hz, $J = 3.4$ Hz), 1.89 (3H, s);
 ^{13}C NMR: δ 206.6, 198.1, 142.2, 142.1, 136.4, 134.4, 130.2, 129.1, 129.0, 128.8, 128.4,
128.13, 128.10, 127.9, 126.7, 126.0, 64.87, 45.24, 43.71, 30.27. LRMS (ESI+) Calcd for
 $\text{C}_{26}\text{H}_{24}\text{O}_2$ ($\text{M} + \text{H}$) $^{+}$: 369.2, Found ($\text{M} + \text{H}$) $^{+}$: 369.2.



SpinWorks 2.5: js3-94 column




js3-94column

expl std13c

SAMPLE DEC. 8 VT
date Jan 21 2008 dfrq 400.029
solvent /export/home/~dprv nl
file /export/home/~dprv 48
jsm/obs/js313c.fid dprv yyy
ACQUISITION
sfrq 100.599 daf PROCESSING 8617
in C13
at 0.640 wfile
np 32876 proc ft
sw 25683.4 fn not used
fb 14200
bs 4 weff
tdvr 57 wds
pw 8.7 wnt
d1 4.000
tof 2271.7
nt 1600
ct 186
clock + not used
gain FLAGS
fl n
in p
dp y
DISPLAY 135.9
sp 21936.8
wp 120
vs 0
sc 250
wc 87.75
hzmm 500.00
ts 8624.2
ftf 7745.6
tpp 100.000
nm no ph



General procedure for the non-enantioselective conjugate allylation (Table 2.3):





An oven-dried 20 mL scintillation vial, equipped with a magnetic stir-bar, was charged with 6.0 mg (0.022 mmol) of bis(1,5-cyclooctadiene)nickel, 12.3 mg (0.0438 mmol) of tricyclohexylphosphine, and 1.46 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 44.2 mg (0.263 mmol) of **2.51** was added followed by 50.0 mg (0.219 mmol) of **2.62**. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for the time period given in Table 2.3 (page 120). After this time period, ~15 mL of water was added, and the mixture transferred to a separatory funnel with CH₂Cl₂. After gently swirling the layers, the organic layer was collected, and the aqueous layer washed with CH₂Cl₂ (1x). The combined organic layers were dried with anhydrous Na₂SO₄, and volatile material was removed under reduced pressure. NMR analysis of the unpurified mixture was used to determine the chemoselectivity of the reaction. Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 50.7 mg (0.187 mmol, 86%) of **2.63** contaminated with a small amount of the minor constitutional isomer formed from benzylidene conjugate allylation.

(E)-5-Allyl-1-phenyldec-1-en-3-one (2.63). The minor constitutional isomer could be removed using iterative chromatography. An oil. R_f = 0.22 (18:1 hexanes:EtOAc); IR (neat): 3064 (w), 2948 (s), 2917 (s), 2858 (s), 1942 (w), 1821 (w), 1689 (s), 1654 (s),

1608 (s), 1445 (s), 1320 (m), 1167 (m) cm^{-1} ; ^1H NMR: δ 7.48-7.59 (3H, m), 7.37 (3H, m), 6.72 (1H, d, $J = 16$ Hz), 5.76 (1H, m), 5.01 (2H, d, $J = 12$ Hz), 2.61 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz), 2.52 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz), 1.95-2.20 (3H, m), 1.15-1.40 (8H, m), 0.86 (3H, t, $J = 7.2$ Hz); ^{13}C NMR: δ 200.4, 142.2, 136.6, 134.6, 130.3, 128.9, 128.2, 126.6, 116.6, 45.23, 38.26, 34.07, 33.80, 31.97, 26.40, 22.58, 14.03. LRMS (ESI+) Calcd for $\text{C}_{19}\text{H}_{26}\text{O}$ ($\text{M} + \text{Na}$) $^+$: 293.2, Found ($\text{M} + \text{Na}$) $^+$: 293.2.

^1H NMR analysis of unpurified reaction mixture (400 MHz, CDCl_3):






(E)-5-Methyl-1-phenylocta-1,7-dien-3-one (2.202).


Prepared

according to the general procedure used to synthesize **2.63** on page

245. The minor constitutional isomer was not observed in the ^1H NMR spectrum of the unpurified reaction mixture. An oil. $R_f = 0.24$ (12:1 hexanes:EtOAc); IR (neat): 3064 (m), 2955 (s), 2917 (s), 1946 (w), 1829 (w), 1689 (s), 1666 (s), 1611 (s), 1452 (s), 1324 (s), 1173 (s) cm^{-1} ; ^1H NMR: δ 7.45-7.57 (3H, m), 7.28-7.44 (3H, m), 6.72 (1H, d, $J = 16$ Hz), 5.77 (1H, m), 5.02 (2H, d, $J = 14$ Hz), 2.66 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz), 2.43 (1H, dd, $J = 16$ Hz, $J = 8.0$ Hz), 2.19 (1H, m), 1.95-2.12 (2H, m), 0.95 (3H, d, $J = 6.4$ Hz); ^{13}C NMR: δ 200.2, 142.4, 136.7, 134.5, 130.4, 128.9, 128.2, 126.5, 116.5, 47.43, 41.20, 29.52, 19.77. LRMS (ESI+) Calcd for $\text{C}_{15}\text{H}_{18}\text{O}$ ($\text{M} + \text{Na}$) $^+$: 237.1, Found ($\text{M} + \text{Na}$) $^+$: 237.1.

^1H NMR analysis of unpurified reaction mixture (400 MHz, CDCl_3):







(E)-5-Cyclohexyl-1-phenylocta-1,7-dien-3-one (2.203). Prepared

according to the general procedure used to synthesize **2.63** on page 245. Isolated as an inseparable mixture of constitutional isomers. An oil. $R_f = 0.29$ (12:1 hexanes:EtOAc); IR (neat): 3056 (m), 2913 (s), 2842 (s), 1953 (w), 1817 (w), 1685 (s), 1662 (s), 1603 (s), 1441 (s), 1320 (m), 1181 (m) cm^{-1} ; ^1H NMR: δ 7.52 (3H, m, major), 7.37 (3H, m, major), 7.25 (2H, m, minor), 7.16 (3H, m, minor), 6.72 (1H, d, $J = 16$ Hz, major), 6.65 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz, minor), 5.93 (1H, d, $J = 16$ Hz, minor), 5.74 (1H, m, major), 5.63 (1H, m, minor), 4.90-5.05 (4H, m, major & minor), 3.28 (1H, p, $J = 7.2$ Hz, minor), 2.81 (2H, m, minor), 2.61 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz, major), 2.52 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz, major), 2.37 (2H, t, $J = 7.2$ Hz, minor), 2.16 (1H, m, major), 1.91-2.10 (3H, m, major & minor), 1.57-1.78 (10H, m, major & minor), 1.36 (1H, m, major), 0.94-1.29 (10H, m, major & minor); ^{13}C NMR: δ 200.6, 199.7, 152.5, 144.2, 142.1, 137.4, 136.2, 134.6, 130.3, 128.9, 128.3, 128.2, 128.1, 127.5, 126.5, 126.3, 116.6, 116.3, 50.06, 42.50, 41.00, 40.56, 40.52, 40.33, 39.16, 35.85, 31.67, 30.00, 29.63, 26.66, 25.86, 25.64. LRMS (ESI+) Calcd for $\text{C}_{20}\text{H}_{26}\text{O}$ ($\text{M} + \text{Na}$) $^+$: 305.2, Found ($\text{M} + \text{Na}$) $^+$: 305.2.

¹H NMR analysis of unpurified reaction mixture (400 MHz, CDCl₃):



Vinyl CH
(minor + major isomers)




(E)-5-*t*-Butyl-1-phenylocta-1,7-dien-3-one (2.204).

Prepared

according to the general procedure used to synthesize **2.63** on page


245. The minor constitutional isomer was not observed in the ^1H NMR spectrum of the unpurified reaction mixture. An oil. $R_f = 0.24$ (15:1 hexanes:EtOAc); IR (neat): 3064 (m), 2959 (s), 2870 (m), 1953 (w), 1814 (w), 1689 (s), 1654 (s), 1608 (s), 1445 (s), 1363 (s), 1185 (m) cm^{-1} ; ^1H NMR: δ 7.48-7.57 (3H, m), 7.37 (3H, m), 6.73 (1H, d, $J = 16$ Hz), 5.72 (1H, m), 4.97 (1H, d, $J = 17$ Hz), 4.91 (1H, d, $J = 10$ Hz), 2.66 (1H, dd, $J = 17$ Hz, $J = 4.0$ Hz), 2.46 (1H, dd, $J = 17$ Hz, $J = 6.0$ Hz), 2.32-2.41 (1H, m), 2.13 (1H, m), 1.78 (1H, m), 0.89 (9H, s); ^{13}C NMR: δ 200.1, 141.6, 138.4, 134.5, 130.1, 128.7, 128.0, 126.3, 115.8, 42.47, 41.99, 35.65, 33.28, 27.50. LRMS (ESI+) Calcd for $\text{C}_{18}\text{H}_{24}\text{O}$ ($\text{M} + \text{Na}$) $^+$: 279.2, Found ($\text{M} + \text{Na}$) $^+$: 279.2.



(E)-5-[(*t*-Butyldimethylsilyloxy)methyl]-1-phenylocta-1,7-dien-3-one (2.205).

Prepared according to the general procedure used to synthesize **2.61** on page 245. Isolated as the major constitutional isomer when **2.70** was used as the substrate in the conjugate allylation. Note that pH 7 buffer was used in the aqueous workup rather than water. Constitutional isomers were separable using column chromatography (SiO_2 /hexanes:EtOAc). An oil. $R_f = 0.26$ (14:1 hexanes:EtOAc); IR (neat): 3076 (m), 2955 (s), 2924 (s), 2854 (s), 1689 (s), 1662 (s), 1611 (s), 1471 (m), 1363 (m), 1251 (s), 1103 (s) cm^{-1} ; ^1H NMR: δ 7.49-7.59 (3H, m), 7.37 (3H, m), 6.72 (1H, d, $J = 16$ Hz), 5.77 (1H, m), 5.03 (2H, m), 3.56 (1H, dd, $J = 10$ Hz, $J = 4.8$ Hz), 3.51 (1H, dd, $J = 10$ Hz, $J = 5.6$ Hz), 2.77 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.53 (1H, dd, $J =$

16 Hz, $J = 6.0$ Hz), 2.11-2.31 (2H, m), 2.07 (1H, m), 0.87 (9H, s), 0.015 (3H, s), 0.0030 (3H, s); ^{13}C NMR: δ 200.2, 142.3, 136.5, 134.6, 130.3, 128.9, 128.2, 126.7, 116.6, 64.73, 41.91, 36.94, 35.63, 25.89, 18.26, -5.46. LRMS (ESI+) Calcd for $\text{C}_{21}\text{H}_{32}\text{O}_2\text{Si}$ ($\text{M} + \text{Na}$) $^+$: 367.2, Found ($\text{M} + \text{Na}$) $^+$: 367.3.




(*E*)-1-(*t*-Butyldimethylsilyloxy)-6-phenylnona-2,8-dien-4-one


(2.206). Prepared according to the general procedure used to

synthesize **2.63** on page 245. Isolated as the minor constitutional isomer when **2.70** was used as the substrate in the conjugate allylation. Note that pH 7 buffer was used in the aqueous workup rather than water. Constitutional isomers were separable using column chromatography (SiO_2 /hexanes:EtOAc). An oil. $R_f = 0.18$ (14:1 Hexanes:EtOAc); IR (neat): 3056 (m), 3029 (m), 2948 (s), 2924 (s), 2858 (s), 1697 (s), 1670 (s), 1635 (s), 1468 (m), 1359 (m), 1251 (s), 1134 (s) cm^{-1} ; ^1H NMR: δ 7.20-7.30 (2H, m), 7.10-7.20 (3H, m), 6.76 (1H, dt, $J = 16$ Hz, $J = 3.2$ Hz), 6.28 (1H, d, $J = 16$ Hz), 5.63 (1H, m), 4.96 (1H, d, $J = 16$ Hz), 4.93 (1H, d, $J = 10$ Hz), 4.28 (2H, m), 3.31 (1H, p, $J = 7.2$ Hz), 2.85 (2H, m), 2.37 (2H, t, $J = 7.2$ Hz), 0.90 (9H, s), 0.045 (6H, s); ^{13}C NMR: δ 198.9, 145.3, 144.2, 136.2, 128.4, 127.9, 127.5, 126.3, 116.7, 62.23, 46.70, 40.81, 40.60, 25.84, 18.35, -5.45. LRMS (ESI+) Calcd for $\text{C}_{21}\text{H}_{32}\text{O}_2\text{Si}$ ($\text{M} + \text{Na}$) $^+$: 367.2, Found ($\text{M} + \text{Na}$) $^+$: 367.3.

^1H NMR analysis of unpurified reaction mixture (400 MHz, CDCl_3):



Carbinol (minor isomer) Carbinol (major isomer)




(*E*)-5-[(2-*t*-Butyldimethylsilyloxy)ethyl]-1-phenylocta-1,7-dien-3-one (2.207). Prepared according to the general procedure

used to synthesize **2.63** on page 245. Note that pH 7 buffer was used in the aqueous workup rather than water. Constitutional isomers were separable using iterative column chromatography (SiO_2 /hexanes:EtOAc). An oil. R_f (major) = 0.25 (15:1 hexanes:EtOAc); R_f (minor) = 0.18 (15:1 hexanes:EtOAc); IR (major, neat): 3068 (m), 2948 (s), 2932 (s), 2847 (s), 1953 (w), 1817 (w), 1682 (s), 1658 (s), 1611 (s), 1468 (m), 1328 (m), 1254 (s) cm^{-1} ; ^1H NMR: δ (major) 7.45-7.57 (3H, m), 7.37 (3H, m), 6.72 (1H, d, $J = 16$ Hz), 5.76 (1H, m), 5.02 (2H, m), 3.66 (2H, m), 2.63 (2H, d, $J = 6.4$ Hz), 2.27 (1H, heptet, $J = 6.4$ Hz), 2.01-2.20 (2H, m), 1.45-1.67 (2H, m), 0.86 (9H, s), 0.020 (6H, s); δ (minor) 7.21-


7.30 (2H, m), 7.10-7.20 (3H, m), 6.74 (1H, dt, $J = 16$ Hz, $J = 7.2$ Hz), 6.04 (1H, d, $J = 16$ Hz), 5.62 (1H, m), 4.96 (1H, d, $J = 15$ Hz), 4.93 (1H, d, $J = 10$ Hz), 3.67 (2H, t, $J = 6.4$ Hz), 3.30 (1H, p, $J = 7.2$ Hz), 2.83 (2H, m), 2.35 (4H, m), 0.87 (9H, s), 0.023 (6H, s); ^{13}C NMR: δ (major) 200.1, 142.3, 136.4, 134.6, 130.3, 128.9, 128.2, 126.5, 116.9, 61.20, 45.29, 38.42, 36.61, 31.23, 25.92, 18.26, -5.34. LRMS (ESI+) Calcd for $\text{C}_{22}\text{H}_{34}\text{O}_2\text{Si}$ ($\text{M} + \text{Na}$) $^+$: 381.2, Found ($\text{M} + \text{Na}$) $^+$: 381.3.

^1H NMR analysis of unpurified reaction mixture (400 MHz, CDCl_3):




Vinyl CH
(minor + major
isomers)

Vinyl CH
(minor isomer)



(E)-5,5-Dimethyl-1-phenylocta-1,7-dien-3-one (2.208). Prepared

according to the general procedure used to synthesize **2.63** on page 245 using 2.0 equiv of **2.51** and 65 °C reaction temperature. The minor constitutional isomer was not observed in the ^1H NMR spectrum of the unpurified reaction mixture. An oil. R_f = 0.26 (14:1 hexanes:EtOAc); IR (neat): 3072 (m), 2959 (s), 2862 (s), 1950 (w), 1814 (w), 1685 (s), 1651 (s), 1608 (s), 1440 (s), 1332 (s), 1200 (s) cm^{-1} ; ^1H NMR: δ 7.52 (2H, m), 7.49 (1H, d, J = 16 Hz), 7.36 (3H, m), 6.72 (1H, d, J = 16 Hz), 5.84 (1H, m), 5.05 (2H, m), 2.52 (2H, s), 2.12 (2H, d, J = 7.6 Hz), 1.03 (6H, s); ^{13}C NMR: δ 200.0, 141.9, 135.0, 134.6, 130.3, 128.9, 128.3, 127.6, 117.7, 51.49, 46.85, 34.30, 27.40. LRMS (ESI+) Calcd for $\text{C}_{16}\text{H}_{20}\text{O}$ ($\text{M} + \text{Na}$) $^+$: 251.1, Found ($\text{M} + \text{Na}$) $^+$: 251.2.




(E)-4,5-dimethyl-1-phenylocta-1,7-dien-3-one (2.209). Prepared

according to the general procedure used to synthesize **2.63** on page 245. Note that the reaction was performed at 65 °C and quenched with 2.2 equiv of 1.0 M HCl in Et_2O at room temperature before the addition of water. The minor constitutional isomer was not observed in the ^1H NMR spectrum of the unpurified reaction mixture. Isolated as an inseparable mixture of diastereomers using column chromatography (SiO_2 /hexanes:EtOAc). An oil. R_f = 0.26 (14:1 hexanes:EtOAc); IR (neat): 3076 (m), 2967 (s), 2920 (s), 1942 (w), 1821 (w), 1685 (s), 1651 (s), 1608 (s), 1449 (s), 1324 (s), 1188 (m) cm^{-1} ; ^1H NMR: δ 7.58 (2H, d, J = 16 Hz, major & minor), 7.54 (4H, m, major & minor), 7.37 (6H, major & minor), 6.80 (1H, d, J = 16 Hz, major), 6.79 (1H, d, J = 16 Hz, minor), 5.76 (2H, m, major & minor), 5.02 (4H, m, major &

minor), 2.80 (1H, p, $J = 6.4$ Hz, minor), 2.73 (1H, p, $J = 6.8$ Hz, major), 1.85-2.28 (6H, m, major & minor), 1.13 (3H, d, $J = 6.8$ Hz, major), 1.06 (3H, d, $J = 7.2$ Hz, minor), 0.92 (3H, d, $J = 6.4$ Hz, major), 0.84 (3H, d, $J = 6.4$ Hz, minor); ^{13}C NMR: δ 203.6, 203.5, 142.3, 136.9, 136.65, 136.59, 134.6, 130.4, 130.3, 128.9, 128.3, 128.27, 125.3, 125.1, 116.5, 116.4, 49.60, 48.47, 39.67, 37.42, 35.31, 34.59, 17.72, 15.25, 13.65, 11.35. LRMS (ESI+) Calcd for $\text{C}_{16}\text{H}_{20}\text{O}$ ($\text{M} + \text{Na}$) $^{+}$: 251.1, Found ($\text{M} + \text{Na}$) $^{+}$: 251.1.

Asymmetric Synthesis of **2.52**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.5 mg (0.0027 mmol) of tris(dibenzylideneacetone)dipalladium, 6.3 mg (0.0064 mmol) of chiral ligand **2.111**, and 0.71 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 19.8 mg (0.118 mmol) of **2.51** was added followed by 25.0 mg (0.107 mmol) of dba. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 24 h. After this time period, water was added, and the mixture transferred to a separatory funnel with CH_2Cl_2 . After gently swirling the layers, the organic layer was collected, and the aqueous layer washed with CH_2Cl_2 (2x). The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure.

Silica gel chromatography (hexanes/EtOAc) afforded 26.3 mg (0.0952 mmol, 89%) of (*R*)-**2.52**. Spectral data were consistent with the literature.¹

Proof of stereochemistry. Stereochemical ratios were determined by comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction when using Ni(cod)₂ as the precatalyst. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH₂Cl₂, to afford 5-phenyl-2-cyclohexen-1-one (see page 358). The optical rotation was measured ($[\alpha]_D^{20} = +40^\circ$ ($c = 0.5$, CHCl₃)) and compared to the known literature value.²


Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 3 mL/min, 4% MeOH) analysis of conjugate allylation product (2.52):



¹ Mandal, S. K.; Amin, Sk. R.; Crowe, W. E.; *J. Am. Chem. Soc.* **2001**, 123, 6457.

² Hareau, G. P.-J.; Koiwa, M.; Hikichi, S.; Sato, F. *J. Am. Chem. Soc.* **1999**, 121, 3640.

General procedure for the asymmetric Ni-catalyzed conjugate allylation (Table 2.7):




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 3.0 mg (0.011 mmol) of bis(1,5-cyclooctadiene)nickel, 22.4 mg (0.0219 mmol) of chiral ligand **2.114**, and 0.44 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 44.2 mg (0.263 mmol) of **2.51** was added followed by 50.0 mg (0.219 mmol) of **2.62**. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for the time indicated in Table 2.7 (page 140). After this time period, degassed water (N_2 sparge) was added, and the mixture transferred to a separatory funnel with CH_2Cl_2 . After swirling the layers, the organic layer was collected, and the aqueous layer washed with CH_2Cl_2 (2x). The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Analysis of the unpurified reaction mixture using either GLC or ^1H NMR was used to determine the chemoselectivity of the reaction. Silica gel chromatography (hexanes/EtOAc) afforded 43.7 mg (0.162 mmol, 74%) of **(S)-2.64**.

(S,E)-4-Phenyltrideca-1,7-dien-6-one (2.64). An oil. R_f = 0.19 (SiO_2 , 40:1 hexanes:EtOAc); IR (neat): 3030 (m), 2926 (s), 1697 (s), 1667 (s), 1452 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.14-7.24 (2H, m), 7.04-7.14 (3H, m), 6.65 (1H, dt, J = 16 Hz, J = 7.2


Hz), 5.92 (1H, d, $J = 16$ Hz), 5.56 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 4.82-4.94 (2H, m) 3.22 (1H, p, $J = 7.2$ Hz), 2.76 (1H, app dd, $J = 16$ Hz, $J = 6.6$ Hz), 2.75 (1H, app dd, $J = 16$ Hz, $J = 7.6$ Hz), 2.30 (2H, t, $J = 7.2$ Hz), 2.06 (2H, q, $J = 6.8$ Hz), 1.32 (2H, p, $J = 7.2$ Hz), 1.10-1.28 (4H, m), 0.796 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 199.1, 147.6, 144.2, 136.2, 130.5, 128.3, 127.5, 126.2, 116.6, 46.04, 40.99, 40.62, 32.39, 31.31, 27.74, 22.42, 13.97. LRMS (ESI+) Calcd for $\text{C}_{19}\text{H}_{26}\text{O}$ (M) $^+$: 270.2, Found (M) $^+$: 270.7. $[\alpha]_{\text{D}}^{20} = +5.9^\circ$ ($c = 1.0$, CHCl_3).


Proof of stereochemistry. Chemoselectivity was determined using achiral GLC. Stereochemical ratios were determined by comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH_2Cl_2 , to afford 5-phenyl-2-cyclohexen-1-one (see page 358). The optical rotation was measured ($[\alpha]_{\text{D}}^{20} = +40^\circ$ ($c = 0.5$, CHCl_3)) and compared to the known literature value.⁷⁸

Achiral GLC (Ultra 1, Hewlett-Packard, 140 °C) analysis of unpurified reaction mixture:



Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 2.0 % MeOH) analysis of conjugate allylation product (**2.64**):





(*S,E*)-4-(3-Chlorophenyl)trideca-1,7-dien-6-one (2.132).

Prepared according to the general procedure used to synthesize

2.64 on page 257. An oil. R_f (major) = 0.18 (SiO₂, 40:1

hexanes:EtOAc), R_f (minor) = 0.24 (SiO₂, 40:1 hexanes:EtOAc); IR (neat): 2957 (s),

2923 (s), 2853 (m), 1673 (s), 1624 (s), 1434 (m), 1367 (m) cm⁻¹; ¹H NMR (CDCl₃): δ

7.00-7.20 (4H, m), 6.74 (1H, dt, $J = 16$ Hz, $J = 6.8$ Hz), 5.99 (1H, d, $J = 16$ Hz), 5.54

(1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 4.87-5.03 (2H, m), 3.28 (1H, p, $J = 7.2$ Hz)

2.82 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz), 2.80 (1H, dd, $J = 16$ Hz, $J = 7.6$ Hz), 2.34 (2H, m),

2.14 (2H, q, $J = 7.2$ Hz), 1.40 (2H, p, $J = 7.2$ Hz), 1.15-1.35 (4H, m), 0.86 (3H, t, $J = 7.2$

Hz); ¹³C NMR (CDCl₃): δ 198.5, 147.9, 146.4, 135.7, 134.1, 130.4, 129.6, 127.5, 126.5,


125.9, 117.0, 45.65, 40.62, 40.48, 32.43, 31.33, 27.74, 22.43, 13.97. LRMS (ESI⁺)

Calcd for C₁₉H₂₅ClO (M)⁺: 304.2, Found (M)⁺: 304.7. $[\alpha]_D^{20} = +4.4^\circ$ ($c = 2.5$, CHCl₃).



Proof of stereochemistry. Chemoselectivity was determined using achiral GLC (Note: the starting material had the same retention time as the major isomer. The reported ratios are calculated assuming 95% conversion as evident by the appearance of no starting material in the ¹H NMR spectrum of the unpurified reaction mixture.) Stereochemical ratios were determined by comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH₂Cl₂, to afford 5-(*m*-chlorophenyl)-2-cyclohexen-1-one

(see page 358). The optical rotation was measured ($[\alpha]_D^{20} = +32^\circ$ ($c = 1.0$, CHCl_3)) and compared to the known literature value.⁵⁰

Achiral GLC (Ultra 1, Hewlett-Packard, 180 °C) analysis of unpurified reaction mixture:



Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 2.0 mL/min, 3.0 % MeOH) analysis of conjugate allylation product (**2.132**):



(*S,E*)-4-(4-Methoxyphenyl)trideca-1,7-dien-6-one (**2.133**).


Prepared according to the general procedure used to synthesize **2.64** on page 257. An oil. R_f = 0.18 (SiO_2 , 17:1 pentane: Et_2O); IR (neat): 3074 (w), 2962 (s), 2924 (s), 2861 (s), 2055 (w), 1879 (w), 1671 (s), 1620 (s), 1514 (s), 1464 (m), 1243 (s), 1180 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.08 (2H, d, J = 8 Hz), 6.79 (2H, d, J = 8 Hz), 6.72 (1H, dt, J = 16 Hz, J = 7.2 Hz), 5.98 (1H, d, J = 16 Hz), 5.63 (1H, ddt, J = 17 Hz, J = 10 Hz, J = 6.8 Hz), 4.90-5.02 (2H, m), 3.75 (3H, s), 3.24 (1H, p, J = 7.6 Hz), 2.78 (2H, m), 2.34 (2H, t, J = 6.8 Hz), 2.13 (2H, q, J = 7.2 Hz), 1.39 (2H, p, J = 7.2 Hz).

Hz), 1.18-1.34 (4 H, m), 0.863 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 199.3, 157.9, 147.5, 136.3, 136.2, 130.5, 128.3, 116.5, 113.7, 55.16, 46.33, 40.80, 40.28, 32.40, 31.33, 27.75, 22.42, 13.99. LRMS (ESI+) Calcd for $\text{C}_{20}\text{H}_{28}\text{O}_2$ (M^+): 300.2, Found (M^+): 300.8. $[\alpha]_{\text{D}}^{20} = +9.5^\circ$ ($c = 3.0$, CHCl_3).


Proof of stereochemistry. Chemoselectivity was determined using achiral GLC. Stereochemical ratios were determined by comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH_2Cl_2 , to afford 5-(*p*-methoxyphenyl)-2-cyclohexen-1-one (see page 358). Subsequent 1,4-reduction with Pd/C and H_2 gave the corresponding saturated cyclohexanone derivative.⁵⁰ The optical rotation was measured ($[\alpha]_{\text{D}}^{20} = -12^\circ$ ($c = 1.0$, CHCl_3)) and compared to the known literature value.¹


¹ Takaya, Y.; Ogasawara, M.; Hayashi, T. *Tetrahedron Lett.* **1999**, 40, 6957.

Achiral GLC (Ultra 1, Hewlett-Packard, 190 °C) analysis of unpurified reaction mixture:



Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 3.0 % MeOH) analysis of conjugate allylation product (2.133):






(*S,E*)-4-*o*-Tolyltrideca-1,7-dien-6-one (2.134). Prepared

according to the general procedure used to synthesize **2.64** on page 257. An oil. $R_f = 0.19$ (SiO₂, 30:1 hexanes:EtOAc); IR (neat): 3069 (m), 3018 (m), 2930 (s), 2861 (s), 1829 (w), 1671 (s), 1627 (s), 1457 (m), 1262 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.00-7.18 (4H, m), 6.73 (1H, dt, $J = 16$ Hz, $J = 7.2$ Hz), 5.99 (1H, dt, $J = 16$ Hz, $J = 1.2$ Hz), 5.63 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.88-5.02 (2H, m) 3.60 (1H, p, $J = 6.8$ Hz), 2.83 (2H, d, $J = 6.8$ Hz), 2.35 (5H, m), 2.13 (2H, q, $J = 6.8$ Hz), 1.39 (2H, p, $J = 7.2$ Hz), 1.18-1.35 (4 H, m), 0.87 (3H, t, $J = 6.8$ Hz); ¹³C NMR (CDCl₃): δ 199.2, 147.5, 142.5, 136.2, 135.9, 130.5, 130.3, 126.0, 125.9, 125.7, 116.5, 45.79, 40.44, 35.65, 32.40, 31.32, 27.76, 22.44, 19.84, 13.97. LRMS (ESI+) Calcd for C₂₀H₂₈O (M)⁺: 284.2, Found (M)⁺: 284.8. $[\alpha]_D^{20} = +6.4^\circ$ ($c = 0.6$, CHCl₃).


Proof of stereochemistry. Chemoselectivity was determined using ¹H NMR spectroscopy; the minor isomer was not observed in the ¹H NMR spectrum of the unpurified reaction mixture. Stereochemical ratios were determined by comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH₂Cl₂, to afford 5-(*o*-tolyl)-2-cyclohexen-1-one (see page 358). Subsequent 1,4-reduction with Pd/C and H₂ gave the corresponding saturated cyclohexanone derivative.⁵⁰

The optical rotation was measured ($[\alpha]_D^{20} = -37^\circ$ ($c = 0.8$, CCl_4)) and compared to the known literature value.²

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 3.0 % MeOH) analysis of conjugate allylation product (2.134):



² Ek, M.; Ahlberg, P. *Acta. Chem. Scand. Ser. B* **1984**, 38, 211.




(*S,E*)-4-[2-(Trifluoromethyl)phenyl]trideca-1,7-dien-6-one (2.135). Prepared according to the general procedure used to

synthesize **2.64** on page 257. An oil. $R_f = 0.18$ (SiO_2 , 30:1 hexanes:EtOAc); IR (neat): 2961 (s), 2930 (s), 2861 (s), 1828 (w), 1696 (s), 1457 (m), 1312 (s), 1155 (s), 1117 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.60 (1H, d, $J = 8.0$ Hz), 7.48 (1H, t, $J = 7.5$ Hz), 7.39 (1H, d, $J = 8.0$ Hz), 7.26 (1H, t, $J = 7.5$ Hz), 6.76 (1H, dt, $J = 16$ Hz, $J = 7$ Hz), 6.03 (1H, d, $J = 16$ Hz), 5.61 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7$ Hz), 4.93 (1H, d, $J = 17$ Hz), 4.91 (1H, d, $J = 10$ Hz), 3.74 (1H, p, $J = 7$ Hz), 2.84 (1H, dd, $J = 16$ Hz, $J = 7.5$ Hz), 2.80 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz), 2.40 (2H, t, $J = 7.5$ Hz), 2.15 (2H, q, $J = 7.0$ Hz), 1.40 (2H, p, $J = 7.5$ Hz), 1.18-1.35 (4 H, m), 0.87 (3H, t, $J = 7$ Hz); ^{13}C NMR (CDCl_3): δ 198.3, 148.0, 143.3, 135.6, 131.8, 130.0, 128.5 (q, $^2J_{\text{CF}} = 29$ Hz), 128.0, 126.2, 125.9 (q, $^3J_{\text{CF}} = 6.1$ Hz), 124.4 (q, $^1J_{\text{CF}} = 272$ Hz), 117.0, 46.45, 40.43, 36.08, 32.38, 31.28, 27.69, 22.39, 13.90. ^{19}F NMR (CDCl_3): δ -59.81. LRMS (ESI+) Calcd for $\text{C}_{20}\text{H}_{25}\text{F}_3\text{O}$ (M) $^+$: 338.2, Found (M) $^+$: 338.8.


Proof of stereochemistry. Chemoselectivity was determined using achiral GLC. To determine the enantiomeric purity, the title compound was subjected to ring-closing metathesis (RCM) conditions using the Hoveyda-Grubbs' second generation catalyst, in CH_2Cl_2 , to afford 5-(*o*-trifluoromethylphenyl)-2-cyclohexen-1-one (see page 358). This derivative was then analyzed by chiral SFC. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction followed by


RCM. Absolute stereochemistry was assumed to be analogous to the configuration determined for others.

Achiral GLC (Ultra 1, Hewlett-Packard, 150 °C) analysis of unpurified reaction mixture:



Chiral SFC ((R,R)-Whelk-O, Pirkle Covalent, 150 bar, 50°C, flow = 2.0 mL/min, 2.0 % MeOH) analysis of conjugate allylation-RCM product:






(*S,E*)-4-[4-(Trifluoromethyl)phenyl]trideca-1,7-dien-6-one (2.136). Prepared according to the general procedure used to synthesize **2.64** on page 257. An oil. R_f (major) = 0.19 (SiO₂, 30:1 hexanes:EtOAc), R_f (minor) = 0.24 (SiO₂, 30:1 hexanes:EtOAc); IR (neat): 2961 (s), 2930 (s), 2854 (m), 1702 (m), 1677 (s), 1620 (s), 1331 (s), 1167 (s), 1123 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.51 (2H, d, J = 8.0 Hz), 7.29 (2H, d, J = 8.0 Hz), 6.74 (1H, dt, J = 16 Hz, J = 7 Hz), 5.99 (1H, d, J = 16 Hz), 5.61 (1H, ddt, J = 17 Hz, J = 10 Hz, J = 7 Hz), 4.96 (1H, d, J = 17 Hz), 4.95 (1H, d, J = 10 Hz), 3.38 (1H, p, J = 7.5 Hz), 2.86 (1H, dd, J = 17 Hz, J = 7.0 Hz), 2.83 (1H, dd, J = 17 Hz, J = 7.5 Hz), 2.39 (1H, dd, J = 14 Hz, J = 7.0 Hz), 2.36 (1H, dd, J = 14 Hz, J = 7.0 Hz), 2.14 (2H, q, J = 7.0 Hz), 1.39 (2H, p, J = 7.0 Hz), 1.20-1.33 (4 H, m), 0.86 (3H, t, J = 7.0 Hz); ¹³C NMR (CDCl₃): δ 198.5, 148.4, 148.0, 135.6, 130.4, 128.6 (q, $^2J_{CF}$ = 32 Hz), 127.9, 125.3 (q, $^3J_{CF}$ = 3.9 Hz), 124.2 (q, $^1J_{CF}$ = 271 Hz), 117.2, 45.60, 40.65, 40.39, 32.38, 31.30, 27.69, 22.36, 13.89. ¹⁹F NMR (CDCl₃): δ -63.72. LRMS (ESI+) Calcd for C₂₀H₂₅F₃O (M)⁺: 338.2, Found (M)⁺: 338.8.


Proof of stereochemistry. Chemoselectivity was determined using achiral GLC. To determine the enantiomeric purity, the title compound was subjected to ring-closing metathesis (RCM) conditions using the Hoveyda-Grubbs' second generation catalyst, in CH₂Cl₂, to afford 5-(*p*-trifluoromethylphenyl)-2-cyclohexen-1-one (see page 358). This derivative was then analyzed by chiral GLC. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or


triphenylphosphite as the achiral ligand in the conjugate allylation reaction followed by RCM. Absolute stereochemistry was determined by comparing the optical rotation ($[\alpha]_D^{20} = +37^\circ$ ($c = 1.0$, CHCl_3)) of the RCM product to the known value.⁵⁰

Achiral GLC (Ultra 1, Hewlett-Packard, 170 °C) analysis of unpurified reaction mixture:



Chiral GLC (β -dex, Supelco, 130°C) analysis of conjugate allylation-RCM product:






(*S,E*)-4-(Furan-2-yl)trideca-1,7-dien-6-one (2.140). Prepared

according to the general procedure used to synthesize **2.64** on



page 257. An oil. Isolated as an inseparable mixture of constitutional isomers. $R_f = 0.16$ (SiO₂, 30:1 hexanes:EtOAc, both constitutional isomers); IR (neat): 3074 (w), 2955 (s), 2930 (s), 2854 (m), 1671 (s), 1626 (s), 1444 (m), 1362 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.47 (1H, d, $J = 3.2$ Hz, minor), 7.28 (1H, d, $J = 16$ Hz, minor), 7.27 (1H, dd, $J = 2.0$ Hz, $J = 1.2$ Hz, major), 6.77 (1H, dt, $J = 16$ Hz, $J = 7.2$ Hz, major), 6.64 (1H, d, $J = 2.0$ Hz, minor), 6.61 (1H, d, $J = 16$ Hz, minor), 6.46 (1H, dd, $J = 3.2$ Hz, $J = 2.0$ Hz, minor), 6.23 (1H, dd, $J = 3.2$ Hz, $J = 2.0$ Hz, major), 6.03 (1H, dt, $J = 16$ Hz, $J = 1.2$ Hz, major), 5.98 (1H, d, $J = 3.2$ Hz, major), 5.67 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz, major), 4.94-5.04 (4H, m, major + minor), 3.43 (1H, p, $J = 7.2$ Hz, major), 2.87 (1H, dd, $J = 16$ Hz, $J = 7.2$ Hz, major), 2.75 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz, major), 2.53 (1H, dd, $J = 16$ Hz, $J = 6.4$ Hz, minor), 2.45 (1H, dd, $J = 16$ Hz, $J = 5.6$ Hz, minor), 2.28-2.46 (3H, m, major + minor), 1.42 (2H, p, $J = 7.6$ Hz, major), 1.18-1.35 (12H, m, major + minor), 0.87 (3H, t, $J = 6.8$ Hz, major), 0.85 (3H, t, $J = 7.2$ Hz, minor); ¹³C NMR (CDCl₃): δ (major) 198.7, 157.1, 147.8, 140.9, 135.7, 130.4, 116.9, 109.9, 105.1, 43.06, 38.00, 34.27, 32.46, 31.34, 27.76, 22.45, 13.98. LRMS (ESI+) Calcd for C₁₇H₂₄O₂ (M)⁺: 260.2, Found (M)⁺: 260.7.

Proof of stereochemistry. Chemoselectivity was determined using achiral GLC. Stereochemical ratios were determined by comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was assumed to be analogous to the configuration determined for others.

Achiral GLC (Ultra 1, Hewlett-Packard, 145 °C) analysis of unpurified reaction mixture:



Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 1.0 % MeOH) analysis of conjugate allylation product (**2.140**):




(*S,E*)-4-(5-Fluoro-2-methylphenyl)trideca-1,7-dien-6-one

(2.137). Prepared according to the general procedure used to synthesize **2.64** on page 257. An oil. R_f = 0.18 (SiO₂, 30:1 hexanes:EtOAc); IR (neat): 3075 (w), 3031 (w), 2955 (s), 2924 (s), 2861 (s), 1841 (w), 1697 (s), 1665 (s), 1621 (s), 1495 (s), 1457 (m), 1237 (m) cm⁻¹; ¹H NMR (CDCl₃): 7.03 (1H, dd, J = 8.4 Hz, J = 6.0 Hz), 6.68-6.85 (3H, m), 5.99 (1H, dt, J = 16 Hz, J = 1.6 Hz), 5.60 (1H, ddt, J = 17 Hz, J = 10 Hz, J = 6.8 Hz), 4.96 (1H, d, J = 17 Hz), 4.93 (1H, d, J =


10 Hz), 3.57 (1H, p, $J = 7.2$ Hz), 2.82 (1H, app dd, $J = 16$ Hz, $J = 6.4$ Hz), 2.78 (1H, app dd, $J = 16$ Hz, $J = 7.2$ Hz), 2.22-2.39 (5H, m), 2.13 (2H, q, $J = 7.2$ Hz), 1.40 (2H, p, $J = 7.2$ Hz), 1.17-1.34 (4H, m), 0.86 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 198.6, 161.4 (d, $^1J_{\text{CF}} = 241$ Hz), 147.7, 144.7 (d, $^3J_{\text{CF}} = 6.2$ Hz), 135.7, 131.42, 131.36 (d, $^3J_{\text{CF}} = 4.7$ Hz), 130.4, 116.9, 112.5 (d, $^2J_{\text{CF}} = 21$ Hz), 112.4 (d, $^2J_{\text{CF}} = 21$ Hz), 45.43, 40.29, 35.73, 32.37, 31.28, 27.70, 22.39, 19.06, 13.91; ^{19}F NMR (CDCl_3): δ 93.93 (m). LRMS (ESI+) Calcd for $\text{C}_{20}\text{H}_{27}\text{FO}$ ($\text{M} + \text{Na}$) $^+$: 325.2, Found ($\text{M} + \text{Na}$) $^+$: 324.7.


Proof of stereochemistry. GLC analysis of the unpurified reaction mixture was used to determine the chemoselectivity of the reaction (Note: the starting material had the same retention time as the major isomer. The reported ratios are calculated assuming 95% conversion as evident by the appearance of no starting material in the ^1H NMR spectrum of the unpurified reaction mixture.) The enantioselectivity of the reaction was determined by measuring the enantiomeric excess of the cyclic enone formed after ring-closing metathesis (see page 362). Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction followed by RCM. Absolute stereochemistry was assumed to be analogous to the configuration determined for others.

Achiral GLC (Ultra 1, Hewlett-Packard, 180 °C) analysis of unpurified reaction mixture:



Chiral GLC (β -dex, Supelco, 135 °C) analysis of conjugate allylation-RCM product (2.142):






(*S,E*)-4-(Benzo[*d*][1,3]dioxol-5-yl)trideca-1,7-dien-6-one

(2.138). Prepared according to the general procedure used to


synthesize **2.64** on page 257. An oil. $R_f = 0.27$ (SiO₂, 15:1 pentane:Et₂O); IR (neat): 3075 (m), 2962 (s), 2930 (s), 2855 (s), 1841 (w), 1671 (s), 1627 (s), 1483 (s), 1444 (s), 1350 (m), 1243 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 6.55-6.80 (4H, m), 5.99 (1H, d, $J = 16$ Hz), 5.89 (2H, s), 5.62 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 4.90-5.05 (2H, m), 3.21 (1H, p, $J = 7.2$ Hz), 2.76 (2H, d, $J = 6.8$ Hz), 2.31 (2H, m), 2.13 (2H, q, $J = 7.6$ Hz), 1.40 (2H, p, $J = 7.2$ Hz), 1.16-1.36 (4H, m), 0.86 (3H, t, $J = 6.8$ Hz); ¹³C NMR (CDCl₃): δ 199.0, 147.6, 147.5, 145.8, 138.1, 136.2, 130.5, 120.5, 116.6, 108.1, 107.7, 100.8, 46.27, 40.84, 40.83, 32.41, 31.33, 27.76, 22.42, 13.96. LRMS (ESI+) Calcd for C₂₀H₂₆O₃ (M)⁺: 314.2, Found (M)⁺: 314.8. $[\alpha]_D^{20} = +11^\circ$ ($c = 3.0$, CHCl₃).


Proof of stereochemistry. Chemoselectivity was determined using achiral GLC. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was assumed to be analogous to the configuration determined for others.

Achiral GLC (Ultra 1, Hewlett-Packard, 190 °C) analysis of unpurified reaction mixture:



Chiral SFC (AD-H, Chiralpak, 150 bar, 50 °C, flow = 2.0 mL/min, 3.0 % MeOH) analysis of conjugate allylation product (**2.138**):






(*S,E*)-4-(1-Benzyl-1*H*-indol-3-yl)trideca-1,7-dien-6-one

(2.139). Prepared according to the general procedure used to


synthesize **2.64** on page 257. An oil. $R_f = 0.10$ (SiO₂, 2:1 pentane:CH₂Cl₂); IR (neat): 3062 (m), 2924 (s), 2848 (s), 1697 (s), 1671 (s), 1463 (s), 1350 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.68 (1H, d, $J = 7.5$ Hz), 7.00-7.32 (8H, m), 6.91 (1H, s), 6.72 (1H, dt, $J = 16$ Hz, $J = 6.8$ Hz), 6.02 (1H, d, $J = 16$ Hz), 5.75 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 5.25 (2H, s), 5.00 (1H, d, $J = 17$ Hz), 4.95 (1H, d, $J = 10$ Hz), 3.66 (1H, p, $J = 7.2$ Hz), 2.97 (2H, dd, $J = 16$ Hz, $J = 7.6$ Hz), 2.93 (1H, dd, $J = 16$ Hz, $J = 7.6$ Hz), 2.56 (1H, dd, $J = 14$ Hz, $J = 6.8$ Hz), 2.52 (1H, dd, $J = 14$ Hz, $J = 6.4$ Hz), 2.09 (2H, q, $J = 6.8$ Hz), 1.15-1.40 (6H, m), 0.87 (3H, t, $J = 7.2$ Hz); ¹³C NMR (CDCl₃): δ 199.8, 147.4, 137.6, 136.7, 130.5, 128.6, 127.4, 127.2, 126.5, 125.4, 121.6, 119.5, 118.9, 118.0, 116.3, 109.7, 49.84, 45.36, 39.74, 32.65, 32.38, 31.31, 27.68, 22.41, 13.97. LRMS (ESI+) Calcd for C₂₈H₃₃NO (M)⁺: 399.3, Found (M)⁺: 399.8. $[\alpha]_D^{20} = +15^\circ$ ($c = 3.5$, CHCl₃).

Proof of stereochemistry. Chemoselectivity was determined using ¹H NMR spectroscopy; the minor isomer was not observed in the ¹H NMR spectrum of the unpurified reaction mixture. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was assumed to be analogous to the configuration determined for others.


Chiral SFC (AS-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 4.0 % MeOH) analysis of conjugate allylation product (2.139):




Allylation
product



Racemic



Allylation product
+ racemic
coinjection



(*S,E*)-6-Phenylnona-2,8-dien-4-one (2.210). Prepared according to the general procedure used to synthesize **2.64** on page 257. An oil. R_f

= 0.20 (SiO₂, 15:1 pentane:Et₂O); IR (neat): 3069 (m), 3024 (m), 2911 (m), 1697 (s), 1671 (s), 1627 (s), 1495 (m), 1439 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.10-7.35 (5H, m), 6.76 (1H, dq, $J = 16$ Hz, $J = 6.8$ Hz), 6.03 (1H, dq, $J = 16$ Hz, $J = 1.6$ Hz), 5.63 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.86-5.05 (2H, m), 3.30 (1H, p, $J = 7.2$ Hz), 2.83 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.81 (1H, dd, $J = 16$ Hz, $J = 7.2$ Hz), 2.37 (2H, t, $J = 7.2$ Hz), 1.82 (3H, dd, $J = 6.8$ Hz, $J = 1.6$ Hz); ¹³C NMR (CDCl₃): δ 198.7, 144.2, 142.4, 136.2, 132.1, 128.3, 127.5, 126.2, 116.6, 46.03, 40.85, 40.60, 18.18. LRMS (ESI+) Calcd for C₁₅H₁₈O (M + Na)⁺: 237.1, Found (M + Na)⁺: 236.7. $[\alpha]_D^{20} = +9.3^\circ$ ($c = 2.5$, CHCl₃).

Proof of stereochemistry. Chemoselectivity was determined by ¹H NMR spectroscopy; the minor isomer was not observed in the ¹H NMR spectrum of the unpurified reaction mixture. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring-closing metathesis conditions using the Hoveyda-Grubbs second generation catalyst, in CH₂Cl₂, to afford 5-phenyl-2-cyclohexen-1-one (see page 358). The optical rotation was measured ($[\alpha]_D^{20} = +37^\circ$ ($c = 0.5$, CHCl₃)) and compared to the known literature value.⁷⁸


Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 2.0 % MeOH) analysis of conjugate allylation product (2.210):




Allylation product



Racemic



Allylation product
+ racemic
coinjection



(*S,E*)-7-Phenyldeca-3,9-dien-5-one (2.211). Prepared according to the general procedure used to synthesize **2.64** on page 257. An oil. R_f


= 0.22 (SiO₂, 20:1 hexanes:EtOAc); IR (neat): 3068 (m), 3024 (m), 2968 (s), 2924 (s), 1948 (w), 1804 (w), 1697 (s), 1627 (s), 1451 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.12-7.32 (5H, m), 6.78 (1H, dt, $J = 16$ Hz, $J = 6.8$ Hz), 5.99 (1H, d, $J = 16$ Hz), 5.63 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.86-5.05 (2H, m), 3.29 (1H, p, $J = 7.2$ Hz), 2.84 (1H, app dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.82 (1H, app dd, $J = 16$ Hz, $J = 7.2$ Hz), 2.37 (2H, t, $J = 7.2$ Hz), 2.16 (2H, p, $J = 7.2$ Hz), 1.00 (3H, t, $J = 7.2$ Hz); ¹³C NMR (CDCl₃): δ 199.1, 148.7, 144.2, 136.2, 129.6, 128.3, 127.5, 126.3, 116.6, 46.12, 40.96, 40.63, 25.50, 12.25. LRMS (ESI+) Calcd for C₁₆H₂₀O (M + Na)⁺: 251.2, Found (M + Na)⁺: 250.7. $[\alpha]_D^{20} = +8.8^\circ$ ($c = 1.0$, CHCl₃).

Proof of stereochemistry. Chemoselectivity was determined by ¹H NMR spectroscopy; the minor isomer was not observed in the ¹H NMR spectrum of the unpurified reaction mixture. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH₂Cl₂, to afford 5-phenyl-2-cyclohexen-1-one (see page 358). The optical rotation was measured ($[\alpha]_D^{20} = +41^\circ$ ($c = 0.5$, CHCl₃)) and compared to the known literature value.⁷⁸


Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 1.0 % MeOH) analysis of conjugate allylation product (2.211):




Allylation
product



Racemic



Allylation product
+ racemic
coinjection




(*S,E*)-1-Cyclohexyl-5-phenylocta-1,7-dien-3-one

(2.212).

Prepared according to the general procedure used to synthesize **2.64** on page 257. An oil. Isolated as an inseparable mixture of constitutional isomers. $R_f = 0.19$ (SiO₂, 30:1 hexanes:EtOAc, major+minor); IR (neat): 3069 (m), 3024 (m), 2924 (s), 2855 (s), 1948 (w), 1810 (w), 1700 (s), 1671 (s), 1627 (s), 1445 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.10-7.30 (5H, m), 6.64 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 5.93 (1H, dd, $J = 16$ Hz, $J = 1.2$ Hz), 5.63 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.85-5.05 (2H, m), 3.28 (1H, p, $J = 7.2$ Hz), 2.83 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.81 (1H, dd, $J = 16$ Hz, $J = 7.6$ Hz), 2.37 (2H, t, $J = 7.2$ Hz), 2.05 (1H, m), 0.92-1.80 (10H, m); ¹³C NMR (CDCl₃, major + minor isomers): δ 200.4, 199.4, 152.3, 144.2, 142.0, 137.4, 136.2, 134.6, 130.2, 128.8, 128.3, 128.2, 128.0, 127.5, 126.5, 126.2, 116.5, 116.2, 46.13, 42.58, 41.09, 40.59, 40.54, 40.44, 39.22, 35.91, 37.74, 30.09, 29.71, 26.72, 25.92, 25.69. LRMS (ESI+) Calcd for C₂₀H₂₆O (M + Na)⁺: 305.2, Found (M + Na)⁺: 304.8.


Proof of stereochemistry. Chemoselectivity was determined by ¹H NMR spectroscopy. Stereochemical ratios were determined in comparison with authentic racemic material prepared using either triphenylphosphine or triphenylphosphite as the achiral ligand in the conjugate allylation reaction. Absolute stereochemistry was determined by subjecting the conjugate allylation product to ring-closing metathesis conditions using the Hoveyda-Grubbs' second generation catalyst, in CH₂Cl₂, to afford 5-phenyl-2-cyclohexen-1-one (see page 358). The optical rotation was measured ($[\alpha]_D^{20} = +45^\circ$ ($c = 0.25$, CHCl₃)) and compared to the known literature value.⁷⁸

^1H NMR analysis of unpurified reaction mixture (400 MHz, CDCl_3):




Vinyl CH
(minor + major
isomers)


Chiral SFC (AS-H, Chiralpak, 150 bar, 50°C, flow = 0.5 mL/min, 1.0 % MeOH) analysis of conjugate allylation product (**2.212**):



Allylation
product




Racemic



Allylation product
+ racemic
coinjection

Synthesis of **2.157**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 3.0 mg (0.0033 mmol) of tris(dibenzylideneacetone)dipalladium(0), 8.2 mg (0.0079 mmol) of **2.162**, and 0.26 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 26.4 mg (0.157 mmol) of **2.51** and 20.0 mg (0.131 mmol) of **2.156** were added sequentially. The mixture was allowed to stir at ambient temperature for 14 h, and water was then added. The mixture was extracted with CH_2Cl_2 , and the combined organic layers were dried with anhydrous Na_2SO_4 . Volatile material was removed under reduced pressure, and the mixture was purified using silica gel chromatography (hexanes/EtOAc). This afforded 13.2 mg (0.0679 mmol, 52%) of **2.157**. Achiral GLC of the unpurified mixture was used to determine the chemoselectivity of the reaction. The enantiomeric purity was measured using chiral GLC analysis of **2.157**. The absolute stereochemistry of **2.157** was not determined.

5-Allyldec-1-en-3-one (2.157). An oil. $R_f = 0.26$ (SiO_2 , 30:1 hexanes:EtOAc); IR (neat): 3076 (m), 2958 (s), 2924 (s), 2848 (s), 1700 (s), 1683 (s), 1641 (m), 1611 (m), 1459 (m), 1396 (m), 1379 (m), 1299 (w), 1202 (w) cm^{-1} ; ^1H NMR (CDCl_3): δ 6.31 (1H, dd, $J = 18$ Hz, $J = 10$ Hz), 6.16 (1H, d, $J = 18$ Hz), 5.76 (1H, d, $J = 10$ Hz), 5.64-5.80 (1H, m), 4.98 (1H, d, $J = 12$ Hz), 4.98 (1H, d, $J = 16$ Hz), 2.51 (1H, dd, $J = 16$ Hz, $J =$

6.4 Hz), 2.42 (1H, dd, $J = 16$ Hz, $J = 6.0$ Hz), 1.91-2.14 (3H, m), 1.12-1.36 (8H, m), 0.84 (3H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 200.6, 136.9, 136.5, 127.6, 116.5, 44.00, 38.29, 33.84, 33.81, 32.01, 26.43, 22.61, 14.05. LRMS (ESI+) Calcd for $\text{C}_{13}\text{H}_{22}\text{O}$ ($\text{M} + \text{H}$) $^+$: 195.2, Found ($\text{M} + \text{H}$) $^+$: 195.2.

Achiral GLC (Ultra 1, Hewlett-Packard, 100 °C) analysis of unpurified reaction mixture:



Error storing signal to M:SIGNAL .BNA
DIRECTORY FULL
Storing processed peaks to M:SIGNAL .PRA


RUN# 1268 NOV 30, 1904 10:24:01

PEAK FILE : M:SIGNAL.PRA
AREA%

RT	AREA	TYPE	WIDTH	AREA%
.906	51095456	SPB	.019	99.71270
1.209	6059	BB	.022	.01182
1.680	2058	BB	.024	.00402
2.607	5165	PB	.031	.01008
7.938	4436	BB	.104	.00866
8.707	129519	PB	.120	.25276

TOTAL AREA=5.1243E+07
MUL FACTOR=1.0000E+00

Chiral GLC (β -dex, Supelco, 75°C) analysis of conjugate allylation product (**2.157**):





J56-198column

expt stidh

SAMPLE DEC. & VT 0
 date Feb 26 2008
 solvent CDCl3
 file exp dnr 30
 ACQUISITION dof 0
 sfreq 400.029 dm mn
 tn H1 dm c
 at 3.000 dmf 200
 np 35992 wifile
 sw 5998.8 proc ft
 to 3400 Tn not used
 ds 4
 tpwr 63
 pw 7.1 weft
 st 4.000 weft
 tof 4.000 wbs
 nt 8 wnt
 ct 8
 atlock n
 gain not used
 flags not used
 il n
 in n
 dp y


DISPLAY
 sp 58.0
 wp 2932.3
 vs 32
 sc 250
 vc 11.23
 hzmm 3185.51
 is 3902.2
 rfi 2896.2
 th 20
 ins 2.000
 na ph





(E)-Trideca-1,7-dien-6-one (2.158). An oil. $R_f = 0.18$ (SiO₂, 30:1 hexanes:EtOAc); IR (CDCl₃ solution): 2966 (s), 2932 (s), 2856 (s), 1700 (s), 1675 (s), 1632 (s), 1455 (s), 1358 (m), 1253 (m), 1189 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 6.80 (1H, dt, $J = 16$ Hz, $J = 7.0$ Hz), 6.07 (1H, dt, $J = 16$ Hz, $J = 1.5$ Hz), 5.76 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.0$ Hz), 5.00 (1H, dq, $J = 17$ Hz, $J = 1.5$ Hz), 4.93-4.97 (1H, m), 2.52 (2H, t, $J = 7.5$ Hz), 2.18 (2H, q, $J = 7.0$ Hz), 2.06 (2H, q, $J = 7.0$ Hz), 1.70 (2H, p, $J = 7.0$ Hz), 1.44 (2H, p, $J = 7.5$ Hz), 1.21-1.35 (4H, m), 0.87 (3H, t, $J = 7.0$ Hz); ¹³C NMR (CDCl₃): δ 200.6, 147.6, 138.1, 130.3, 115.1, 39.15, 33.17, 32.40, 31.34, 27.77, 23.27, 22.41, 13.95. LRMS (ESI+) Calcd for C₁₃H₂₂O (M + H)⁺: 195.2, Found (M + H)⁺: 195.2.

Achiral GLC (Ultra 1, Hewlett-Packard, 100 °C) analysis of 2.158:



Error storing signal to M:SIGNAL .BNA
 DIRECTORY FULL
 Storing processed peaks to M:SIGNAL .PRA

RUN# 967 JUN 20, 1984 13:19:29


PEAK FILE : M:SIGNAL.PRA

RT	AREA	TYPE	WIDTH	AREA%
0.902	17605616	BB	.050	99.18816
8.632	9891	PB	.110	.05547
12.036	7244	PB	.473	.04002
17.565	129064	PB	.279	.72379

TOTAL AREA=1.7832E+07

MUL FACTOR=1.0000E+00

Representative procedure for asymmetric allylation of **2.164**:




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 3.0 mg (0.0033 mmol) of tris(dibenzylideneacetone)dipalladium(0), 8.2 mg (0.0079 mmol) of **2.162**, and 0.66 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 26.4 mg (0.157 mmol) of **2.51** and 20.0 mg (0.131 mmol) of **2.164** were added sequentially. The vial was capped, sealed with electrical tape, removed from the dry-box, and heated at 70°C for 18 h. Degassed water (N_2 sparge) was then added, and the mixture was extracted with CH_2Cl_2 . The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Conversion and chemoselectivity were determined by achiral GLC analysis of the unpurified reaction mixture. Stereochemical ratios were determined on purified material using chiral GLC analysis of **2.165**. Racemic **2.165** was prepared by conducting the conjugate allylation with 5% $\text{Ni}(\text{cod})_2$ and 10% racemic **2.114** as ligand. This gave a 1.35:1 mixture of constitutional isomers favoring **2.165** in 36% combined yield. Separation was achieved using silica gel chromatography (hexanes/EtOAc). The absolute stereochemistry of **2.165** was not determined.


5-Allyl-5-methylnon-1-en-3-one (2.165). An oil. $R_f = 0.24$ (SiO_2 , 40:1 hexanes:EtOAc); ^1H NMR (CDCl_3): δ 6.33 (1H, dd, $J = 18$ Hz, $J = 10$ Hz), 6.14 (1H, d,


$J = 18$ Hz), 5.70-5.84 (1H, m), 5.70 (1H, d, $J = 10$ Hz), 4.86-5.07 (2H, m), 2.43 (2H, s), 2.02-2.18 (2H, m), 1.14-1.36 (6H, m), 0.95 (3H, s), 0.87 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 200.4, 138.0, 134.9, 127.1, 117.5, 47.98, 44.00, 39.39, 36.66, 36.11, 25.96, 25.06, 14.15. LRMS (ESI+) Calcd for $\text{C}_{13}\text{H}_{22}\text{O}$ ($\text{M} + \text{H}$) $^+$: 195.2, Found ($\text{M} + \text{H}$) $^+$: 195.2.

Achiral GLC (Ultra 1, Hewlett-Packard, 90 °C) analysis of unpurified reaction mixture:




Chiral GLC (β -dex, Supelco, 75°C) analysis of conjugate allylation product (**2.165**):





Synthesis of **2.168**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 4.5 mg (0.016 mmol) of bis(1,5-cyclooctadiene)nickel(0), 8.6 mg (0.033 mmol) of triphenylphosphine, and 0.82 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 5 min. Next, 33.1 mg (0.197 mmol) of **2.51** and 25.0 mg (0.164 mmol) of **2.164** were added sequentially. The vial was capped, taped with electrical tape, removed from the dry-box, and heated at 70 °C for 26 h. Degassed water (N₂ sparge) was then added, and the mixture was extracted with CH₂Cl₂. The combined organic layers were dried with anhydrous Na₂SO₄, and volatile material was removed under reduced pressure. Conversion and chemoselectivity was determined by achiral GLC analysis of the unpurified reaction mixture. Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 9.7 mg (0.050 mmol, 30%) of **2.168**.

(E)-8-Methyldodeca-1,7-dien-6-one (2.168). An oil. $R_f = 0.20$ (SiO₂, 40:1 hexanes:EtOAc); IR (neat): 3080 (m), 2958 (s), 2932 (s), 2860 (s), 1691 (s), 1620 (s), 1464 (m), 1442 (m), 1413 (m), 1362 (m), 1223 (w), 1130 (w) cm⁻¹; ¹H NMR (CDCl₃): δ 6.02 (1H, s), 5.77 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.4$ Hz), 4.90-5.04 (2H, m), 2.40 (2H, t, $J = 7.2$ Hz), 2.00-2.14 (7H, m), 1.67 (2H, p, $J = 7.6$ Hz), 1.36-1.48 (2H, m), 1.29 (2H, h, $J = 7.2$ Hz), 0.89 (3H, t, $J = 7.2$ Hz); ¹³C NMR (CDCl₃): δ 201.0, 158.6, 138.2,

123.0, 114.9, 43.57, 40.96, 33.25, 29.72, 23.43, 22.39, 19.30, 13.90. LRMS (ESI+)

Calcd for $C_{13}H_{22}O$ (M + H)⁺: 195.2, Found (M + H)⁺: 195.2.

Achiral GLC (Ultra 1, Hewlett-Packard, 90 °C) analysis of 2.168:



Closing signal file M:SIGNAL .BNC
Storing processed peaks to M:Q9126D36.PRO
DIRECTORY FULL


RUN# 1157 SEP 26, 1984 14:29:09

SIGNAL FILE: M:SIGNAL.BNC
PEAK FILE : M7Q9126D36.PRO
AREAX

RT	AREA	TYPE	WIDTH	AREAX
.907	11851168	PB	.027	98.54816
4.506	5298	PB	.066	.04406
10.237	4311	UV	.140	.03585
21.516	164988	PB	.348	1.37195

TOTAL AREA=1.2026E+07
MUL FACTOR=1.0000E+00

Synthesis of **2.170**



An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 4.2 mg (0.015 mmol) of bis(1,5-cyclooctadiene)nickel(0), 23.5 mg (0.0230 mmol) of **2.114**, and 0.50 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 30.9 mg (0.184 mmol) of **2.51** and 35.0 mg (0.153 mmol) of **2.169** were added sequentially. The vial was capped, sealed with electrical tape, removed from the dry-box, and heated at 70°C for 20 h. The reaction was quenched by the addition of unpurified, degassed (N_2 sparge) MeOH. Water was then added, and the mixture was extracted with CH_2Cl_2 . The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Chemoselectivity was determined by achiral GLC analysis of the unpurified reaction mixture.¹ Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 8.7 mg (0.032 mmol, 21%) of **2.170** contaminated with some of the other constitutional isomer derived from benzyldiene conjugate allylation. Stereochemical ratios were determined using chiral SFC analysis of the alcohol obtained from 1,2-


¹ The minor constitutional isomer was prepared by another route to obtain a GLC standard. This synthesis is given on page 303.

reduction of **2.170** using LAH at -78 °C in THF.² The absolute configuration of **2.170** was not determined.

(E)-5-Allyl-5-methyl-1-phenylnon-1-en-3-one (2.170). An oil. $R_f = 0.17$ (SiO₂, 25:1 hexanes:EtOAc); IR (CDCl₃ solution): 3067 (w), 3025 (w), 2962 (s), 2932 (s), 2860 (m), 1687 (s), 1649 (s), 1603 (s), 1573 (m), 1497 (m), 1451 (s), 1371 (m), 1328 (m), 1198 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.49-7.56 (2H, m), 7.48 (1H, d, $J = 16$ Hz), 7.34-7.40 (3H, m), 6.72 (1H, d, $J = 16$ Hz), 5.81 (1H, ddt, $J = 17$ Hz, $J = 11$ Hz, $J = 7.5$ Hz), 4.98-5.09 (2H, m) 2.52 (2H, s), 2.08-2.20 (2H, m), 1.30-1.43 (2H, m), 1.18-1.30 (4H, m), 1.00 (3H, s), 0.88 (3H, t, $J = 7.0$ Hz); ¹³C NMR (CDCl₃): δ 200.1, 141.7, 135.0, 134.6, 130.3, 128.9, 128.3, 127.6, 117.6, 49.54, 44.04, 39.40, 36.92, 25.94, 25.11, 23.40, 14.16. LRMS (ESI+) Calcd for C₁₉H₂₆O (M + H)⁺: 271.2, Found (M + H)⁺: 271.2.

² The reduction product existed as a ~1:1 mixture of diastereomers that were inseparable by column chromatography on silica gel [$R_f = 0.28$ (9:1 hexanes:EtOAc)].

Achiral GLC (Ultra 1, Hewlett-Packard, 135 °C) analysis of unpurified reaction mixture:



Closing signal file M:SIGNAL .BNC
Storing processed peaks to M:Q067E5BC.PRO
DIRECTORY FULL


RUN# 1577 MAR 17, 1985 19:23:39

SIGNAL FILE: M:SIGNAL.BNC
PEAK FILE : M:Q067E5BC.PRO
AREA%

RT	AREA	TYPE	WIDTH	AREA%
.961	48107520	SPB	.025	99.43021
1.170	4715	TBB	.029	.00975
22.545	14249	PV	.361	.02945
24.061	6625	UU	.442	.01369
28.895	7556	BU	.352	.01562
30.163	31959	UU	.365	.06605
35.592	7524	PV	.523	.01555
37.330	111549	UB	.646	.23055
42.724	5680	PP	.398	.01174
48.642	5801	PV	.586	.01199
62.859	80061	UB	.927	.16542


TOTAL AREA=4.8383E+07
MUL FACTOR=1.0000E+00

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 2.0 % MeOH) analysis of the alcohol from reduction of **2.170** with LAH at -78 °C:



13031323334353637383940414243444:

conjugate allylation-reduction
product





28293031323334353637383940414243444

racemic


js6-200column

expi szpu1

SAMPLE DEC. 8 VT
date Mar 1 2008 dfrq 499.774
solvent CDC13 dn H1
title exp dpr 30
file ACQUISITION exp 30
sfreq 499.774 da 0
in H1 dm 0
at 5.000 dmf 200
nd 70058 dseq C
sw 7005.9 dres 1.0
fb 4000 homo n
bs 4 DEC2
tpvr 57 dfrq2 0
pw 4.6 dn2
dl 4.000 dpr2- 1
tof 497.0 dot2 0
nt 16 dm2 n
ct 8 dm2 C
alock n dmt2 200
gain not used dseq2 1.0
flags not used dres2 n
il n homo2 PROCESSING n
in n y wfile
dp n y proc
hs n fn not used f
SD DISPLAY 23.3 math
WD 3903.3 weff
VS 162 weff
SC 0 weff
WC 250 wds
hzmm 15.61 wnt
IS 215.07 wft
IFI 4147.5
rfp 3618.3
th 7
ins 1.000
nm ph



Synthesis of **2.214**



To 28.1 mg (0.148 mmol) of **2.80** (see page 360) and 21.7 mg (0.222 mmol) of (*N,O*)-dimethylhydroxylamine hydrochloride in 0.60 mL of CH₂Cl₂ in a 2-dram vial with magnetic stir-bar was added 3.6 mg (0.030 mmol) of 4-(dimethylamino)pyridine (DMAP) followed by 42.6 mg (0.222 mmol) of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDCI), all in one portion. This mixture was put under N₂, and 0.03 mL (0.2 mmol) of triethylamine was added. The mixture was then allowed to stir for 3 h. The final mixture was transferred to a separatory funnel, washed with 3 M HCl (2x) followed by saturated aqueous NaHCO₃. The organic layer was dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. Silica gel chromatography (hexanes:EtOAc) afforded 26.8 mg (0.115 mmol, 78%) of Weinreb amide **2.213**.

To 48.2 mg (0.215 mmol) of **2.196** in 0.50 mL of THF at -78 °C was added 0.33 mL (0.43 mmol) of a 1.3 M solution of *t*-BuLi in pentane dropwise. This solution was allowed to stir at this temperature for 30 min and then transferred by canula to a solution of 22.8 mg (0.0977) of **2.213** (dried by azeotropic removal of water with benzene) in 0.90 mL of THF at -78 °C. After complete addition (5 min), the reaction was allowed to stir for 1 h at -78 °C. To the mixture was then added 0.2 mL of MeOH, the dry-ice acetone bath was removed, and water was added. After reaching room temperature, the mixture was transferred to a separatory funnel with water and Et₂O. The organic layer was

collected after shaking, and the aqueous layer was extracted with Et₂O (1x). The combined organics were washed with saturated aqueous NaHCO₃ and dried with anhydrous Na₂SO₄. Volatile material was removed under reduced pressure, and purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 13.8 mg (0.0510 mmol, 52%, 79% brsm) of **2.214** along with 7.7 mg of unreacted **2.213**.

***N*-Methoxy-*N*-methyl-3-phenylhex-5-enamide (2.213).** An oil. $R_f = 0.27$ (SiO₂, 2:1 hexanes:EtOAc); IR (CDCl₃ solution): 3392 (br, s), 3067 (m), 3029 (m), 2932 (s), 2852 (m), 1945 (w), 1729 (m), 1661 (s), 1493 (m), 1455 (s), 1421 (s), 1383 (s), 1180 (m), 1117 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.10-7.40 (5H, m), 5.65 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.97 (1H, d, $J = 17$ Hz), 4.93 (1H, d, $J = 10$ Hz), 3.53 (3H, s), 3.31 (1H, p, $J = 7.6$ Hz), 3.08 (3H, s), 2.72 (2H, d, $J = 6.8$ Hz), 2.30-2.50 (2H, m); ¹³C NMR (CDCl₃): δ 181.2, 144.3, 136.3, 128.2, 127.5, 126.2, 116.4, 61.11, 41.13, 40.41, 38.07, 29.68. LRMS (ESI+) Calcd for C₁₄H₁₉NO₂ (M + H)⁺: 234.2, Found (M + H)⁺: 234.2.



***(E)*-8-Allyl-5-methyl-1-phenylnon-1-en-3-one (2.214).** An oil. $R_f = 0.15$ (SiO₂, 30:1 hexanes:EtOAc); IR (CDCl₃ solution): 3076 (w), 3029 (w), 2958 (s), 2928 (s), 2864 (m), 1683 (s), 1619 (s), 1493 (m), 1451 (m), 1383 (m), 1227 (w), 1130 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.20-7.30 (2H, m), 7.12-7.20 (3H, m), 5.93 (1H, q, $J = 1.0$ Hz), 5.63 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.0$ Hz), 4.89-5.00 (2H, m), 3.26 (1H, p, $J = 7.0$ Hz), 2.74 (1H, dd, $J = 15$ Hz, $J = 6.5$ Hz), 2.69 (1H, dd, $J = 15$ Hz, $J = 7.5$ Hz), 2.35 (2H, t, $J = 8.0$ Hz), 2.04 (2H, t, $J = 7.5$ Hz), 2.02 (3H, s), 1.32-1.42 (2H, m), 1.25 (2H, h, $J = 8.0$ Hz), 0.88

(3H, t, $J = 8.0$ Hz); ^{13}C NMR (CDCl_3): δ 199.8, 159.0, 144.4, 136.4, 128.3, 127.5, 126.2, 123.4, 116.5, 50.21, 41.07, 40.84, 40.76, 29.56, 22.30, 19.27, 13.89. LRMS (ESI+) Calcd for $\text{C}_{19}\text{H}_{26}\text{O}$ ($\text{M} + \text{H}$) $^+$: 271.2, Found ($\text{M} + \text{H}$) $^+$: 271.2.


js6-186column

exp1 s2pu1

SAMPLE DEC. & VT
date Feb 23 2008 dfrq 499.774
solvent CDCl3 dn H1
title exp dpr 30
ACQUISITION exp 30
sfrq 499.774 de nmh
in H1 dm c
at 5.000 dseq 200
np 70058 dres 1.0
sw 7005.9
fb 4000 homo
bs 4
tdwr 57 dfrq2 DEC2 0
pw 4.6 dn2
dl 4.000 dpr2 1
to 497.0 dof2 0
nt 8 dm2 n
ct 8 dm2 c
atlock n dm2 200
gain not used dseq2
il FLAGS not used dres2 1.0
in n homo2
dp n
hs n
DISPLAY nm wfile ft
sp 5.2 match not used f
wd 3774.4
vs 152 weff
sc 0 wepd
wc 250 wds
hzm 15.10 wnt
is 274.73
rfi 528.8
rfd 0
th 7
ins 1.000
ph



Synthesis of **2.174**



An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.3 mg (0.0025 mmol) of tris(dibenzylideneacetone)dipalladium(0), 1.7 mg (0.0060 mmol) of tricyclohexylphosphine, and 0.20 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 25 mg (0.15 mmol) of **2.51** and 0.15 mL (0.10 mmol) of a 0.68 M solution of **2.173**¹ in THF were added sequentially. The mixture was allowed to stir at ambient temperature for 22 h, and water was then added. The mixture was extracted with CH_2Cl_2 , and the combined organic layers were dried with anhydrous Na_2SO_4 . Volatile material was removed under reduced pressure, and the mixture was purified using silica gel chromatography (hexanes/EtOAc). This afforded 23.5 mg (0.0954 mmol, 94%) of **2.174**.

(E)-Ethyl 2-hydroxy-2-styrylpent-4-enoate (2.174). An oil. $R_f = 0.21$ (SiO_2 , 8:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3510 (br, s), 3080 (m), 3025 (m), 2983 (m), 2920 (m), 1966 (w), 1734 (s), 1645 (m), 1451 (m), 1366 (m), 1286 (m), 1223 (s), 1160 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.38 (2H, d, $J = 7.2$ Hz), 7.30 (2H, t, $J = 7.2$ Hz), 7.18-7.25 (1H, m), 6.83 (1H, d, $J = 16$ Hz), 6.31 (1H, d, $J = 16$ Hz), 5.80 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 5.14 (1H, d, $J = 17$ Hz), 5.13 (1H, d, $J = 10$ Hz), 4.15-4.35 (2H, m), 3.45 (1H, s), 2.66 (1H, dd, $J = 14$ Hz, $J = 8.0$ Hz), 2.53 (1H, dd, $J = 14$ Hz, $J = 6.8$ Hz),

¹ Evans, D. A.; Johnson, J. S.; Olhava, E. J. *J. Am. Chem. Soc.* **2000**, 122, 1635.

1.30 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 174.4, 136.4, 132.0, 130.1, 129.6, 128.5, 127.8, 126.7, 119.2, 77.02, 62.39, 44.03, 14.28. LRMS (ESI+) Calcd for $\text{C}_{15}\text{H}_{18}\text{O}_3$ ($\text{M} + \text{NH}_4$) $^+$: 264.2, Found ($\text{M} + \text{NH}_4$) $^+$: 264.2.

js3-33column

exp2 std1h

SAMPLE DEC. & VT

date Jan 28 2008 dfrq 0

solvent CDCl3 dn 30

file exp dpr 30

ACQUISITION da 30

sfrq 400.029 dm 200

tn H1 dm 200

at 3.000 dmf 200

np 35992 wfile

sw 5998.8 proc

fb 3400 ft

bs 4

tpwr 63

pv 7.1 werr

dl 4.000 wexp

tof 0 wbs

nt 8 wnt

ct 8

alock n

gain not used

11 . FLAGS

in n

dp y

SP DISPLAY 33.6

wp 3042.7

vs 144

sc 0

WC 250

hzmm 12.17



is 2750.91

rtf 3903.0


th 2896.2

ins 20

nm 1.000





Asymmetric Synthesis of **2.174**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.3 mg (0.0025 mmol) of tris(dibenzylideneacetone)dipalladium(0), 3.9 mg (0.0060 mmol) of **2.110**, and 0.20 mL of C_6D_6 in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 25 mg (0.15 mmol) of **2.51** and 0.15 mL (0.10 mmol) of a 0.68 M solution of **2.173** in C_6D_6 were added sequentially. The mixture was transferred to an NMR tube and monitored by NMR. After 1 h, water was added, and the mixture was extracted with CH_2Cl_2 . The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. The residue was purified using silica gel chromatography (hexanes/EtOAc). This afforded 14 mg (0.057 mmol, 56%) of **2.174** that was determined to have an enantiomeric purity of 27% ee by analysis of **2.174** using chiral SFC. The absolute configuration was not determined.

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 1.0 mL/min, 4.0 % MeOH) analysis of **2.174**:



Synthesis of **2.181**



An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.3 mg (0.0025 mmol) of tris(dibenzylideneacetone)dipalladium(0), 1.7 mg (0.0060 mmol) of tricyclohexylphosphine, and 0.20 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 0.40 mL of THF, 27 mg (0.15 mmol) of **(Z)-2.180**,¹ and 0.15 mL (0.10 mmol) of a 0.68 M solution of **2.173** in THF were added sequentially. The mixture was allowed to stir at ambient temperature for 16 h, and water was then added. The mixture was extracted with CH_2Cl_2 , and the combined organic layers were dried with anhydrous Na_2SO_4 . Volatile material was removed under reduced pressure, and the mixture was purified using silica gel chromatography (hexanes/EtOAc). This afforded 24.1 mg (0.0926 mmol, 93%) of **2.181** as a single diastereomer by ^1H NMR analysis. The relative stereochemistry was not determined.



(E)-Ethyl 2-hydroxy-3-methyl-2-styrylpent-4-enoate (2.181). An oil. $R_f = 0.32$ (SiO_2 , 8:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3510 (br, s), 3080 (m), 3025 (m), 2979 (s), 2936 (m), 2869 (m), 1953 (w), 1877 (w), 1725 (s), 1637 (m), 1497 (m), 1447 (m), 1371 (m), 1295 (m), 1244 (s), 1142 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.35-7.47 (2H, m), 7.31

¹ Roush, W. R.; Ando, K.; Powers, D. B.; Palkowitz, A. D.; Halterman, R. L. *J. Am. Chem. Soc.* **1990**, *112*, 6339.

(2H, t, $J = 7.2$ Hz), 7.18-7.26 (1H, m), 6.86 (1H, d, $J = 16$ Hz), 6.25 (1H, d, $J = 16$ Hz), 5.79 (1H, ddd, $J = 17$ Hz, $J = 10$ Hz, $J = 8.4$ Hz), 5.00-5.15 (2H, m), 4.14-4.34 (2H, m), 3.40 (1H, s), 2.71 (1H, p, $J = 7.2$ Hz), 1.30 (3H, t, $J = 7.2$ Hz), 1.04 (3H, d, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 174.7, 138.6, 136.5, 130.7, 129.0, 128.5, 127.6, 126.6, 116.3, 79.65, 62.34, 45.77, 14.25, 13.74. LRMS (ESI+) Calcd for $\text{C}_{16}\text{H}_{20}\text{O}_3$ ($\text{M} - \text{H}_2\text{O} + \text{H}$) $^+$: 243.1, Found ($\text{M} - \text{H}_2\text{O} + \text{H}$) $^+$: 243.1.


js3-90column
exp1 std1h

SAMPLE DEC. & VT 0
date Jan 28 2008 dfreq 0
solvent CDCl3 dn 30
file ACQUISITION exp dpr 0
sfrq 400.029 da 0
in 3.001 dam mn 0
at 3.000 dnt c
nd 35982 wifile PROCESSING 200
sp 5398.8 proc ft
fb 3400 fn not used
bs 4
tdwr 63
pw 7.1 weff
dl 4.000 wexp
tof 0 wds
nt 8 wnt
ct 4
alock n
gain not used
11 . FLAGS
in n
dp y
SP DISPLAY 2.9
vd 3113.5
vs 148
sc 0
wc 250
h2mm 12.45
is 3136.05
rf1 3302.2
rfp 2896.2
th 20
ins 1.000
nm ph




js3-96column
exp2 std13c

SAMPLE DEC. 8 VT
date Jan 28 2008 dfrq 400.029
solvent CDCl3 dn H1
file exp dpr 45
ACQUISITION 0
stfrq 100.599 dm yyy
in C13 dm w
at 0.648 dnt 8617
cp 32276 lb PROCESSING 2.00
fb 258334 wftje ft
bs 14200 4 proc not used
tpwr 57 fn
pw 8.7 weff
dl 4.000 weff
tof 2271.7 wexp
nt 1600 wbs
ct 64 wnt
a1ock n
gain not used
* FLAGS
i1 n
i2 n
dp y
SP DISPLAY -102.4
WD 22071.6
VS 162
WC 0
SC 250
hzmm 88.29
ts 500.00
rfl 8825.8
rtp 7745.2
th 100.000
nm no ph



One-Pot Synthesis of **2.182**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 6.0 mg (0.0065 mmol) of tris(dibenzylideneacetone)dipalladium(0), 4.4 mg (0.016 mmol) of tricyclohexylphosphine, and 0.52 mL of toluene in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 78.7 mg (0.310 mmol) of bis(pinacolato)diboron [$B_2(\text{pin})_2$] was added followed by 30.0 mg (0.258 mmol) of phenyl allene (**2.47**). The vial was capped and allowed to stir at ambient temperature for 14 h. Next, 0.25 mL (0.17 mmol) of a 0.68 M stock solution of **2.173** in THF was added, and this mixture was allowed to stir for 3 h. After removal of volatile material under reduced pressure, 1.0 mL of THF, 0.76 mL of pH 7 buffer, and 0.30 mL of aqueous 30% H_2O_2 were added sequentially, and the resultant mixture was allowed to stir under N_2 for 6 h. The mixture was transferred to a separatory funnel and saturated aqueous $Na_2S_2O_3$ was added. This solution was extracted with CH_2Cl_2 , and the combined organic layers were washed with brine and dried with anhydrous Na_2SO_4 . Volatile material was removed under reduced pressure, and silica gel chromatography of the mixture afforded 41.2 mg (0.121 mmol, 71%) of **2.182** as a single diastereomer by 1H NMR analysis. The relative stereochemistry was not determined.

(E)-Ethyl 2-hydroxy-4-oxo-3-phenyl-2-styrylpentanoate (2.182). A white solid. Mp = 110-120 °C. R_f = 0.31 (SiO_2 , 4:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3510 (br, s),

3063 (m), 3025 (m), 2982 (m), 2924 (m), 2852 (m), 1881 (w), 1717 (s), 1607 (m), 1497 (m), 1455 (m), 1358 (s), 1231 (s), 1202 (s), 1134 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.10-7.40 (10H, m), 6.49 (1H, d, $J = 16$ Hz), 5.94 (1H, d, $J = 16$ Hz), 4.44 (1H, s), 4.27 (2H, q, $J = 7.0$ Hz), 4.21 (1H, s), 2.05 (3H, s), 1.31 (3H, t, $J = 7.0$ Hz); ^{13}C NMR (CDCl_3): δ 209.2, 174.1, 136.3, 133.0, 131.5, 130.6, 128.42, 128.36, 127.9, 127.7, 127.5, 126.5, 78.40, 65.09, 62.26, 29.50, 14.17. LRMS (ESI+) Calcd for $\text{C}_{21}\text{H}_{22}\text{O}_4$ ($\text{M} - \text{H}_2\text{O} + \text{H}$) $^+$: 321.1, Found ($\text{M} - \text{H}_2\text{O} + \text{H}$) $^+$: 321.2.

SpinWorks 2.5: js3-132B column



js3-132Bco1umh

exps std13c

SAMPLE DEC. & VT



date Jan 26 2008 dfreq 400.029
solvent CDC13 dn H1
file /export/home/~
jpw/js3-132Bco1umh do 45
js3-132Bco1umh do 0
js3-132Bco1umh do 0

ACQUISITION


sfreq 100.599 dm 8617
at 0.640 lb 2.00
np 32876 wfile
sw 25683.4 proc ft
fb 14200 fn not used
bs 4
tpwr 57 weff
pw 8.7 wexp
dl 4.000 wbs
tof 2271.7 wnt
nt 1600
ct 376

atlock n
gain not used
FLAGS

ij n
in p
dp y
DISPLAY -171.4
sp 22139.0
wp 40
vs 0
sc 250
wc 88.56
hzmm 500.00
ts 6842.4
ft1 7745.5
t1p 100.000
ins nm no ph




Enantioselective Synthesis of **2.182**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 1.6 mg (0.0018 mmol) of tris(dibenzylideneacetone)dipalladium(0) and 0.14 mL (0.0042 mmol) of a 0.030 M stock solution of triisopropylphosphine in THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 0.14 mL of THF, 0.15 mL (0.077 mmol) of a 0.50 M stock solution of *(S)*-**2.48** in THF, and 0.10 mL (0.070 mmol) of a 0.68 M stock solution of **2.173** in THF were added sequentially. The mixture was allowed to stir at ambient temperature for 17 h. After this time period, 0.30 mL of pH 7 buffer and 0.13 mL of aqueous 30% H₂O₂ were added, and the mixture allowed to stir for an additional 6 h. The reaction was quenched by the addition of saturated aqueous Na₂S₂O₃ and extracted with CH₂Cl₂. The combined organic layers were dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 14.5 mg (0.0420 mmol, 60%) of **2.182** as a single diastereomer by ¹H NMR analysis. The enantiomeric purity of **2.182** was determined to be 70% ee using chiral SFC analysis. The absolute and relative stereochemistry of **2.182** was not determined.

Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 3.0 mL/min, 3.0 % MeOH) analysis of 2.182:




allylation-oxidation product



racemic

Synthesis of **2.184**




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.0 mg (0.0073 mmol) of bis(1,5-cyclooctadiene)nickel(0), 14.8 mg (0.0145 mmol) of **2.114**, and 0.29 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 5 min. Next, 29.2 mg (0.174 mmol) of **2.51** and 25.0 mg (0.145 mmol) of **2.183**¹ were sequentially added. This mixture was allowed to stir at ambient temperature for 20 h, and then water was added, and the mixture was extracted with CH₂Cl₂. The combined organics were dried with anhydrous Na₂SO₄, and volatile material was removed under reduced pressure. Purification using silica gel chromatography (hexanes/EtOAc) afforded 23.7 mg (0.111 mmol, 76%) of **2.184** as a white solid. The enantiomeric purity of **2.184** was determined to be 37% ee using chiral HPLC analysis. The absolute stereochemistry was not determined.

(5Z,7E)-4-Methyl-8-phenylocta-1,5,7-trien-4-ol (2.184). $R_f = 0.15$ (SiO₂, 15:1 hexanes:EtOAc); IR (CDCl₃ solution): 3430 (br, s), 3080 (s), 3000 (s), 2970 (s), 2928 (s), 1848 (w), 1708 (w), 1637 (m), 1599 (m), 1489 (s), 1447 (s), 1371 (s), 1341 (m), 1260 (m), 1151 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.70 (1H, dd, $J = 16$ Hz, $J = 12$ Hz), 7.42 (2H,


¹ **2.183** was prepared in 82% yield by Wittig reaction between (*E*)-cinnamaldehyde and 1-(triphenylphosphoranylidene)-2-propanone in refluxing toluene for 17 h.

d, $J = 7.6$ Hz), 7.30 (1H, t, $J = 7.2$ Hz), 7.21 (1H, t, $J = 6.8$ Hz), 6.48 (1H, d, $J = 16$ Hz), 6.14 (1H, t, $J = 12$ Hz), 5.80-5.95 (1H, m), 5.50 (1H, d, $J = 12$ Hz), 5.18 (1H, d, $J = 10$ Hz), 5.17 (1H, d, $J = 17$ Hz), 2.45 (1H, dd, $J = 14$ Hz, $J = 6.8$ Hz), 2.34 (1H, d, $J = 14$ Hz, $J = 8.4$ Hz), 1.88 (1H, s), 1.42 (3H, s); ^{13}C NMR (CDCl_3): δ 137.4, 136.6, 134.0, 133.6, 129.6, 128.5, 127.5, 126.6, 125.6, 119.3, 73.88, 48.50, 29.28. LRMS (ESI+) Calcd for $\text{C}_{15}\text{H}_{18}\text{O}$ ($\text{M} + \text{H}$) $^+$: 215.2, Found ($\text{M} + \text{H}$) $^+$: 215.2.

Chiral HPLC (Chiralcel-OD, Daicel, 0.7 % iPrOH in hexanes, 1.0 mL/min) analysis of 2.184:



allylation product




racemic

js6-197column


expl stidh

SAMPLE DEC. & VT 0
date Feb 26 2008 dfrq 0
solvent CDC13 dn 30
title ACQUISITION exp 0
sfrq 400.029 dm 0
in H1 dm 0
at 3.000 dm 200
ap 3598.2 dnt PROCESSING
sp 5998.2 wfile
fb 33400 proc ft
bs 4 fn not used
tpwr 63
pw 7.1 weff
dl 4.000 wexp
tof 0 wbs
nt 8 wnt
ct alock n
gain not used
* FLAGS
il n
in n
dp y
DISPLAY
sp 58.0
wp 3153.1
vs 126
vc 250
hzmm 12.61
is 1793.98
rfi 3902.2
rfp 2896.2
th 20
ins 1.000
nm ph



5. Mechanistic Experiments

Synthesis of **2.87**




An oven-dried 20 mL scintillation vial, equipped with a magnetic stir-bar, was charged with 5.9 mg (0.021 mmol) of bis(1,5-cyclooctadiene)nickel, 11.9 mg (0.0426 mmol) of tricyclohexylphosphine, 60 mg of activated, crushed 4Å molecular sieves, and 1.42 mL of THF in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 43.0 mg (0.256 mmol) of **2.51** was added followed by 48.6 mg (0.213 mmol) of **2.62**. The vial was capped, sealed with electrical tape, and allowed to stir at ambient temperature for 4 h. After this time period, 37.5 μ L (0.256 mmol) of *N*-(trimethylsilyl)imidazole (**2.86**) was added and stirring was continued for an additional 1 h. Next, the mixture was filtered through a pad of celite using 10:1 hexanes:EtOAc, and volatile material was removed under reduced pressure. Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 52.3 mg (0.153 mmol, 72%) of **2.87**.

((1*E*,3*E*)-5-Allyl-1-phenyldeca-1,3-dien-3-yloxy)trimethylsilane (2.87**)**. An oil. R_f = 0.12 (SiO₂, 150:1 hexanes:EtOAc); IR (neat): 3080 (m), 3063 (m), 3029 (m), 2962 (s), 2924 (s), 2852 (s), 1945 (w), 1810 (w), 1641 (m), 1607 (m), 1493 (m), 1447 (m), 1341 (m), 1248 (s), 1181 (m), 1147 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.42 (2H, d, J = 8.0 Hz), 7.32 (2H, t, J = 7.2 Hz), 7.12-7.25 (1H, m), 6.88 (1H, d, J = 16 Hz), 6.78 (1H, d, J = 16

Hz), 5.78 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.92-5.08 (2H, m), 4.70 (1H, d, $J = 10$ Hz), 2.35-2.60 (1H, m), 2.11-2.24 (1H, m), 1.99-2.11 (1H, m), 1.41-1.56 (1H, m), 1.15-1.41 (7H, m), 0.86 (3H, t, $J = 6.8$ Hz), 0.24 (9H, s); ^{13}C NMR (CDCl_3): δ 147.7, 137.4, 137.0, 128.9, 128.5, 127.4, 126.7, 121.9, 119.4, 115.8, 41.01, 37.17, 35.96, 32.04, 27.07, 22.67, 14.12, 0.36. LRMS (ESI+) Calcd for $\text{C}_{22}\text{H}_{34}\text{OSi}$ ($\text{M} + \text{H}$) $^+$: 343.3, Found ($\text{M} + \text{H}$) $^+$: 343.2.



exp1	std1h
------	-------

date	AMPF	DEC.	& VI
Aug 30 2006	CD013	dfreq	259.909
solvent	dm	dn	H1
file	dpr	dpr	30
file /export/home/~	dof	dof	0
jpm/jbsjds/164-fid3	dm	dm	nmn
AcqISP8.fid	c	c	200
ACQUISITION	dmsf	dmsf	200
259.909	H1 dsesd	H1 dsesd	1.0
at	dres	dres	n
in	37612	37612	
mp	5.000	5.000	
sw	3761.2	3761.2	
nb	2100	2100	
nb	59	59	
tpwr	7.0	7.0	
dm	4.000	4.000	
tof	499.9	499.9	
nt	16	16	
ct	16	16	
atoclk	n	n	
gain	not used	not used	
flags	n	n	
11	n	n	
in	y	y	
dp	nm	nm	
hs			
DISPLAY	-71.9	-71.9	
sp	2362.4	2362.4	
wd	766	766	
ve	0	0	
sc	250	250	
wc	9.45	9.45	
hznm	402.45	402.45	
1s	2486.6	2486.6	
rfl	2171.3	2171.3	
rflp	20	20	
th	1.000	1.000	
ins			
nm			
ph			




NOESY of **2.87**:

js4-143bc01sp81 = spot 8 in CDC13
Pulse Sequence: NOESY
Solvent: CDC13
Ambient temperature
File: temp "nmr2"
UNITY-300 "nmr2"
Relax. delay 1.000 sec
Mixing 0.400 sec
Acq. time 0.212 sec
NUC1 13C
NUC2 1H
P1 2418.8 Hz
SFO 500.136 MHz
2 x 200 increments
OBSERVE H1, 299.307653 MHz
DATA PROCESSING
Gauss apodization 0.098 sec
F1 DATA PROCESSING
Gauss apodization 0.076 sec
F1 size 2048 x 2048
Total time 1 hr, 29 min, 36 sec



Synthesis of **2.215**




An oven-dried 2-dram vial equipped with a magnetic stir-bar was charged with 2.3 mg (0.0084 mmol) of bis(1,5-cyclooctadiene)nickel, 17.2 mg (0.0168 mmol) of **2.114**, and 0.84 mL of toluene-*d*₈ in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 42.3 mg (0.252 mmol) of **2.51** was added followed by 38.4 mg (0.168 mmol) of **2.62**. The vial was capped, sealed with electrical tape, and allowed to stir at ambient temperature for 19 h. After this time period, 36.9 μ L (0.252 mmol) of *N*-(trimethylsilyl)imidazole (**2.86**) was added and stirring was continued for an additional 1 h. Next, the mixture was filtered through a pad of celite using 10:1 hexanes:EtOAc, and volatile material was then removed under reduced pressure. Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 29.8 mg (0.0870 mmol, 52%) of **2.215**. In addition, 8.2 mg (0.024 mmol) of a 1:4.5 mixture of **2.87**:(*Z*)-**2.215** was isolated (66% combined yield). Following the reaction by ¹H NMR verified that (*E*)-**2.215** was initially formed, and that as the reaction progressed, (*E*)-**2.215** slowly isomerized to (*Z*)-**2.215**.¹


Trimethyl[(*S*,5*E*,7*E*)-4-phenyltrideca-1,5,7-trien-6-yloxy]silane (2.215**).** An oil. *R*_f = 0.17 (SiO₂, 70:1 hexanes:EtOAc); IR (neat): 3067 (m), 3029 (m), 2958 (s), 2920 (s), 2860 (s), 1658 (m), 1599 (m), 1497 (m), 1455 (m), 1366 (m), 1244 (s), 1189 (m), 1130

¹ Note that the *E* and *Z* descriptors used here represent the geometry of the silyl enol ether.

(m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.21-7.30 (2H, m), 7.10-7.21 (3H, m), 6.35 (1H, d, $J = 15$ Hz), 5.93 (1H, dt, $J = 15$ Hz, $J = 6.8$ Hz), 5.64-5.78 (1H, m), 5.01 (1H, d, $J = 18$ Hz), 4.96 (1H, d, $J = 10$ Hz), 4.88 (1H, d, $J = 10$ Hz), 3.58-3.68 (1H, m), 2.44-2.54 (1H, m), 2.30-2.42 (1H, m), 2.08 (2H, q, $J = 7.2$ Hz), 1.18-1.42 (6H, m), 0.86 (3H, t, $J = 6.4$ Hz), 0.15 (9H, s); ^{13}C NMR (CDCl_3): δ 147.5, 145.3, 136.6, 132.8, 128.3, 127.2, 125.9, 122.8, 116.1, 114.3, 42.47, 41.93, 32.57, 31.52, 28.95, 22.55, 14.08, 0.035. LRMS (ESI+) Calcd for $\text{C}_{22}\text{H}_{34}\text{OSi}$ ($\text{M} + \text{H}$) $^+$: 343.3, Found ($\text{M} + \text{H}$) $^+$: 343.3.




NOESY of **2.215**:



Deuterium labeling experiments:

Synthesis of Allylboronic Acid Pinacol Ester- d_2 (**2.51- d_2**)¹




To 0.665 g (4.32 mmol) of vinylboronic acid pinacol ester (**2.216**) and 1.00 g (5.61 mmol) of chloriodomethane- d_2 (**2.217**)² in 17 mL of THF at $-78\text{ }^{\circ}\text{C}$ was added 2.07 mL (5.2 mmol) of 2.5 M n -BuLi in hexane dropwise over 15 min. The reaction was then allowed to warm to room temperature and allowed to stir overnight. The reaction was then concentrated under reduced pressure. The residue was diluted with pentane and filtered through celite. Saturated aqueous NH_4Cl was added, and the two layers were filtered through celite. The organic layer was collected, and the aqueous layer extracted with pentane. The organic material was dried with anhydrous MgSO_4 and concentrated under reduced pressure. Purification of the mixture using silica gel chromatography (pentane/ CH_2Cl_2) gave 0.249 g (1.46 mmol, 34%) of **2.51- d_2** as a colorless oil that was contaminated with 8% of **2.216** as determined by GLC analysis.


¹ For lead reference, see: Sadhu, K. M.; Matteson, D. S. *Organometallics* **1985**, *4*, 1687.

² Purchased from Cambridge Isotopes or prepared from CD_2Cl_2 according to: Miyano, S.; Hashimoto, H. *Bull. Chem. Soc. Jpn.* **1971**, *44*, 2864.

Achiral GLC (Ultra 1, Hewlett-Packard, 35 °C) analysis of isolated **2.51-d₂**:



Non-Enantioselective Conjugate Allylation with **2.51-d₂** and Ni(cod)₂




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 3.0 mg (0.011 mmol) of bis(1,5-cyclooctadiene)nickel, 6.1 mg (0.022 mmol) of PCy₃, and 0.73 mL of THF-*d*₈ in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 53.7 μL (0.131 mmol) of a 2.44 M stock solution of **2.51-d₂** in toluene-*d*₈ was added, followed by 25.0 mg (0.109 mmol) of **2.62**. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at

ambient temperature for 12 h. After this time period, degassed water (N₂ sparge) was added, and the mixture transferred to a separatory funnel with CH₂Cl₂. After swirling the layers, the organic layer was collected, and the aqueous layer washed with CH₂Cl₂ (2x). The combined organic layers were dried with anhydrous Na₂SO₄, and volatile material was removed under reduced pressure. Analysis of the unpurified reaction mixture using achiral GLC was used to determine the chemoselectivity of the reaction. ¹H NMR analysis of the unpurified material verified that the label in unreacted **2.51-*d*₂** had not been scrambled. Silica gel chromatography of the mixture (hexanes/EtOAc) afforded 16.8 mg (0.0617 mmol, 57%) of **2.63-*d*₂**. The deuterium label ratios were determined by ²H NMR analysis of the purified material.


CCCCC1C(=O)C(=C)C(C1)C(=O)C=Cc2ccccc2 + CCCC1C(=O)C(=O)C(C1)C(=O)C=Cc2ccccc2

Non-Enantioselective Conjugate Allylation with **2.51-*d*₂** and Pd₂(dba)₃




An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.5 mg (0.0027 mmol) of bis(dibenzylideneacetone)dipalladium, 1.8 mg (0.0065 mmol) of PCy₃, and 0.72 mL of THF-*d*₈ in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 62.7 μL (0.153 mmol) of a 2.44 M stock solution of **2.51-*d*₂** in toluene-*d*₈ was added, followed by 25.0 mg (0.109 mmol) of **2.62**. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 24 h. After this time period, degassed water (N₂ sparge) was added, and the mixture transferred to a separatory funnel with CH₂Cl₂. After swirling the layers, the organic layer was collected, and the aqueous layer washed with CH₂Cl₂ (2x). The combined organic layers were dried with anhydrous Na₂SO₄, and volatile material was removed under reduced pressure. Analysis of the unpurified reaction mixture using achiral GLC was used to determine the chemoselectivity of the reaction. ¹H NMR analysis of the unpurified mixture verified that the label in unreacted **2.51-*d*₂** had not been scrambled. Silica gel chromatography of the mixture (hexanes/EtOAc) afforded 9.3 mg (0.034 mmol, 31%) of **2.63-*d*₂**. The deuterium label ratios were determined by ²H NMR analysis of the purified material.

338



CCCCC(=O)C(C)C(C)C(=O)C

Enantioselective Conjugate Allylation with **2.51-*d*₂**



An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 2.2 mg (0.0082 mmol) of bis(1,5-cyclooctadiene)nickel, 16.7 mg (0.0163 mmol) of **2.114**, and 0.23 mL of toluene-*d*₈ in a dry-box under an argon atmosphere. The vial was capped and allowed to stir for 45 min. Next, 100.0 μL (0.244 mmol) of a 2.44 M stock solution of **2.51-*d*₂** in toluene-*d*₈ was added, followed by 37.2 mg (0.163 mmol) of **2.62**. The vial was capped, taped with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 12 h. After this time period, degassed water (N_2 sparge) was added, and the mixture transferred to a separatory funnel with CH_2Cl_2 . After swirling the layers, the organic layer was collected, and the aqueous layer washed with CH_2Cl_2 (2x). The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Analysis of the unpurified reaction mixture using achiral GLC was used to determine the chemoselectivity of the reaction. ^1H NMR analysis of the unpurified mixture verified that the label in unreacted **2.51-*d*₂** had not been scrambled. Silica gel chromatography of the mixture (hexanes/EtOAc) afforded 9.1 mg (0.033 mmol, 20%) of **(S)-2.64-*d*₂**. The deuterium label ratios were determined by ^2H NMR analysis of the purified material.

SAMPLE DEC. & VT
date Sep 15 2007 dfreq 499.774
solvent CDCl₃ dn H₁
f16 / export/home/~dpuv 42
jpw/js6-47col-nmr-100t 42
ACQUISITION f16 dam yyy
dfreq 76.718 daf 7200
tn H2 dseq
at 3.000 dres 1.0
np 9208 homo n
sw 1534.7 DEC2 0
fb 1000 dfreq2 0
bs 4 dn2 1
cpwr 58 dpwr2 1
pw 11.0 dot2 0
dl 0 dm2 0
rt 0 dm2 0
ct 16 dm2 9900
n 16 dres2 1.0
gain 36 homo2 n
PROCESSING
FLAGS
f1 n wfile ft
in n proc ft
dp v tn 4036
hs nm math t
DISPLAY
sp -8.1 weff
wp 624.2 wepp
vs 14142 wds wft dssh dsst
sc 0
mcmm 250
is 250
rfl 64147.36
rfp 955.2
th 555.4
ins 22
ph 1.000
al

499.774
7.240
4.993
2.356


1.00
1.07

ppm

499.7 D →
Ph R

521. D

Synthesis of **2.98**




A 2-dram vial with magnetic stir-bar was charged with 2.2 mg (0.0082 mmol) of bis(1,5-cyclooctadiene)nickel and 40.0 mg (0.0815 mmol) of **2.96** in a dry-box under an argon atmosphere. Toluene (0.41 mL) was then added followed by 16.4 mg (0.0978 mmol) of allylboronic acid pinacol ester (**2.51**). The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 18 h. The reaction was quenched with the addition of 2 drops (18 gauge needle) of MeOH and subsequently, concentrated under reduced pressure. Chemoselectivity was determined by ¹H NMR analysis of the unpurified mixture. Purification by flash chromatography (SiO₂, hexanes/EtOAc) afforded 17.1 mg (0.0321 mmol) of **2.98** along with 9.3 mg (0.017 mmol) of oxidized material (**2.218**) (combined yield = 60%).

(E)-5-Oxo-7-phenyldeca-3,9-dienyl 2-(diphenylphosphino)benzoate (2.98). *R*_f = 0.17 (SiO₂, 6:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3400 (br w), 3067 (m), 2919 (m), 28528 (m), 1953 (w), 1716 (s), 1670 (s), 1632 (m), 1429 (s), 1265 (s), 1243 (s), 1138 (s), 1113 (s) cm⁻¹; ¹H NMR (CDCl₃): δ 7.95-8.02 (1H, m), 7.31-7.41 (2H, m), 7.26-7.31 (6H, m), 7.19-7.26 (6H, m), 7.10-7.18 (3H, m), 6.88-6.94 (1H, m), 6.63 (1H, dt, *J* = 16 Hz, *J* =

6.8 Hz), 6.01 (1H, d, $J = 16$ Hz), 5.61 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.88-4.99 (2H, m), 4.20 (2H, t, $J = 6.4$ Hz), 3.27 (1H, p, $J = 7.2$ Hz), 2.82 (2H, app d, $J = 7.6$ Hz), 2.40 (2H, q, $J = 6.8$ Hz), 2.36 (2H, t, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3): δ 198.6, 166.6, 144.1, 142.1, 140.4, 140.1, 137.8, 137.7, 136.1, 134.3, 133.9, 133.7, 132.3, 132.0, 130.6, 128.6, 128.5, 128.4, 128.3, 128.2, 127.5, 126.3, 63.03, 46.20, 40.96, 40.63, 31.53. ^{31}P NMR (CDCl_3): δ -3.10. LRMS (ESI+) Calcd for $\text{C}_{35}\text{H}_{33}\text{O}_3\text{P}$ ($\text{M} + \text{H}$) $^+$: 533.2, Found ($\text{M} + \text{H}$) $^+$: 533.1.

(*E*)-5-Oxo-7-phenyldeca-3,9-dienyl 2-(diphenylphosphoryl)benzoate (2.218). $R_f = 0.17$ (SiO_2 , 1:3 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3422 (br m), 3058 (m), 2928 (m), 2856 (m), 1966 (w), 1733 (s), 1661 (s), 1628 (m), 1433 (s), 1281 (s), 1256 (s), 1197 (s), 1121 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.80-7.88 (1H, m), 7.52-7.68 (5H, m), 7.30-7.52 (8H, m), 7.10-7.29 (5H, m), 6.51 (1H, dt, $J = 16$ Hz, $J = 6.8$ Hz), 5.89 (1H, d, $J = 16$ Hz), 5.61 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 4.88-4.98 (2H, m), 4.00 (2H, t, $J = 6.8$ Hz), 3.26 (1H, p, $J = 7.2$ Hz), 2.74-2.86 (2H, m), 2.35 (2H, t, $J = 7.2$ Hz), 2.20 (2H, q, $J = 6.8$ Hz); ^{31}P NMR (CDCl_3): δ 32.72. LRMS (ESI+) Calcd for $\text{C}_{35}\text{H}_{33}\text{O}_4\text{P}$ ($\text{M} + \text{H}$) $^+$: 549.2, Found ($\text{M} + \text{H}$) $^+$: 549.1.

Synthesis of **2.100**




An oven-dried vial with stir-bar was charged with 2.0 mg (0.0074 mmol) of bis(1,5-cyclooctadiene)nickel, 4.7 mg (0.015 mmol) of ligand **2.219**, and 0.98 mL of toluene in a dry-box under an argon atmosphere. To this solution was added 29.6 mg (0.176 mmol) of **2.51** and 45.0 mg (0.147 mmol) of **2.99**. The vial was sealed, removed from the dry-box, and allowed to stir at ambient temperature for 12 h. After this time period, degassed water (N_2 sparge) was added, and the mixture transferred to a separatory funnel with CH_2Cl_2 . The organic layer was collected, and the aqueous layer washed with CH_2Cl_2 (2x). The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Chemoselectivity was determined by ^1H NMR analysis of the unpurified mixture. Purification of the mixture using silica gel chromatography (hexanes/EtOAc) afforded 38.4 mg (0.110 mmol, 75%) of the conjugate allylation products **2.100** and **2.220**.

(E)-3-Allyl-5-oxo-7-phenylhept-6-enyl benzoate (2.100). An oil. $R_f = 0.23$ (SiO_2 , 6:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3067 (m), 2915 (m), 2852 (m), 1961 (w), 1716 (s), 1666 (s), 1610 (s), 1450 (m), 1272 (s), 1117 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 8.01 (2H, d, $J = 8$ Hz), 7.28-7.61 (9H, m), 6.72 (1H, d, $J = 16$ Hz), 5.79 (1H, ddt, $J = 17$ Hz, $J = 10$

Hz, $J = 7.2$ Hz), 5.00-5.12 (2H, m), 4.30-4.45 (2H, m), 2.71 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.66 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.37 (1H, h, $J = 6.4$ Hz), 2.09-2.29 (2H, m), 1.72-1.92 (2H, m); ^{13}C NMR (CDCl_3): δ 199.3, 166.5, 142.5, 135.8, 134.4, 132.8, 130.4, 130.2, 129.5, 128.8, 128.3, 128.2, 126.3, 117.3, 63.00, 44.88, 38.26, 32.58, 31.23. LRMS (ESI+) Calcd for $\text{C}_{23}\text{H}_{24}\text{O}_3$ ($\text{M} + \text{H}$) $^+$: 349.2, Found ($\text{M} + \text{H}$) $^+$: 349.1.

(E)-5-Oxo-7-phenyldeca-3,9-dienyl benzoate (2.220). An oil. $R_f = 0.17$ (SiO_2 , 6:1 hexanes:EtOAc); IR (CH_2Cl_2 solution): 3071 (m), 3025 (m), 2919 (m), 2851 (m), 1720 (s), 1673 (s), 1636 (s), 1454 (m), 1272 (s), 1104 (m) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.96-8.04 (2H, m), 7.52-7.58 (1H, m), 7.38-7.46 (2H, m), 7.18-7.27 (2H, m), 7.10-7.18 (3H, m), 6.74 (1H, dt, $J = 16$ Hz, $J = 6.8$ Hz), 6.12 (1H, dt, $J = 16$ Hz, $J = 2$ Hz), 5.61 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 6.8$ Hz), 4.86-5.00 (2H, m), 4.37 (2H, t, $J = 6.4$ Hz), 3.27 (1H, p, $J = 7.2$ Hz), 2.85 (1H, dd, $J = 17$ Hz, $J = 6.4$ Hz), 2.82 (1H, dd, $J = 17$ Hz, $J = 7.6$ Hz), 2.61 (2H, q, $J = 6.8$ Hz), 2.36 (2H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 198.6, 166.2, 144.0, 141.9, 136.1, 133.0, 132.4, 129.9, 129.5, 128.34, 128.32, 127.4, 126.3, 116.7, 62.71, 46.31, 41.00, 40.62, 31.80. LRMS (ESI+) Calcd for $\text{C}_{23}\text{H}_{24}\text{O}_3$ ($\text{M} + \text{H}$) $^+$: 349.2, Found ($\text{M} + \text{H}$) $^+$: 349.1.

Synthesis of Ni-enone Complex 2.101





An oven dried 20 mL vial with magnetic stir-bar was charged with 27.5 mg (0.0999 mmol) of bis(1,5-cyclooctadiene)nickel, 49.0 mg (0.0999 mmol) of **2.96**, and 1.0 mL of THF in a dry-box under an argon atmosphere. After stirring for 5 min, a deep purple solution formed, to which was then added 26.2 mg (0.0999 mmol) of triphenylphosphine. This mixture was allowed to stir for an additional 1 h providing a deep red solution. Volatile material was removed under reduced pressure in the dry-box. The residue was triturated with degassed pentane (3x) and dried under vacuum to afford 88.1 mg of a deep red solid.

Complex 2.101. Mp 152-172 °C (sealed capillary, decomp.). ^1H NMR (C_6D_6): δ 7.89 (1H, t, $J = 8.5$ Hz), 7.73 (1H, d, $J = 16$ Hz, H_f), 7.34-7.60 (8H, m), 7.33 (2H, t, $J = 8.5$ Hz), 6.78-7.11 (27H, m), 6.71 (1H, t, $J = 7.0$ Hz), 6.64 (1H, t, $J = 7.5$ Hz), 5.80 (1H, d, $J = 16$ Hz, H_e), 4.66 (1H, br t, H_a), 3.80 (1H, m, H_c), 3.71 (1H, m, H_d), 3.50 (1H, d, $J = 11$ Hz, H_a'), 1.15-1.32 (2H, m, H_b); ^{13}C NMR (C_6D_6), diagnostic peaks: δ 187.8 (C_g), 168.1, 137.21 (C_f), 130.65 (C_e), 71.15 (C_d , d, $^2J_{\text{CP}} = 11$ Hz), 62.52 (C_a), 54.23 (C_c , d, $^2J_{\text{CP}} = 20$ Hz), 27.98 (C_b). Note that the aromatic region was too complex for further assignment. ^{31}P NMR (C_6D_6): δ 41.21 (d, $^2J_{\text{PP}} = 32$ Hz), 28.59 (d, $^2J_{\text{PP}} = 32$ Hz).

js5-86 in c606

exp5 szput1

SAMPLE DEC. & VT 499.774
 date Oct 22 2007 /dfrq
 solvent Benzene /dn
 title /export/home/~dpw
 jsm/js5-86.tld /dot
 ACQUISITION
 stfrq 499.774
 tn 499.774
 at 5.000
 np 4317.6
 sw 4317.6
 fb 2000
 bs 4
 tpwr 57
 pw 4.6
 dt 0
 tof 0
 nt -322.2
 ct 8
 atlock 8
 gain not used
 flos not used
 i1 n
 in n
 dp n
 hs n
 DISPLAY 5.4
 sp 3999.6
 wp 317
 vs wepp
 sc 0
 wc wnt
 hzmm 15.00
 ts 448.74
 fti 3596.4
 ffp 3573.4
 fns 1.000
 at ph




```

js6-97
exp5 sepul

SAMPLE
date Oct 26 2007
time 14:00
jsm/js6-97-13C-bb-~

ACQUISITION
sfreq 125.680
in C13
at 3.000
np 200168
sw 33361.1
f2 18000
bs 4
tpwr 55
pw 7.0
dl 2.000
tot 816.6
nt 4520
atock 4520
gain not used
flask not used
f1 n
f2 n
f3 n
f4 n
f5 n
f6 n
f7 n
f8 n
f9 n
f10 n
f11 n
f12 n
f13 n
f14 n
f15 n
f16 n
f17 n
f18 n
f19 n
f20 n
f21 n
f22 n
f23 n
f24 n
f25 n
f26 n
f27 n
f28 n
f29 n
f30 n
f31 n
f32 n
f33 n
f34 n
f35 n
f36 n
f37 n
f38 n
f39 n
f40 n
f41 n
f42 n
f43 n
f44 n
f45 n
f46 n
f47 n
f48 n
f49 n
f50 n
f51 n
f52 n
f53 n
f54 n
f55 n
f56 n
f57 n
f58 n
f59 n
f60 n
f61 n
f62 n
f63 n
f64 n
f65 n
f66 n
f67 n
f68 n
f69 n
f70 n
f71 n
f72 n
f73 n
f74 n
f75 n
f76 n
f77 n
f78 n
f79 n
f80 n
f81 n
f82 n
f83 n
f84 n
f85 n
f86 n
f87 n
f88 n
f89 n
f90 n
f91 n
f92 n
f93 n
f94 n
f95 n
f96 n
f97 n
f98 n
f99 n
f100 n
f101 n
f102 n
f103 n
f104 n
f105 n
f106 n
f107 n
f108 n
f109 n
f110 n
f111 n
f112 n
f113 n
f114 n
f115 n
f116 n
f117 n
f118 n
f119 n
f120 n
f121 n
f122 n
f123 n
f124 n
f125 n
f126 n
f127 n
f128 n
f129 n
f130 n
f131 n
f132 n
f133 n
f134 n
f135 n
f136 n
f137 n
f138 n
f139 n
f140 n
f141 n
f142 n
f143 n
f144 n
f145 n
f146 n
f147 n
f148 n
f149 n
f150 n
f151 n
f152 n
f153 n
f154 n
f155 n
f156 n
f157 n
f158 n
f159 n
f160 n
f161 n
f162 n
f163 n
f164 n
f165 n
f166 n
f167 n
f168 n
f169 n
f170 n
f171 n
f172 n
f173 n
f174 n
f175 n
f176 n
f177 n
f178 n
f179 n
f180 n
f181 n
f182 n
f183 n
f184 n
f185 n
f186 n
f187 n
f188 n
f189 n
f190 n
f191 n
f192 n
f193 n
f194 n
f195 n
f196 n
f197 n
f198 n
f199 n
f200 n
f201 n
f202 n
f203 n
f204 n
f205 n
f206 n
f207 n
f208 n
f209 n
f210 n
f211 n
f212 n
f213 n
f214 n
f215 n
f216 n
f217 n
f218 n
f219 n
f220 n
f221 n
f222 n
f223 n
f224 n
f225 n
f226 n
f227 n
f228 n
f229 n
f230 n
f231 n
f232 n
f233 n
f234 n
f235 n
f236 n
f237 n
f238 n
f239 n
f240 n
f241 n
f242 n
f243 n
f244 n
f245 n
f246 n
f247 n
f248 n
f249 n
f250 n
f251 n
f252 n
f253 n
f254 n
f255 n
f256 n
f257 n
f258 n
f259 n
f260 n
f261 n
f262 n
f263 n
f264 n
f265 n
f266 n
f267 n
f268 n
f269 n
f270 n
f271 n
f272 n
f273 n
f274 n
f275 n
f276 n
f277 n
f278 n
f279 n
f280 n
f281 n
f282 n
f283 n
f284 n
f285 n
f286 n
f287 n
f288 n
f289 n
f290 n
f291 n
f292 n
f293 n
f294 n
f295 n
f296 n
f297 n
f298 n
f299 n
f300 n
f301 n
f302 n
f303 n
f304 n
f305 n
f306 n
f307 n
f308 n
f309 n
f310 n
f311 n
f312 n
f313 n
f314 n
f315 n
f316 n
f317 n
f318 n
f319 n
f320 n
f321 n
f322 n
f323 n
f324 n
f325 n
f326 n
f327 n
f328 n
f329 n
f330 n
f331 n
f332 n
f333 n
f334 n
f335 n
f336 n
f337 n
f338 n
f339 n
f340 n
f341 n
f342 n
f343 n
f344 n
f345 n
f346 n
f347 n
f348 n
f349 n
f350 n
f351 n
f352 n
f353 n
f354 n
f355 n
f356 n
f357 n
f358 n
f359 n
f360 n
f361 n
f362 n
f363 n
f364 n
f365 n
f366 n
f367 n
f368 n
f369 n
f370 n
f371 n
f372 n
f373 n
f374 n
f375 n
f376 n
f377 n
f378 n
f379 n
f380 n
f381 n
f382 n
f383 n
f384 n
f385 n
f386 n
f387 n
f388 n
f389 n
f390 n
f391 n
f392 n
f393 n
f394 n
f395 n
f396 n
f397 n
f398 n
f399 n
f400 n
f401 n
f402 n
f403 n
f404 n
f405 n
f406 n
f407 n
f408 n
f409 n
f410 n
f411 n
f412 n
f413 n
f414 n
f415 n
f416 n
f417 n
f418 n
f419 n
f420 n
f421 n
f422 n
f423 n
f424 n
f425 n
f426 n
f427 n
f428 n
f429 n
f430 n
f431 n
f432 n
f433 n
f434 n
f435 n
f436 n
f437 n
f438 n
f439 n
f440 n
f441 n
f442 n
f443 n
f444 n
f445 n
f446 n
f447 n
f448 n
f449 n
f450 n
f451 n
f452 n
f453 n
f454 n
f455 n
f456 n
f457 n
f458 n
f459 n
f460 n
f461 n
f462 n
f463 n
f464 n
f465 n
f466 n
f467 n
f468 n
f469 n
f470 n
f471 n
f472 n
f473 n
f474 n
f475 n
f476 n
f477 n
f478 n
f479 n
f480 n
f481 n
f482 n
f483 n
f484 n
f485 n
f486 n
f487 n
f488 n
f489 n
f490 n
f491 n
f492 n
f493 n
f494 n
f495 n
f496 n
f497 n
f498 n
f499 n
f500 n
f501 n
f502 n
f503 n
f504 n
f505 n
f506 n
f507 n
f508 n
f509 n
f510 n
f511 n
f512 n
f513 n
f514 n
f515 n
f516 n
f517 n
f518 n
f519 n
f520 n
f521 n
f522 n
f523 n
f524 n
f525 n
f526 n
f527 n
f528 n
f529 n
f530 n
f531 n
f532 n
f533 n
f534 n
f535 n
f536 n
f537 n
f538 n
f539 n
f540 n
f541 n
f542 n
f543 n
f544 n
f545 n
f546 n
f547 n
f548 n
f549 n
f550 n
f551 n
f552 n
f553 n
f554 n
f555 n
f556 n
f557 n
f558 n
f559 n
f560 n
f561 n
f562 n
f563 n
f564 n
f565 n
f566 n
f567 n
f568 n
f569 n
f570 n
f571 n
f572 n
f573 n
f574 n
f575 n
f576 n
f577 n
f578 n
f579 n
f580 n
f581 n
f582 n
f583 n
f584 n
f585 n
f586 n
f587 n
f588 n
f589 n
f590 n
f591 n
f592 n
f593 n
f594 n
f595 n
f596 n
f597 n
f598 n
f599 n
f600 n
f601 n
f602 n
f603 n
f604 n
f605 n
f606 n
f607 n
f608 n
f609 n
f610 n
f611 n
f612 n
f613 n
f614 n
f615 n
f616 n
f617 n
f618 n
f619 n
f620 n
f621 n
f622 n
f623 n
f624 n
f625 n
f626 n
f627 n
f628 n
f629 n
f630 n
f631 n
f632 n
f633 n
f634 n
f635 n
f636 n
f637 n
f638 n
f639 n
f640 n
f641 n
f642 n
f643 n
f644 n
f645 n
f646 n
f647 n
f648 n
f649 n
f650 n
f651 n
f652 n
f653 n
f654 n
f655 n
f656 n
f657 n
f658 n
f659 n
f660 n
f661 n
f662 n
f663 n
f664 n
f665 n
f666 n
f667 n
f668 n
f669 n
f670 n
f671 n
f672 n
f673 n
f674 n
f675 n
f676 n
f677 n
f678 n
f679 n
f680 n
f681 n
f682 n
f683 n
f684 n
f685 n
f686 n
f687 n
f688 n
f689 n
f690 n
f691 n
f692 n
f693 n
f694 n
f695 n
f696 n
f697 n
f698 n
f699 n
f700 n
f701 n
f702 n
f703 n
f704 n
f705 n
f706 n
f707 n
f708 n
f709 n
f710 n
f711 n
f712 n
f713 n
f714 n
f715 n
f716 n
f717 n
f718 n
f719 n
f720 n
f721 n
f722 n
f723 n
f724 n
f725 n
f726 n
f727 n
f728 n
f729 n
f730 n
f731 n
f732 n
f733 n
f734 n
f735 n
f736 n
f737 n
f738 n
f739 n
f740 n
f741 n
f742 n
f743 n
f744 n
f745 n
f746 n
f747 n
f748 n
f749 n
f750 n
f751 n
f752 n
f753 n
f754 n
f755 n
f756 n
f757 n
f758 n
f759 n
f760 n
f761 n
f762 n
f763 n
f764 n
f765 n
f766 n
f767 n
f768 n
f769 n
f770 n
f771 n
f772 n
f773 n
f774 n
f775 n
f776 n
f777 n
f778 n
f779 n
f780 n
f781 n
f782 n
f783 n
f784 n
f785 n
f786 n
f787 n
f788 n
f789 n
f790 n
f791 n
f792 n
f793 n
f794 n
f795 n
f796 n
f797 n
f798 n
f799 n
f800 n
f801 n
f802 n
f803 n
f804 n
f805 n
f806 n
f807 n
f808 n
f809 n
f810 n
f811 n
f812 n
f813 n
f814 n
f815 n
f816 n
f817 n
f818 n
f819 n
f820 n
f821 n
f822 n
f823 n
f824 n
f825 n
f826 n
f827 n
f828 n
f829 n
f830 n
f831 n
f832 n
f833 n
f834 n
f835 n
f836 n
f837 n
f838 n
f839 n
f840 n
f841 n
f842 n
f843 n
f844 n
f845 n
f846 n
f847 n
f848 n
f849 n
f850 n
f851 n
f852 n
f853 n
f854 n
f855 n
f856 n
f857 n
f858 n
f859 n
f860 n
f861 n
f862 n
f863 n
f864 n
f865 n
f866 n
f867 n
f868 n
f869 n
f870 n
f871 n
f872 n
f873 n
f874 n
f875 n
f876 n
f877 n
f878 n
f879 n
f880 n
f881 n
f882 n
f883 n
f884 n
f885 n
f886 n
f887 n
f888 n
f889 n
f890 n
f891 n
f892 n
f893 n
f894 n
f895 n
f896 n
f897 n
f898 n
f899 n
f900 n
f901 n
f902 n
f903 n
f904 n
f905 n
f906 n
f907 n
f908 n
f909 n
f910 n
f911 n
f912 n
f913 n
f914 n
f915 n
f916 n
f917 n
f918 n
f919 n
f920 n
f921 n
f922 n
f923 n
f924 n
f925 n
f926 n
f927 n
f928 n
f929 n
f930 n
f931 n
f932 n
f933 n
f934 n
f935 n
f936 n
f937 n
f938 n
f939 n
f940 n
f941 n
f942 n
f943 n
f944 n
f945 n
f946 n
f947 n
f948 n
f949 n
f950 n
f951 n
f952 n
f953 n
f954 n
f955 n
f956 n
f957 n
f958 n
f959 n
f960 n
f961 n
f962 n
f963 n
f964 n
f965 n
f966 n
f967 n
f968 n
f969 n
f970 n
f971 n
f972 n
f973 n
f974 n
f975 n
f976 n
f977 n
f978 n
f979 n
f980 n
f981 n
f982 n
f983 n
f984 n
f985 n
f986 n
f987 n
f988 n
f989 n
f990 n
f991 n
f992 n
f993 n
f994 n
f995 n
f996 n
f997 n
f998 n
f999 n
f1000 n



```



COSY of **2.101** in C₆D₆:



js6-06-gCOSY in C6D6
Pulse Sequence: gCOSY
Solvent: Benzene
Ambient temperature
File: js6-06gCOSY
INNOVA-500 "nmr3"

Relax. delay 1.000 sec
Acq. time 0.237 sec
Width 4317.6 Hz
2D Width 4317.6 Hz
16 repetitions
512 increments
OBSERVE H1, 499.7714819 MHz
DATA PROCESSING
Sf. time bell 0.119 sec
F1 DATA PROCESSING
Sf. time bell 0.030 sec
F1 size 2182 x 0192
Total time 2 hr, 59 min, 6 sec





js6-06-gCOSY in C6D6
Pulse Sequence: gCOSY
Solvent: Benzene
Ambient temperature
INNOVA-500 "nmr3"

Relax. delay 1.000 sec
Acq. time 0.237 sec
Width 4317.6 Hz
2D Width 4317.6 Hz
16 repetitions
512 increments
OBSERVE H1, 499.7714819 MHz
DATA PROCESSING
Sf. time bell 0.119 sec
F1 DATA PROCESSING
Sf. time bell 0.030 sec
F1 size 2182 x 0192
Total time 2 hr, 59 min, 6 sec





HSQC of **2.101** in C₆D₆:

jse-97
 Pulse Sequence: ghsqc
 Solvent: Benzene
 Ambient temperature
 User: j-14-97
 File: jse-97ghsdc-rp
 INOVA-500 "nars"
 Relax. delay 1.000 sec
 Acq. time 0.100 sec
 Width 4197.8 Hz
 2D Width 22707.9 Hz
 64 repetitions
 2 x 256 increments
 OBSERVE H1, 499.7714051 MHz
 DECOUPLE C13, 125.6009937 MHz
 Power 44 dB
 on during acquisition
 off during delay
 GARP-1 modulated
 DATA PROCESSING
 Gauss apodization 0.003 sec
 F1 DATA PROCESSING
 Gauss apodization 0.011 sec
 F1 size 6192 x 4096
 Total time 11 hr, 25 min, 47 sec





jse-97
 Pulse Sequence: ghsqc
 Solvent: Benzene
 Ambient temperature
 User: j-14-97
 File: jse-97ghsdc-rp
 INOVA-500 "nars"
 Relax. delay 1.000 sec
 Acq. time 0.100 sec
 Width 4197.8 Hz
 2D Width 22707.9 Hz
 64 repetitions
 2 x 256 increments
 OBSERVE H1, 499.7714051 MHz
 DECOUPLE C13, 125.6009937 MHz
 Power 44 dB
 on during acquisition
 off during delay
 GARP-1 modulated
 DATA PROCESSING
 Gauss apodization 0.003 sec
 F1 DATA PROCESSING
 Gauss apodization 0.011 sec
 F1 size 6192 x 4096
 Total time 11 hr, 25 min, 47 sec





NOESY of **2.101** in C₆D₆:

j96-127
Pulse Sequence: NOESY
Solvent: Benzene
Ambient temperature
INOVA-500 "nuc3"
Relax. delay 1.000 sec
Mixing 0.500 sec
Acq. time 0.230 sec
Width 4455.4 Hz
ZD Width 4455.4 Hz
16 repetitions
2 x 256 increments
OBSERVE H1, 499.7714022 MHz
DATA PROCESSING
Gauss apodization 0.100 sec
F1 DATA PROCESSING
Gauss apodization 0.041 sec
FT size 2048 x 2048
Total time 4 hr, 2 min, 53 sec



j96-127
Pulse Sequence: NOESY
Solvent: Benzene
Ambient temperature
File: j96-127-noesy
INOVA-500 "nuc3"
Relax. delay 1.000 sec
Mixing 0.500 sec
Acq. time 0.230 sec
Width 4455.4 Hz
ZD Width 4455.4 Hz
16 repetitions
2 x 256 increments
OBSERVE H1, 499.7714022 MHz
DATA PROCESSING
Gauss apodization 0.100 sec
F1 DATA PROCESSING
Gauss apodization 0.041 sec
FT size 4096 x 4096
Total time 4 hr, 2 min, 53 sec



$$\text{dba} + \text{Ni(cod)}_2 + \text{PCy}_3 \xrightarrow{\text{d}_8\text{-THF}} \text{NMR at } -27^\circ\text{C}$$

d_8 -THF

NMR at -27 °C

5.500


—COD

3.580

2.336

—COD


1.730



js4-200NPD-Nicod2 + PCy3 + dba at -27C
exp4 s2pu1



SAMPLE DEC. 8 VT
date Sep 30 2006 dfrq 489.785
solvent thf dn H1
file /export/home/~ dpr 39
js4-200NPD-26~ dot 0
-6-31P.fid dm yvy
ACQUISITION
sfrq 202.312 dmf 11905
in P31 dseq 1.0
at 2.000 homo
np 279966 temp -30.0
sw 6931.5
lb 39000 dfrq2 DEC2 0
4 56 dn2
tpwf 15.0 dpr2 1
np 1.000 dorf2 0
dl 1500.0 dm2 n
tof 128 dm2 c
ct 48 dm2 9900
a1ock n dseq2 1.0
gain not used
11 n PROCESSING 2.00
in n lb file
dp y wfile
ns ny prod ft
ns DISPLAY ny fn not used
f
SP -38987.8 math
VE 69991.0
VC 46 weff
WC 250 wepp
h2mm 279.96 vds
15 500.00 wnt
rf1 46463.6
rfp 7475.8
t1n 100.000
nm cdc ph

dba + Ni(cod)₂ + PCy₃ $\xrightarrow{\text{d}_8\text{-THF}}$ NMR at -27 °C



SpinWorks 2.5: js4-57, Ni + PCy₃ + dba, after 30 min

46.1084
38.1908
11.2343




file: D:\sieber\2\js4-57\21\fid exp: <gpg30>
transmitter freq: 161.946247 MHz
time domain size: 65536 points
width: 51282.05 Hz = 316.660943 ppm = 0.782502 Hz/pt
number of scans: 80

freq. of 0 ppm: 161.942715 MHz
processed size: 32768 complex points
LB: 1.000 GB: 0.0000
Hz/cm: 2051.282 ppm/cm: 12.66644

Non-linear effects (Figures 2.6 and 2.7):

Effect of Ligand Enantiomeric Purity on Enantioselectivity



An oven-dried 2-dram vial, equipped with a magnetic stir-bar, was charged with 3.0 mg (0.011 mmol) of bis(1,5-cyclooctadiene)nickel. The two enantiopodes of ligand **2.114** were then added as 0.100 M stock solutions in toluene, using gas-tight syringes, to make the desired enantiomeric purity of ligand for the reaction (0, 25, 50, and 75 % ee were examined). Next, 221 μL of toluene was added, and this mixture was allowed to stir for 45 min. Next, 44.2 mg (0.263 mmol) of **2.51** was added, followed by 50.0 mg (0.219 mmol) of **2.62**. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 20 h. After this time period, degassed water (N_2 sparge) was added, and the mixture transferred to a separatory funnel with CH_2Cl_2 . After swirling the layers, the organic layer was collected, and the aqueous layer washed with CH_2Cl_2 (2x). The combined organic layers were dried with anhydrous Na_2SO_4 , and volatile material was removed under reduced pressure. Analysis of the unpurified mixture using GLC was used to determine the chemoselectivity of the reaction. Purification using silica gel chromatography (hexanes/ EtOAc) afforded the


conjugate allylation product. Enantioselectivity was determined by SFC analysis of the purified material.

Effect of Ligand Enantiomeric Purity on Conversion

Enantiomerically pure and racemic catalyst stock solutions, consisting of a mixture of Ni(cod)₂ and ligand **2.114**, were prepared using 3.4 mg of Ni(cod)₂ and 24.5 mg of enantiomerically pure or racemic ligand **2.114** in 240 μ L of toluene-*d*₈, respectively. The catalyst stock solutions were then allowed to stand at room temperature for 45 min before use. To two oven-dried J-Young NMR tubes, in a dry-box, was added 350 μ L of a 0.750 M stock solution of **2.51** in toluene-*d*₈ to each tube. To one tube was added 175 μ L of the enantiomerically pure catalyst stock solution, and to the other tube was added 175 μ L of the racemic catalyst solution. Finally, 350 μ L of a 0.500 M stock solution of **2.62** in toluene-*d*₈ was added to each tube, and each tube was capped and inverted several times. The reactions were followed by ¹H NMR, and the ratio of starting material to product¹ was used to calculate conversion. After complete consumption of starting material, the reactions were worked up as described for the general procedure for the Ni-catalyzed asymmetric conjugate allylation of **2.62** on page 257.


¹ Note that the product observed during the reaction was the expected intermediate boron enolate.

Sample ^1H NMR spectrum for conversion calculation:




6. Synthetic Transformations

Synthesis of **2.81**



To 47.9 mg (0.173 mmol) of **2.52** was added 2.2 mg (0.0035 mmol) of the Hoveyda-Grubbs' 2nd generation catalyst (**2.221**) in a dry-box under an argon atmosphere. This mixture was then diluted with 5.7 mL of dry, degassed CH_2Cl_2 , and a magnetic stir-bar was added. A septum was fitted on the reaction flask, and the reaction was removed from the dry-box and allowed to stir at room temperature under N_2 . After 1 h, 3 drops (22 gauge needle) of *t*-butyl vinyl ether was added, and volatile material was removed under reduced pressure. Silica gel chromatography (pentane/ Et_2O) of the mixture afforded 28.1 mg (0.163 mmol, 94%) of 5-phenylcyclohex-2-en-1-one (**2.81**). Spectral data were consistent with the literature.¹

Synthesis of **2.79**




¹ Rutherford, A. P.; Gibb, C. S.; Hartley, R. C.; Goodman, J. M. *J. Chem. Soc. Perkin Trans. 1* **2001**, 1051.

In a 2-dram vial with magnetic stir-bar, in a dry-box under Ar, was weighed ~30 mg of crushed 4 Å molecular sieves. Next, 33.8 µL (0.034 mmol) of a 1.0 M solution of (±)-*trans*-1,2-diaminocyclohexane (**2.78**) in CH₂Cl₂ was added by syringe, followed by dilution with 0.32 mL of CH₂Cl₂. Next, 33.8 µL (0.034 mmol) of a 1.0 M solution of SnCl₄ in CH₂Cl₂ was added, and the vial was capped with a septum, removed from the dry-box, and cooled to 0 °C (ice/brine). TMS₂O₂ (**2.77**) was added dropwise as a 1.0 M solution in CH₂Cl₂ (0.27 mL, 0.27 mmol). The mixture was allowed to stir for 10 min at this temperature, and then, 37.3 mg (0.135 mmol) of **2.52** was added in 0.59 mL of CH₂Cl₂ by canula. The reaction became a blue-gray color and was subsequently warmed to room temperature and allowed to stir for 15 h. Sodium sulfite (41 mg) was then added, and the reaction was allowed to stir for an additional 3 h. Finally, the reaction was filtered through a pad of silica gel using EtOAc and concentrated under reduced pressure. Silica gel chromatography (hexanes/EtOAc) of the mixture afforded 31.5 mg (0.108 mmol, 80 %) of **2.79** as a white solid.

(E)-Styryl-3-phenylhex-5-enoate (2.79). Mp = 66-70°C. R_f = 0.24 (30:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 3087 (m), 3024 (m), 2917 (m), 1747 (s), 1652 (m), 1495 (m), 1212 (m), 1142 (s) cm⁻¹; ¹H NMR: δ 7.74 (1H, d, *J* = 13 Hz), 7.24-7.39 (6H, m), 7.10-7.24 (4H, m), 6.03 (1H, d, *J* = 13 Hz), 5.66 (1H, m), 5.02 (2H, m), 3.26 (1H, p, *J* = 7.2 Hz), 2.81 (1H, dd, *J* = 16 Hz, *J* = 6.4 Hz), 2.68 (1H, dd, *J* = 16 Hz, *J* = 8.4 Hz), 2.42 (2H, m); ¹³C NMR: δ 169.3, 143.1, 136.1, 135.6, 134.0, 128.6, 128.5, 127.3,

126.7, 126.1, 117.1, 115.2, 41.56, 40.61, 40.26. LRMS (ESI+) Calcd for C₂₀H₂₀O₂ (M + Na)⁺: 315, Found (M + Na)⁺: 315.


Synthesis of **2.80**



To a solution of 26.3 mg (0.900 mmol) of **2.79** in 0.68 mL of THF was added 0.22 mL of water, and the reaction was subsequently cooled to 0 °C. LiOH·H₂O (7.6 mg, 0.18 mmol) was then added, and the reaction was allowed to stir at this temperature and monitored by TLC. After complete consumption of the starting material (2-2.5 h), the reaction was acidified with 1 M HCl and extracted with EtOAc (3x). The organic layers were combined, washed with brine, and dried over anhydrous Na₂SO₄. Volatile material was removed under reduced pressure, and the product was purified by silica gel chromatography (1% AcOH in CH₂Cl₂/Et₂O, R_f = 0.25 in 1% AcOH in 20:1 CH₂Cl₂:Et₂O) to give 16.5 mg (0.867 mmol, 96%) of **2.80** after removal of AcOH by azeotropic distillation with toluene using a rotary evaporator followed by removal of toluene via azeotropic distillation with CH₂Cl₂. Spectral data were consistent with the literature.²

² Allin, S. M.; Essat, M.; Pita, C. H.; Baird, R. D.; McKee, V.; Elsegood, M.; Edgar, M.; Andrews, D. M.; Shah, P.; Aspinall, I. *Org. Biomol. Chem.* **2005**, 3, 809.

Synthesis of **2.141**




In a 2-dram vial with magnetic stir-bar, in a dry-box under Ar, was weighed ~45 mg of crushed 4 Å molecular sieves. Next, 48.8 μL (0.049 mmol) of a 1.0 M solution of (\pm)-*trans*-1,2-diaminocyclohexane (**2.78**) in CH_2Cl_2 was added by syringe followed by dilution with 0.43 mL of CH_2Cl_2 . Next, 48.8 μL (0.049 mmol) of a 1.0 M solution of SnCl_4 in CH_2Cl_2 was added, and the vial was capped with a septum, removed from the dry-box, and cooled to 0 °C (ice/brine). TMS_2O_2 (**2.77**) was added dropwise as a 1.0 M solution in CH_2Cl_2 (0.39 mL, 0.39 mmol). The mixture was allowed to stir for 10 min at this temperature, and then, 52.8 mg (0.195 mmol) of (*S*)-**2.64** was added in 0.85 mL of CH_2Cl_2 by canula. The reaction was subsequently warmed to room temperature and allowed to stir for 15 h. Sodium sulfite (60 mg) was then added, and the reaction was allowed to stir for an additional 3 h. Finally, the mixture was filtered through a pad of silica gel using EtOAc and concentrated under reduced pressure. Silica gel chromatography (hexanes/EtOAc) of the mixture afforded 37.8 mg (0.132 mmol, 68 %, 92% brsm) of **2.141** as a colorless oil along with 13.8 mg (0.0510 mmol) of unreacted (*S*)-**2.64**.

(*S,E*)-Hept-1-enyl-3-phenylhex-5-enoate (2.141). R_f = 0.30 (30:1 hexanes:EtOAc); IR (neat): 3080 (m), 3029 (m), 2958 (s), 2928 (s), 2856 (s), 1945 (w), 1750 (s), 1675 (m),


1447 (m), 1236 (m), 1160 (s) cm^{-1} ; ^1H NMR (CDCl_3): δ 7.25-7.32 (2H, m), 7.15-7.23 (3H, m), 6.95 (1H, dt, $J = 12$ Hz, $J = 2$ Hz), 5.64 (1H, ddt, $J = 17$ Hz, $J = 10$ Hz, $J = 7.2$ Hz), 5.33 (1H, dt, $J = 12$ Hz, $J = 7.2$ Hz), 4.91-5.05 (2H, m), 3.22 (1H, p, $J = 7.6$ Hz), 2.74 (1H, dd, $J = 16$ Hz, $J = 6.8$ Hz), 2.61 (1H, dd, $J = 16$ Hz, $J = 8.4$ Hz), 2.32-2.48 (2H, m), 1.93 (2H, q, $J = 6.8$ Hz), 1.17-1.39 (6H, m), 0.87 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (CDCl_3): δ 169.5, 143.3, 135.7, 135.2, 128.4, 127.3, 126.6, 117.0, 115.1, 41.55, 40.55, 40.28, 31.24, 27.16, 27.21, 22.45, 14.02. LRMS (ESI+) Calcd for $\text{C}_{19}\text{H}_{26}\text{O}_2$ ($\text{M} + \text{H}$) $^+$: 287.2, Found ($\text{M} + \text{H}$) $^+$: 287.1. $[\alpha]_{\text{D}}^{20} = +15^\circ$ ($c = 3.0$, CHCl_3).

Synthesis of (*S*)-**2.80**



To a solution of 27.4 mg (0.0957 mmol) of **2.141** in 0.96 mL of a 3:1 THF:H₂O mixture at 0 °C was added LiOH·H₂O (8.0 mg, 0.19 mmol). The reaction was allowed to reach ambient temperature and allowed to stir for 20h. The reaction was acidified with 1 M HCl and extracted with EtOAc (3x). The organic layers were combined, washed with brine, and dried over anhydrous Na₂SO₄. Volatile material was removed under reduced pressure, and the product was purified using silica gel chromatography (1% AcOH in CH₂Cl₂/Et₂O, $R_f = 0.25$ in 1% AcOH in 20:1 CH₂Cl₂:Et₂O) to give 17.9 mg (0.0941 mmol, 98%) of (*S*)-**2.80** after removal of AcOH by azeotropic distillation with toluene using a rotary evaporator, followed by removal of toluene via azeotropic distillation with CH₂Cl₂.

Synthesis of **2.142**





To 237 mg (0.759 mmol) of **2.137** in a 20 mL scintillation vial with magnetic stir-bar, in a dry-box under an Ar atmosphere, was added a solution of 9.8 mg (0.0157 mmol) of the Hoveyda-Grubbs' second generation catalyst (**2.221**) in 16 mL of CH₂Cl₂. The vial was capped, sealed with electrical tape, removed from the dry-box, and allowed to stir at ambient temperature for 1 h. Next, 0.15 mL of *t*-butyl vinyl ether was added, and the mixture was allowed to stir for another 30 min. The reaction was concentrated under reduced pressure and purified using column chromatography (SiO₂, pentane:Et₂O) to afford 152 mg (0.743 mmol, 98%) of **2.142** as a colorless oil.

(R)-5-(5-Fluoro-2-methylphenyl)cyclohex-2-en-1-one (2.142). R_f = 0.14 (SiO₂, 7:1 pentane:Et₂O); IR (neat): 3031 (m), 2930 (m), 1879 (w), 1678 (s), 1615 (m), 1584 (m), 1495 (s), 1388 (s), 1243 (s), 1161 (m) cm⁻¹; ¹H NMR (CDCl₃): δ 7.09 (1H, dd, J = 8.4 Hz, J = 6.0 Hz), 6.99-7.07 (1H, m), 6.91 (1H, dd, J = 10 Hz, J = 2.8 Hz), 6.81 (1H, dt, J = 8.0 Hz, J = 2.8 Hz), 6.01 (1H, dd, J = 11 Hz, J = 2.8 Hz), 3.45-3.58 (1H, m), 2.38-2.62 (4H, m); ¹³C NMR (CDCl₃): δ 198.6, 161.4 (d, $^1J_{CF}$ = 242 Hz), 149.2, 143.0 (d, $^3J_{CF}$ = 6.2 Hz), 131.9 (d, $^3J_{CF}$ = 7.8 Hz), 130.6, 129.6, 113.2 (d, $^2J_{CF}$ = 21 Hz), 112.3 (d, $^2J_{CF}$ = 21 Hz), 44.16, 36.85, 32.48, 18.50; ¹⁹F NMR: δ 95.10 (m). LRMS (APPI) Calcd for C₁₃H₁₃FO (M + H)⁺: 205.1, Found (M + H)⁺: 205.1.


jss-155coium

exp1 std1h

SAMPLE		DEC. & VT	
date	Mar 27 2007	dfrq	0
solvent	CDCl3	dn	30
file	ACQUISITION	exp	0
stfrq	400.029	dot	mm
th	400	dm	200
at	3.000	dat	PROCESSING
mp	35992	vtfile	ft
sw	5398.8	proc	not used
fb	3400	fn	
bs	4		
tpwr	63		
pw	7.1	werr	
dl	4.000	wexp	
tof	0	wbs	
nt	8	wnt	
ct	8		
alock	n		
gain	not used		
il	n		
in	n		
dp	y		
sd	18.6		
wd	3034.0		
vs	5.1		
sc	0		
mc	250		
hzmm	12.14		
is	1871.03		
rfl	3902.2		
rfp	2896.2		
th	20		
ins	3.000		
na			
ph			



Synthesis of **2.143**



To 76.4 mg (0.374 mmol) of **2.142** in a 20 mL scintillation vial with magnetic stir-bar was added 1.9 mL of a 1:1 mixture of THF and H₂O. Next, 142 mg (0.561 mmol) of I₂ was added, followed by 62.1 mg (0.449 mmol) of K₂CO₃. 9.1 mg (0.0748 mmol) of 4-(dimethylamino)pyridine (DMAP) was added, and the vial was purged with N₂, capped, and allowed to stir at ambient temperature while monitoring the reaction by TLC. After 2 h, starting material had been consumed, so the mixture was transferred to a separatory funnel with EtOAc and washed with saturated aqueous Na₂S₂O₃. The organic layer was washed with 11 mL of 0.1 M HCl, and then, dried with anhydrous Na₂SO₄. Volatile material was removed under reduced pressure, and 112 mg (0.339 mmol, 91%) of **2.143** was isolated, as a white solid, after purification by silica gel chromatography (hexanes:EtOAc).



(S)-5-(5-Fluoro-2-methylphenyl)-2-iodocyclohex-2-enone (2.143). Mp 100-106 °C. *R*_f = 0.23 (SiO₂, 10:1 hexanes:EtOAc); IR (CH₂Cl₂ solution): 2930 (m), 1684 (s), 1615 (m), 1590 (m), 1495 (s), 1318 (m), 1250 (m) cm⁻¹; ¹H NMR: δ 7.79 (1H, t, *J* = 4.4 Hz), 7.11 (1H, dd, *J* = 8.4 Hz, *J* = 6.0 Hz), 6.89 (1H, dd, *J* = 10 Hz, *J* = 2.8 Hz), 6.83 (1H, dt, *J* = 8.4 Hz, *J* = 2.8 Hz), 3.59 (1H, m), 2.86 (1H, dd, *J* = 16 Hz, *J* = 3.6 Hz), 2.72 (1H, dd, *J* = 16 Hz, *J* = 14 Hz) 2.50-2.64 (2H, m), 2.26 (3H, s); ¹³C NMR: δ 191.4, 161.4 (d, ¹*J*_{CF} =

242 Hz), 157.8, 141.8 (d, $^3J_{\text{CF}} = 6.2$ Hz), 132.1 (d, $^3J_{\text{CF}} = 7.8$ Hz), 130.7, 113.6 (d, $^2J_{\text{CF}} = 21$ Hz), 112.2 (d, $^2J_{\text{CF}} = 22$ Hz), 103.4, 43.09, 36.89, 36.47, 18.56; ^{19}F NMR: δ 95.43 (m). LRMS (APPI) Calcd for $\text{C}_{13}\text{H}_{12}\text{FIO}$ ($\text{M} + \text{H}$) $^+$: 331.0, Found ($\text{M} + \text{H}$) $^+$: 331.0.

J55-111colum

expi stdh



SAMPLE		DEC. & VT	
date	Mar 22 2007	dfrq	0
solvent	CDCl3	dp	30
111	exp	dot	0
ACQUISITION		nm	200
sfrq	400.029	nm	200
in	H1	nm	200
at	3.000	nm	200
np	35992	nm	200
sw	5998.8	nm	200
fb	3400	nm	200
bs	4	nm	200
tdwr	63	nm	200
pw	7.1	nm	200
dl	4.000	nm	200
tot	0	nm	200
nt	8	nm	200
ct	8	nm	200
atlock	n	nm	200
gain	not used	nm	200
11	FLAGS	nm	200
in	n	nm	200
dp	y	nm	200
DISPLAY		nm	200
sp	-21.5	nm	200
wp	3279.2	nm	200
vs	25	nm	200
sc	0	nm	200
wc	250	nm	200
hmm	13.12	nm	200
fs	972.22	nm	200
rfl	3903.0	nm	200
rtp	2896.2	nm	200
th	20	nm	200
ins	1.000	nm	200
nm	ph	nm	200




js5-111colum

exp2 std13c

SAMPLE DEC. & VT
date Mar 22 2007 dfrq 400.029
solvent CDCl3 in AL
f1 acq 45
f2 acq 45
f3 acq 45
ACQUISITION exp 0
sfrq 100.599 dm yy
tn C13 dmf yy
at 0.640 dmf 6617
np 32876 lb PROCESSING 1.00
sw 25683.4 wfile
fb 14200 proc
bs 4 fn
tpwr 57 not used
pw 8.7
dl 4.900 weff
tof 2271.7 wep
nt 1600 wds
ct 80 wnt
clock n
gain not used
flags n
il n
in n
dp y
DISPLAY
sp -412.8
wp 21599.7
vs 51
sc 0
wc 250
hzmm 86.40
is 500.00
f1 8932.9
f2 7745.2
th 100.000
ms no ph



Synthesis of **2.144**




To a mixture of 1.93 g (5.37 mmol) of **2.222**¹ and 1.29 g (10.7 mmol) of MgSO₄, in 18 mL of CH₂Cl₂ at 0 °C under N₂, was added 0.41 mL (5.4 mmol) of dimethylhydrazine, dropwise. After complete addition, the reaction was warmed to room temperature and allowed to stir for 2 h. The mixture was filtered through celite and concentrated under reduced pressure. Column chromatography (neutral alumina, hexanes:EtOAc) afforded 1.81 g (4.51 mmol, 84%) of **2.144** as a slight yellow oil.

(E)-1,1-Dimethyl-2-[3-(tributylstannyl)-but-2-enylidene]hydrazine (2.144). *R_f* = 0.26 (Al₂O₃, 40:1 hexanes:EtOAc); IR (neat): 2961 (s), 2930 (s), 2848 (s), 1860 (w), 1595 (m), 1551 (m), 1469 (m), 1356 (m), 1274 (m), 1129 (m) cm⁻¹; ¹H NMR: δ 7.31 (1H, d, *J* = 8.8 Hz), 6.30 (1H, dq, *J* = 8.8 Hz, *J* = 1.6 Hz, ³*J*_{SnH} = 67.2 Hz), 2.85 (6H, s), 2.03 (3H, d, *J* = 2 Hz, ³*J*_{SnH} = 45.2 Hz), 1.34-1.58 (6H, m), 1.28 (6H, h, *J* = 7.2 Hz) 0.78-1.01 (15H, m); ¹³C NMR: δ 144.1, 136.8, 131.7, 42.90, 29.15, 27.41, 19.96, 13.72, 9.21. LRMS (APPI) Calcd for C₁₈H₃₈N₂Sn (M + H)⁺: 403.2, Found (M + H)⁺: 403.2.

¹ Lipshutz, B. H.; Clososki, G. C.; Chrisman, W.; Chung, D. W.; Ball, D. B.; Howell, J. *Org. Lett.* **2005**, 7, 4561.

Synthesis of **2.145**



To an oven-dried 2-dram vial with magnetic stir-bar, in a dry-box under an Ar atmosphere, was weighed 1.7 mg (0.0046 mmol) of bis(benzonitrile)palladium dichloride, 2.8 mg (0.0091 mmol) of tri-*o*-tolylphosphine, and 30.0 mg (0.0909 mmol) of **2.143**. Next, 0.43 mL of dry, degassed *N,N*-dimethylformamide (DMF) was added, followed by 43.8 mg (0.109 mmol) of **2.144**. The vial was capped, sealed with electrical tape, removed from the dry-box, and heated at 65 °C for 4 h. After cooling to room temperature, the mixture was diluted with Et₂O and washed with 3 % aqueous KF (2x). The combined aqueous layers were back extracted with Et₂O (1x), and the combined organic layers were dried with anhydrous Na₂SO₄. After removal of volatile material under reduced pressure, 20.9 mg (0.0665 mmol, 73%) of **2.145** was isolated as a yellow solid after purification by silica gel chromatography (first column with hexanes:EtOAc, followed by a second column using CH₂Cl₂:Et₂O).

(S)-2-[(2E)-4-(2,2-Dimethylhydrazono)but-2-en-2-yl]-5-(5-fluoro-2-



methylphenyl)cyclohex-2-enone (2.145). Mp 119-122 °C. *R*_f = 0.23 (SiO₂, 2:1 hexanes:EtOAc) and 0.31 (SiO₂, 10:1 CH₂Cl₂:EtOAc); IR (CH₂Cl₂ solution): 2949 (m), 2867 (m), 2779 (w), 1677 (s), 1614 (m), 1551 (m), 1495 (m), 1463 (m), 1343 (m) cm⁻¹; ¹H NMR: δ 7.16 (1H, d, *J* = 9.4 Hz), 7.09 (1H, dd, *J* = 8.4 Hz, *J* = 6.4 Hz), 6.91 (1H, dd,

$J = 10$ Hz, $J = 4.0$ Hz), 6.87 (1H, dd, $J = 6.0$ Hz, $J = 2.8$ Hz), 6.80 (1H, dt, $J = 8.4$ Hz, $J = 2.8$ Hz), 6.25 (1H, d, $J = 9.4$ Hz), 3.48 (1H, m), 2.87 (6H, s), 2.40-2.67 (4H, m), 2.27 (3H, s), 1.99 (3H, s); ^{13}C NMR: δ 197.4, 161.4 (d, $^1J_{\text{CF}} = 242$ Hz), 143.7, 143.4, 143.0 (d, $^3J_{\text{CF}} = 6.3$ Hz), 133.8, 131.9, 131.8 (d, $^3J_{\text{CF}} = 9.3$ Hz), 130.7, 127.4, 113.2 (d, $^2J_{\text{CF}} = 20$ Hz), 112.2 (d, $^2J_{\text{CF}} = 22$ Hz), 45.18, 42.73, 42.71, 36.90, 33.04, 18.53, 16.58; ^{19}F NMR: δ 95.02 (m). LRMS (ESI+) Calcd for $\text{C}_{19}\text{H}_{23}\text{FN}_2\text{O}$ (M) $^+$: 314.2, Found (M) $^+$: 314.9.

js5-152column

exp2 std1h

SAMPLE DEC. & VT 0
date Mar 24 2007 dfrq 0
solvent CDCl3 dn 30
file /export/home/~dpwr 0
jpm/js5-152col- dot 0
nm 6
nm 6
ACQUISITION 200
sfrq 400.029 dmf PROCESSING
tn H1
at 3.000 wfile ft
nd 35392 proc not used
sw 5898.8 fn
fb 3400
bs 4 weff
tpwr 63 wepd
pw 7.1 wbs
dl 4.000 wt
tof 0
nt 0
ct 0
alock n
gain not used
flags
ii n
in n
dp y
DISPLAY 586.5
SP 2412.0
V 7.0
WC 250
h2mm 9.65
is 2066.55
rfi 3901.7
rfp 2896.2
th 20
ins 3.000
ph



j55-15scolumn

expl std13c

SAMPLE DEC. & VT

date Mar 24 2007 dfrq 400.029

solvent CDCl3 dn H1

file /export/home/~dbwr 45

j55-15scolumn 0

ACQUISITION 100.599

sfreq 100.599 dmf 8617

in C13 lb PROCESSING 1.00

at 0.640 wf file

nd 32876 vfile

sw 25683.4 proc ft

fb 14200 fn not used

bs 4

tpwr 57 wert

pw 8.7 wexp

dl 4.000 wds

tof 2271.7 wnt

nt 1600

ct 120

atlock n

gain not used

flags n

il n

in n

dp y

DISPLAY

sp -615.0

wd 21295.6

vs 41

sc 0

vc 250

hzmm 85.18



is 500.00

rfi 8532.1


rfp 7745.2

tp 15

ins no ph 100.000




Synthesis of **2.146**




A solution of 24.8 mg (0.0789 mmol) of **2.145** in 1.5 mL of 1,2-dichlorobenzene (*o*-DCB) was heated at 195 °C for 5 h in a sealed tube. After cooling to room temperature, the solvent was removed under reduced pressure, and 9.0 mg (0.033 mmol, 42%) of **2.146** was isolated, as a white solid, after silica gel chromatography (one column in hexanes/EtOAc, followed by a second using CH₂Cl₂/EtOAc). Spectral data were consistent with the literature.⁴⁹ $[\alpha]_{\text{D}}^{20} = -4.8^{\circ}$ (*c* = 1.0, CCl₄). The enantiomeric purity of **2.146** was determined by comparison to authentic racemic material prepared by the known route.¹

¹ Miki, S.; Suide, H.; Tawada, H.; Iwano, N.; Aoki, I.; Adachi, M. Process for the Production of Optically Active Cyclic Enaminone Derivatives. Eur. Patent 1,229,019 A1, July, 8, 2002.


Chiral SFC (AD-H, Chiralpak, 150 bar, 50°C, flow = 5.0 mL/min, 5.0 % MeOH) analysis of **2.146**:



2.146



Racemic
2.146





2.146 + racemic
coinjection


js5-56 column2

exp2 std1h

SAMPLE DEC. 8 VT
 date Jan 19 2007 dfreq 289.908
 solvent CDCl3 dn H1
 file ACQUISITION exp dpr 30
 sfreq 289.909 dm dof 0
 tn H1 dam mm
 at 5.000 dmf C
 np 37612 dseq 200
 sw 37612 dres 1.0
 fd 2100 homoPROCESSING n
 ds 4.000
 power 59 wfile
 pw 7.0 proc ft
 d1 4.000 fn not used
 tof 494.9 math f
 nt 16 weff
 ct 8 wexp
 atlock n wds
 gain not used wnt
 f1 . FLAGS n
 in dn n
 ds v n
 hs DISPLAY mn
 sp 70.4
 wp 2525.5
 vs 120
 sc 0
 wc 250
 hzmm 10.250
 f1 107.13
 rfp 2086.4
 th 2171.3
 ins 20
 ph 1.000



Synthesis of **2.148**



To 8.8 mg (0.033 mmol) of **2.146**, in a 0.5 dram vial, was added 4.5 mg (0.041 mmol) of aminoguanidine·HCl, followed by 0.10 mL of anhydrous MeOH. The vial was capped, sealed with electrical tape, and heated at 65 °C for 30 min. After cooling to room temperature, 10 µL of concentrated HCl was added, and the vial was heated at 100 °C for 2.5 h. Volatile material was next removed under reduced pressure, after cooling to room temperature. The residue was diluted with 0.25 mL of anhydrous MeOH, and 35.7 mg (0.66 mmol) of NaOMe was then added. This mixture was allowed to stir for 20 min, and water was then added, followed by extraction with EtOAc (10x). The combined organic layers were dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. This gave 13.2 mg of material, which was diluted with 0.1 mL of EtOH and 6.6 mg (4.5 µl, 0.069 mmol) of MeSO₃H was subsequently added. Upon standing overnight in the freezer, a solid precipitate formed that was subsequently collected by filtration. The solid was washed with EtOH that had been cooled to -78 °C (dry ice/acetone) followed by Et₂O that had been cooled to -78 °C (dry ice/acetone). The solid was finally collected by rinsing the material through the filter funnel using MeOH, followed by removal of volatile material under reduced pressure. This afforded 2.0 mg (0.0040 mmol, 12%) of slightly impure **2.148**. Its structure was verified by comparison of the ¹H

NMR spectrum with the reported literature data.⁴⁹ Further purification by crystallization or chromatography on alumina was unsuccessful.

SAMPLE		APR 5 2007		DEC. & VT	
date	Apr 5 2007	dfrq	499.766		
solvent	H ₂ O	dn	11		
title	/export/home/~ jpm/j35-163ctop1.7	dpr	30		
id	16	nm	30		
ACQUISITION		dm	C		
dfrq	499.767	dm	200		
tn	H1	dseq	200		
at	5.000	dres	1.0		
nm	70058	homo	1.0		
sw	7005.9	temp	28.0		
td	4000	DECT	0		
ps	5	dfrq2	0		
dpr	4.5	dprw2	1		
dl	4,000	dof2	0		
tof	487.0	dm2	0		
ct	128	dmw2	C		
nt	35	dm2	200		
clock	n	dseq2	1.0		
gain	not used	dres2	n		
FLOPS		homo2	PROCESSING		
in	n	wf	fit		
dn	y	proc	not used		
id	y	tr	f		
DISPLAY		math			
sp	-0.4	werr			
vc	5477.5	wexp			
vp	102	whs			
bs	25	whs			
h2am	21.91	wfit			
is	879.79				
rft1	506.8				
rftp	0				
th	13				
nm	1.000				
ph					

